



# Evaluation of Summer Marigold Hybrids for Agro Economic Traits under Prayagraj Agro-Climatic Condition

Pooja Patel <sup>a+++\*</sup>, Vijay Bahadur <sup>b#</sup>, Samir E. Topno <sup>b†</sup>,  
V. M. Prasad <sup>b‡</sup> and Anita Kerketta <sup>b†</sup>

<sup>a</sup> Department of Horticulture (Floriculture and Landscaping), Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh, 211007, India.

<sup>b</sup> Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh, 211007, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i173174

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101608>

Original Research Article

Received: 18/04/2023

Accepted: 21/06/2023

Published: 29/06/2023

## ABSTRACT

Evaluation of ten marigold varieties/ hybrids of African marigold, has been taken up at Vegetable farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (UP) during 2022-23 The experiment was laid out in Randomized Block Design (RBD) with three replications. The analysis of variance revealed

<sup>++</sup> M.Sc. Scholar;

<sup>#</sup> Associate Professor & Head;

<sup>†</sup> Assistant Professor;

<sup>‡</sup> Professor;

\*Corresponding author: E-mail: [poojapatel.soil99@gmail.com](mailto:poojapatel.soil99@gmail.com);

significant variation for most of the parameters. The result revealed that highest plant height (T10) (92.93 cm), number of branches (T7) (27), was noted in recorded maximum leaf area index (T7) (3.25 cm) and flower bud initiation (T10) (45). Earliest 50% flowering (T10) (57 days). No. of flower per plant highest in (T7) (78), highest yield per plot (873.34 g) and yield hectare-1 (4.37 t.) in (T7) Pusa Basanti.

**Keywords:** Summer; marigold varieties/ hybrids; climate.

## 1. INTRODUCTION

African Marigold (*Tagetes erecta* L.) belonging to family Asteraceae is the most commonly grown loose flower and is extensively used in religious and social functions. It is gaining popularity on account of its easy culture, wide adaptability, and increasing demand in the Asian subcontinent. Marigold plants originate from Central America, probably in Mexico. Today, they are naturalised in the tropics and sub-tropics of the Old and New worlds. They are cultivated in India and Pakistan as a medicinal, flavouring, dye and ornamental plant. Marigold flowers are available year round with a peak season in the summer and fall months. Marigold is an annual flower and requires full sun to bloom. The blooms may be single or double colored and can be varying hues of yellow, orange, red, and maroon. In Sometimes, the whole plant is used for decorations. It can be planted in beds for mass display, in mixed borders and can also be grown in pots. Different varieties of African marigold vary in plant height and spread, flower size, quality and yield. The flowers are large and globular in shape. Colour shades vary from light yellow to creamy yellow, bright yellow, cadmium orange, deep orange, sulphur yellow and white. "The total area under marigold cultivation was 68.33 thousand hectares with the production of 607.97 thousand MT (NHB 2017) in India and estimated production was 755.10 thousand MT according to NHB 2021-22 (1<sup>st</sup> advanced estimate). Seed production is a critical component in any crop improvement program, and finding good seed is a major challenge in marigold. The success of marigold production is determined by a number of management elements, one of which is nutrition. Nutrition plays a vital role in marigold development, flowering, and seed generation" [1].

"Among the various reasons behind low productivity of marigold, poor soil and nutrient management is a major cause. Therefore, nutrient management has prime importance for successful cultivation. But, the indiscriminate application of chemical fertilizers alters the soil fertility, leading to the pollution of soil and water

bodies. These are an important part of contemporary agriculture for maintaining production sustainability, yet they can be harmful to soil health over time. As a result, chemical fertilizers use must be restricted to a certain extent" [2]. "On the other hand total organic farming may be a desirable proposition for improving the quality of agricultural produce. It may not be possible to maintain the quality of the produce in commercial agriculture, where mostly the stress will be given mainly on yield. It is impossible to meet the nutrient requirement of the crops, exclusively through the organic farming" [3].

"After all the sustainability in agriculture system is a global issue. Under these circumstances, practice of INM is the better option for the improvement of physical (structure and water retention capacity), chemical (nutrients and cation exchange capacity) and biological (microflora and microfauna) properties. As a result, a promising cultivar and performing well in one region, may fail to perform well in another region of varying climatic conditions. The quest for selecting suitable high yielding variety/hybrid for the region leads to the requirement of collection and evaluation of available genotypes" [4]. INM is based on the idea of maintaining and maybe improving soil fertility in order to achieve long-term productivity [5].

"Under given agro-climatic conditions, it is important to study the performance of existing varieties available in market and also to identify the best genotype with desirable characteristics and yield which will fetch remunerative profit to farmers of the state. The ultimate yield and production of quality flowers, pigment contents in flower and resistance to biotic factors depend upon the selection of suitable cultivars for a particular locality" [4].

## 2. MATERIALS AND METHODS

The experiment was conducted at Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (U.P.) during

2022-2023. Prayagraj is situated at an elevation of 78 meters above sea level at 25.87° North latitude and 81.150 E longitudes. This region has a sub-tropical climate prevailing in the South-East part of U.P. with both the extremes in temperature, i.e., the winter and the summer. In cold winters, the temperature sometimes is as low as 32°F in December – January and very hot summer with temperature reaching up to 115°F in the months of May and June. During winter, frosts and during summer, hot scorching winds are also not uncommon. The average rainfall is around 1013.4 (cm) with maximum concentration during July to September months with occasional showers in winter.

The present investigation was carried out on summer, marigold varieties/ hybrids was used as a there were 10 treatments marigold varieties/ hybrids the treatment was common manures and fertilizer application replicated thrice in randomized block design. Cocopeat and vermicompost was used in nursery tray for preparation of marigold seedlings.

**Chart 1. List of treatments and treatment variety used for the study**

Treatments	Treatment variety	Variety
T <sub>1</sub> V <sub>1</sub>	Indus Orange Bunch	Hybrid
T <sub>2</sub> V <sub>2</sub>	Marigold Gulzafri Orange	Variety
T <sub>3</sub> V <sub>3</sub>	Indus Sweet Orange	Hybrid
T <sub>4</sub> V <sub>4</sub>	Pusa Narangi	Hybrid
T <sub>5</sub> V <sub>5</sub>	Double Orange African Marigold	Hybrid
T <sub>6</sub> V <sub>6</sub>	Marigold Yellow F2	Variety
T <sub>7</sub> V <sub>7</sub>	Pusa Basanti	Hybrid
T <sub>8</sub> V <sub>8</sub>	African Marigold Yellow Supreme	Hybrid
T <sub>9</sub> V <sub>9</sub>	Orange African Marigold	Hybrid
T <sub>10</sub> V <sub>10</sub>	Marigold Yellow F1	Hybrid

### 3. RESULTS AND DISCUSSION

#### 3.1 Plant Height (cm)

At 30 DAT, the plant height was reported highest (52.57cm) in the Double orange African marigold (T5) which was found *at par* with the Indus orange bunch (T1) (49.01 cm) (43.58). The plant height was reported minimum (27.98) in Pusanarangi (T4). Similarly, at 60 DAT, the plant height was recorded maximum (82.84 cm) in Indus sweet orange (T3) which was *at par* with (T10) Marigold yellow F1 (81.55) and (T1) Indus orange bunch (77.55) the plant height was reported minimum (47.44) in Pusanarangi (T4)

marigold yellow F1 (T10) observed maximum height (92.93 cm) At 90DAT which was *at par* with (T8) African marigold yellow supreme (88.48), followed by (T7) Pusabasanti (60.26 cm). Variation noted among the treatment might be due to the varietal characters responsible by a gene. As a genetically controlled factor, plant height varied among treatments of marigold. Similar results were also observed in different genotypes by kumar et al. [6], in a Chrysanthemum, Hedge et al. (2022), in Dahlia, Kumari et al. [7], Singh and Singh [8], Khanvilkar et al [9] in Marigold .

#### 3.2 Number of Branches per PLANT

At 30 DAT, the Number of branches was reported highest (11.47cm) in the Pusabasanti (T7) which was found *at par* with the Pusanarangi (T4) (9.26 cm) (9.11cm). the Number of branches was reported minimum (27.98) in Marigold yellow F2 (T6). At 60 DAT, the Number of branches was reported highest (17.47 cm) in the Pusabasanti (T7) which was found *at par* with the Pusanarangi (T4) (15.27 cm) (15.11 cm). the Number of branches was reported minimum (7.89) in African marigold yellow supreme (T8). At 90 DAT, the Number of branches was reported highest (22.47 cm) in the Pusabasanti (T7) which was found *at par* with the Pusanarangi (T4) (18.27 cm) (17.11 cm). the Number of branches was reported minimum (11.89) in Marigold yellow F1 (T6). The variation observed in different treatments for number of primary branch may be due to its genetic inheritance character [10] environmental condition also affected the genetic composition. These finding also followed with Singh and Singh [8], Rao et al. [11], in marigold, Munikrishnappa et al. (2013).

#### 3.3 Leaf Area Index and Days to Flower Bud Initiation

The records made known that the Leaf Area Index (3.25) was recorded in (T7) Pusabasanti which found to be *at par* with (T8) African marigold yellow supreme (2.83) whereas the lowest Leaf Area Index was reported in (T6) marigold yellow F2 (1.89). Illustrated the data relevant to Days to flower bud Initiation (T8) African marigold yellow supreme (37.25 Days) earliest days of flower bud Initiation were noticed in Late bud Initiation was (T10 marigold yellow F2 (45.18). The performance of genotype may vary with climatic condition and their genetic make up. These findings are common with Kumar [12]; in gerbera, vikas et al. (2015).

**Table 1. Plant height, number of branches per plant and leaf area index of marigold variety/hybrid at different time intervals**

Treatments	Treatments	Plant Height (cm)			Number of branches/ Plant			Leaf Area Index
		30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	
T <sub>1</sub>	Indus orange bunch	49.01	77.55	81.36	9	15	17	2.19
T <sub>2</sub>	Marigold gulzafri orange	35.85	61.7	79.33	5	9	13	2.33
T <sub>3</sub>	Indus sweet orange	48.53	82.84	86.72	8	13	16	2.1
T <sub>4</sub>	Pusanarangi	27.98	47.44	76.31	9	15	18	2.33
T <sub>5</sub>	Double orange African marigold	52.57	72.25	85.43	5	9	12	2.03
T <sub>6</sub>	Marigold yellow F2	34.99	70.5	86.36	3	9	13	1.89
T <sub>7</sub>	Pusabasanti	43.58	53.31	60.26	11	17	22	3.25
T <sub>8</sub>	African marigold yellow suprem	37.65	76.63	88.48	4	8	12	2.83
T <sub>9</sub>	Orange african marigold	35.64	61.59	68.82	8	12	14	2.15
T <sub>10</sub>	Marigold yellow F1	38.24	81.55	92.93	5	10	12	2.41
<b>S.Ed(±)</b>		<b>2.24</b>	<b>2.41</b>	<b>2.47</b>	<b>0.16</b>	<b>0.29</b>	<b>0.37</b>	<b>0.12</b>
<b>C.D. at 5%</b>		<b>6.66</b>	<b>7.15</b>	<b>7.34</b>	<b>0.49</b>	<b>0.86</b>	<b>1.1</b>	<b>0.36</b>

**Table 2. Days to flower bud initiation, days to 50% flowering, no. of flowers per plant and flower duration (days) of marigold variety/hybrid at different time intervals**

Treatments	Treatments	Days to flower bud initiation	Days to 50% Flowering	No. of flowers per plant	Flower duration (Days)
T <sub>1</sub>	Indus orange bunch	42	57	59	39
T <sub>2</sub>	Marigold gulzafri orange	37	48	44	26
T <sub>3</sub>	Indus sweet orange	40	52	50	28
T <sub>4</sub>	Pusanarangi	37	48	65	26
T <sub>5</sub>	Double orange African marigold	44	49	35	20
T <sub>6</sub>	Marigold yellow F2	39	48	35	24
T <sub>7</sub>	Pusabasanti	41	50	78	22
T <sub>8</sub>	African marigold yellow suprem	37	47	36	26
T <sub>9</sub>	Orange african marigold	42	49	47	21
T <sub>10</sub>	Marigold yellow F1	45	57	38	36
<b>S.Ed(±)</b>		<b>1.08</b>	<b>1.35</b>	<b>1.22</b>	<b>0.75</b>
<b>C.D. at 5%</b>		<b>3.21</b>	<b>4.01</b>	<b>3.61</b>	<b>2.24</b>

### 3.4 Days to 50% Flowering, No. of flowers per Plant and Flower Duration

The number of days required for the investigated variety to reach 50% flowering showed significant variances. According to data on the number of days needed for 50% flowering (T4) Pusanarangi (48.16 Days) experienced 50% flowering at the earliest possible time.(T10) Marigold yellow F1 (57.41 Days) was flowering late.Varied genotypes may require different flowering times depending on their genetic make-up (Singh and Singh, 2006). The results were consistent with those of Singh and Mishra [13] for marigold and those of Nair et al. (2003) for gerbera.

The records made known that the highest number of flowers plant.the records made known that the highest number of flowers per plant (78.34) was recorded in (T7) Pusabasanti which found to be *at par* with (T4) Pusanarangi (65.44) whereas the lowest number of flower was reported in (T5) Double orange African marigold (35.33).Similar finding have been reported by Singh and Sangama (2000) in China Aster, Singh et al. [14], Singh and Mishra [13], Sunitha et al. (2007), Narsude et al.(2010), Beniwal and Dahiya [15] in marigold. The result related to flower duration was found to be significant.the data relevant to days to first flowering earliest days of first flowering were noticed in (T1) Indus orange bunch (39.31Days).

## 4. CONCLUSION

These findings concluded thatsummer marigold variety/hybrid experiment considering the important vegetative, floral and yield characteristics, (T7) Pusabasanti is found suitable for commercial cultivation. To obtain the higher benefits in prayagraj (U.P.) under Allahabad agro climatic conditions. With better quality along with maximum yield of flower (T7) Pusabasanti followed by (T4) Pusanarangi may be recommended for commercial cultivation.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Swaroop Krishan Raju DVS, Singh K. Effect of nitrogen and phosphorus on growth, flowering and seed yield of African marigold var. Pusa Narangi Gainda

- (*Tagetes erecta* L.). Orissa Journal of Horticulture. 2007;35(2):15-20.
2. Gotmare PT, Damke MM, Gonge VS, Deshmukh S. Influence of integrated nutrient management on vegetative growth parameters of marigold (*Tagetes erecta* L.). Asian J. Hort. 2007;2(2): 33-36.
3. Upadhya AK, Singh R, Singh PKR, Sengar RK, Kumar M, Singh NV. Effect of integrated nutrient management on plant growth, flower yield of African marigold (*Tagetes erecta* L.).The Pharma Innovation Journal. 2022;11(5):2064-2069.
4. Borah K, Bora SS, Rahman SW. Effect of integrated nutrient management (INM) with special reference to floricultural crops: A review. International Journal of Chemical Studies. 2019;7(3):369-373.
5. Panda SC. Soil management and organic farming. Agrobios Publications (India). 2006;207-208.
6. Kumar PJ, Thakur, Mohit. Effect of conjoint application of nutrient sources and PGPR on soil microbiological properties of cauliflower. Journal of Pharmacognosy and Phytochemistry. 2017;6(3): 672-676.
7. Kumari A, Sharma BP, Gupta YC, Sharma U, Sharma S. Effect of biofertilizers applications on growth and flowering of African marigold cv. Pusa Narangi Gainda during different season of the year under mid-Hills conditions of Himachal Pradesh. International Journal of Current Microbiology and Applied Sciences. 2019; 8(9):234-240.
8. Singh D, Singh AK. Characterization of African marigold (*Tagetes erecta* Linn.). Journal of Ornamental Horticulture. 2006; 9(1):40-42.
9. Khanvilkar MH. Performance of African marigold (*Tagetes erecta*) in north Konkan coastal zone of Maharashtra.Journal of Maharashtra Agriculture University. 2003; 28(3):333-334.
10. Narsude PB, Kadam AS, Patil VK. Studies on the growth and yield attributes of different African marigold (*Tagetes erecta* L.) genotypes under Marathwada condition. Asian J. Hort. 2010;5(2):284-286.
11. Rao CC, Goud PV, Reddy KM, Padmaja G. Screening of African marigold (*Tagetes erecta* L.) cultivars for flower yield and carotenoid pigments. Indian Journal of Horticulture. 2005;62(3):276-279.
12. Kumar S, Singh JP, Braj-mohan Nathiram, Rajbeer. Influence of integrated nutrient

- management on growth, flowering and yield parameters of African marigold (*Tagetes erecta* L.) cv. Pusa Basanti Gaiinda. The Asian Journal of Horticulture. 2013;8:118-121.
13. Singh D, Misra KK. Genetic variability in quantitative characters of marigold. Indian Journal Horticulture. 2008;65(2):187-192.
  14. Singh CV, Shankar MV, Kasana BS, Bobede A. Effect of integrated nutrient management on flowering characters of African marigold (*Tagetes erecta* L.).International Journal of Agriculture Sciences. 2016;8(4):1000-1002.
  15. Beniwal BS, Dahiya SS. Variability studies in marigold (*Tagetes* spp.). In: Abstracts of National Seminar on Sustainable Agriculture and Food Security: Challenges in Changing Climate, held at CCS Haryana Agricultural University, Hisar, Haryana. March 27-28, 2012;298.

---

© 2023 Patel et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/101608>