



# Response of Different Moisture Regimes and Nitrogen Sources on Soil Health, Growth and Yield Attributes of Wheat: A Comprehensive Review

Suneel Kumar <sup>a++</sup>, Anant Sharma <sup>b#</sup>, Santosh Pandey <sup>ct</sup>,  
Shantonu Paul <sup>d§</sup>, Hariom Mishra <sup>e‡</sup>, Amit Kesarwani <sup>ft</sup>,  
Sandeep Kumar <sup>gt</sup>, Ayushi Jha <sup>h</sup>, Shubham Singh <sup>i++\*</sup>  
and Himanshu Tiwari <sup>b^</sup>

<sup>a</sup> Department of Soil Science and Agricultural Chemistry, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut – 250110, U.P., India.

<sup>b</sup> Department of Agronomy, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut – 250110, U.P., India.

<sup>c</sup> Department of Agronomy, Institute of Agriculture Sciences, Bundelkhand University, Jhansi – 284128, U.P., India.

<sup>d</sup> Krishi Vigyan Kendra, Assam Agricultural University, Dibrugarh, India.

<sup>e</sup> Department of Agronomy, Krishi Vigyan Kendra, Basti, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya – 224229, U.P., India.

<sup>f</sup> Department of Agronomy, College of Agriculture, G.B. Pant University of Agriculture and Technology, Pantnagar, Uttarakhand, India.

<sup>g</sup> Department of Agriculture Science, Galgoties University Greater Noida, U.P., India.

<sup>h</sup> Department of Soil and Water Engineering, Punjab Agricultural University, Punjab, India.

<sup>i</sup> Department of Agricultural Extension Education, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut – 250110, U.P., India.

## **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## **Article Information**

DOI: 10.9734/IJPSS/2023/v35i203837

<sup>++</sup> Research Scholar;

<sup>#</sup> Master Scholar;

<sup>†</sup> Assistant Professor;

<sup>‡</sup> Subject Matter Specialist;

<sup>^</sup> Assistant Professor (Guest Faculty);

<sup>§</sup> Farm Manager;

<sup>\*</sup>Corresponding author: E-mail: ssagftp@gmail.com;

**Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/106459>

**Review Article**

**Received: 17/07/2023**

**Accepted: 19/09/2023**

**Published: 27/09/2023**

## ABSTRACT

Wheat (*Triticum aestivum* L.), a ancestor of the Poaceae family, is one of the most important cereal crops in the world. On a global scale, wheat is the main cereal that is cultivated and produced most widely. Irrigation has been essential to keep wheat a key food crop around the world. In response to climate change, it is also becoming more important. In response to better irrigation management, a key aspect in increasing water productivity and maintaining the viability of water resource use in agricultural production. At all growth stages of the crop, the wheat crop's growth and yield characteristics improved with rising levels of moisture regime. The performance of the yield component parts has been linked to higher nitrogen fertilization levels.

**Keywords:** *Wheat; irrigation scheduling; water; organic manure; soil health; yield.*

## 1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is a main cereal crop of India and it is second most common crop of the country. In terms of both production and consumption, it is also one of the main staple crops in many parts of the nation. In India it is the second-most significant grain crop and is crucial to the nation's food and nutritional security Kumar et al. [1,2].

The goal of irrigation should be to replenish the soil water content in the root zone to a point where the crop can completely satiate its evapotranspiration needs Kumar et al. [1,2]. In its early phases of growth, wheat need water for root development and germination. Reduced plant populations and therefore lower yields of grains are caused by insufficient soil moisture throughout the initial phases of plant growth. Irrigation should be used before seeding to reduce emergence issues caused by soil crusting and make sure the soil has enough water for germination Gupta et al. [3]. To produce wheat with the maximum yield attainable, water is required at all stages of plant development, from seed germination to plant maturation. A positive association is present between irrigation frequency and grain yield Kumar et al. [4]. The type of crop, growth stage, length of the growing season, plant population, topography of the soil, temperature, relative humidity, wind speed, and crop management practices like tillage,

fertilization, and weeding are all factors that affect a crop's need for water. Three approaches, namely the soil moisture depletion strategy, the climate approach (IW/CPE ratio), and the critical stages, can be used to schedule irrigation in wheat. The climatological method is one of them and is extremely beneficial and well-recognized by scientists and research workers around the world.

To maximize production from various wheat varieties, ideal irrigation facilities are crucial Kumar et al. [5]. Nitrogen is crucial for all of the plant's living tissues. All of a plant's essential processes are associated with protein, of which nitrogen is a critical component. Nitrogen is a constituent of proteins, enzymes, coenzymes, nucleic acids, phytochromes and chlorophyll. It is significant to the plant's biochemical operations. Consequently, it is unique of the nutrients that wheat crops need the most Kutman et al. [6]. Widespread use of better technologies, with fertilizer management particularly that of nitrogen and organic manure playing a crucial role can enable an increase in wheat output that is more environmentally friendly Kumar et al. [1,2]. Vermicompost, crop residues, and organic manures are needed in addition to inorganic fertilizers for increased production and healthy soil. By combining inorganic fertilizers with organic manures, crop wastes, and bio-fertilizers, one can improve soil health and accelerate up nutrient uptake Pandey et al. [7].

An optimal level of soil fertility and plant nutrient delivery is maintained by combining organic and inorganic manures, optimizing the benefits from all potential sources of plant nutrients in order to maintain the required productivity Kumar et al. [1,2].

## 2. RESPONSE OF MOISTURE REGIMES AND NITROGEN SOURCES ON SOIL HEALTH

Shaaban [8] Compared to untreated soil, the organic residue improved bulk density, total porosity, macro and micro pores, soil water retention, and soil hydraulic conductivity. It was discovered that the combination of poultry manure and sunflower residue had a greater impact than either poultry manure or sunflower residue alone. Another finding was reported by Agbede et al. [9] reported that Poultry manure considerably increased soil porosity and moisture content while reducing soil bulk density and temperature. The contents of soil organic matter, soil and leaf N, P, K, Ca and Mg were all significantly elevated by the manure. Similar finding observed by Akanni et al. [10].

Moisture Regimes and nitrogen sources effects on soil pH. The dilution of salts and leaching of ions outside the root zone may be the cause of the fall in soil pH that occurs with irrigation application based on the IW/CPE ratio. By creating organic acids during their decomposition, the usage of organic manures, such as poultry manure and FYM, has also been shown to contribute to lowering the pH of the soil to some extent. This may also account for the increased availability and mobility of nutrients, particularly micronutrients. Kumar et al. [1,2]. Similar results were noted by Jat et al. [11].

Kumar et al. [1,2] reported that that a slight improvement in organic carbon was obtained with increasing level of moisture regime up to 1.2 IW/CPE ration. It might be caused by the right soil moisture levels being maintained along with the breakdown of organic litters and discharge of solubilized plant nutrients.

## 3. RESPONSE OF MOISTURE REGIMES AND NITROGEN SOURCES ON GROWTH OF WHEAT CROP

Saren et al. [12] reported that wheat grown on sandy loam soil produced the highest plant height, dry matter accumulation, leaf area index, and crop growth rate after four irrigations

provided at the crown root initiation, tillering, blooming, and grain development stages. An increase in crop moisture availability may be responsible for the improvement in growth characteristics. These finding support the observation of Rathore *et al.*, (2006).

Shah et al. [13] showed that the 75:25 ratio of N from urea and FYM produced the highest biological yield, straw yield, and grain yield of wheat. The subsequent treatment that received N from the two sources in a 50:50 ratio produced the highest yield.

Channabasanagowda et al. [14] reported the impact of several organic manures on wheat quality, seed output, and growth. When compared to other treatments, the application of vermicompost at 3.8 t per ha and poultry manure at 2.45 t per ha significantly increased plant height, the number of leaves, the number of tillers at 90 DAS, the number of ear heads per metre square, seed weight and seed yield, and protein content.

Rehman et al. [15] said that, it can boost wheat yield components and biomass under rain-fed conditions to apply inorganic fertilizers (NPK) along with organic fertilizers in the form of farmyard manure (FYM). Spikes  $m^{-2}$ , grains spike $^{-1}$ , biological yield, and thousand grain weight were significantly affected by various levels of NPK and FYM either separately or in combination. At 80-60-30 kg NPK  $ha^{-1}$ , the highest spikes  $m^{-2}$  were noted. The highest spikes  $m^{-2}$ , grains spike $^{-1}$ , thousand grain weight, and biological yield are produced by farmyard manure at 45 t  $ha^{-1}$ .

Atikullah et al. [16] reported that the maximum dry matter content, crop growth rate, relative growth rate were found from I<sub>1</sub> (Irrigation at 20 days after sowing at crown root initiation stage) which was statistically identical with I<sub>2</sub> (Irrigation at 55 DAS at flowering stage) whereas minimum from I<sub>0</sub> (No irrigation). Plant height ,number of tiller, number of spike ,number of spikelets, spike length, filled grains, total grains , grains weight, grain yield, straw, biological and harvest index were correspondingly showed higher results as of growth characters.

Gupta et al. [3] reported that irrigation at 1.2 IW/CPE was significantly superior to other irrigation treatments in terms of plant height, number of shoots  $m^{-1}$ , and dry matter accumulation ( $gm^{-1}$ ), with the exception of

irrigation at 1.0 IW/CPE and five irrigations at the CRI, tillering, late jointing, flowering, and milking stages. The availability of sufficient nutrients may have contributed to the enhanced vegetative development of plants, which in turn led to the higher dry matter at higher fertility levels. Both Laghari et al. [17]; Kale et al. [18] reported results that were similar.

Pandey et al. [19] reported that the consisted of three sulphur levels (a) Control (b) 20 kg/ha (c) 40 kg/ha) and four irrigation levels (a) I<sub>1</sub> 0.6 IW/CPE ratio (b) I<sub>2</sub> 0.8 IW/CPE ratio (c) I<sub>3</sub> 1.0 IW/CPE ratio and (d) I<sub>4</sub> 1.2 IW/CPE ratio). At all growth phases, the PBW 502 variety greatly outperformed other cultivars in terms of plant height, number of shoots per square metre and leaf area index (1.2 IW/CPE ratio).

Singh et al. [20] reported that the irrigation levels has significant effects on wheat growth and characteristics, as well as grain and straw yields. In a comparison with two and one irrigation, four irrigations recorded the highest plant height, tiller count, dry matter accumulation at harvest stage, and leaf area index at 90 DAS, as well as greater grain and straw output. The ability of plants to grow taller at higher irrigation levels may result from the maintenance of numerous metabolic processes by the plants' steady water supply. The Similar result was find by Rahman et al. [21]; Brahma et al. (2007).

Kumar et al. [22] studies on the effect of FYM and urea combination on wheat quality, seed production, and growth. Plant height, Leaf area index at 90 DAS, and the number of shoots increased significantly after applying 75% of the prescribed dose of N through inorganic sources and 25% of the recommended dose of N through organic sources (FYM) ha<sup>-1</sup>.

Ullah et al. [23] reported on the number of tillers, plant height, spikelets per spike, seeds per spike, biological yield, 1000 grain weight, grain yield, and harvest index as well as other growth and yield data. All growth and yield indicators were dramatically raised by various nitrogen levels. The treatment with nitrogen applied at 203 kg ha<sup>-1</sup> produced the most tillers, the tallest plants, and the best biological yield; however, the treatment with nitrogen applied at 145 kg ha<sup>-1</sup> produced the highest grain yield, seeds per spike, and 1000 grain yield.

#### **4. RESPONSE OF MOISTURE REGIMES AND NITROGEN SOURCES ON YIELD ATTRIBUTES OF WHEAT CROP**

Prashar and Thaman [24] reported that the best grain yields were achieved with irrigation at 50 mm CPE, followed by irrigation at 40 mm CPE. As irrigation water usage increased, it became less efficient.

Ibrahim et al. [25] observed that in order to achieve maximum crop output, the quantities of different organic manures added with the inorganic fertilizers must be optimized. The determined changes in wheat's growth and yield metrics compared to controls that received only inorganic fertilizer. The impacts of organic amendments were usually favourable. In comparison to the control, the application of organic manures enhanced the wheat yield. In comparison to the control, the wheat plant's height, tiller count, spike length, straw yield, grain yield, and weight per 1000 grains were all significantly different. Similar results are in agreement with those obtained by Nawab et al. [26].

Shah et al. [27] reported that the plant height, spike length, grain per spike, and 1000-grain weight increased with the integrated use in various amounts. The treatments which applied 25% N from FYM, 25% N from city waste, and 50% N from mineral fertilizer produced the highest grain yields. The treatments that applied 25% N from FYM, 25% N from poultry manure, or city waste, and 50% N from mineral fertilizer produced the same results. These results are supported by Iqbal et al. [28]; Arif et al. [29].

Ali et al. [30] studies the impact of various nitrogen levels on the development and yield of wheat crops at four different nitrogen rates, namely 0, 80, 130, and 180 kg ha<sup>-1</sup>. With a seed rate of 100 kg ha<sup>-1</sup>, the sowing date occurred during the previous week. The results demonstrated that raising the nitrogen levels over the control considerably enhanced the number of tillers per unit<sup>-1</sup>, plant height, spike's length, number of grain spikes<sup>-1</sup>, 1000-grain weight, and grain yield.

Singh et al. [31] the findings showed that the diversity of nitrification inhibitors in co-fertilizer caused growth characteristics, such as the number of tillers and plant height, to dramatically increase. These inhibitors slowly delivered nitrogen to the plant while reducing nutrient losses, making more nitrogen available to the

plant. In comparison to treatment T<sub>2</sub>, which applied a full dose of NPK through urea, a single superphosphate, and muriate of potash, the combined application of organic manures and inorganic fertilizers boosted the dry matter accumulation, number of grains Spike<sup>1</sup>, grain yield, straw yield, and NPK uptake by wheat crop.

Brar et al. [32] reported that a maximum wheat grain yield of 1.25 IW: CPE, which was statistically comparable to 1.0 IW: CPE, was achieved when irrigation was applied. This may be attributed to an increase in the number of seeds as a result of better soil moisture and nutrient availability. Similar findings were reported by Tripathi and Singh [33]; Dhindwal et al. [34].

Yousaf et al. [35] reported on the impact of timing of irrigation and various nitrogen fertilization amounts on the agronomic performance of wheat. A total of 0, 80, 100, 120, and 150 kg ha<sup>-1</sup> of nitrogen were added to the soil, with irrigation occurring 15, 20, 25, and 30 days following germination. In comparison to the control and other treatments, nitrogen applied at 120 and 150 kg N ha<sup>-1</sup> and irrigation timing of 25 days after germination produced the highest plant height, most tillers, the greatest number of fertile tillers, and the highest yield.

Ahmad et al. [36] reported that adding organic manure with a high N level under frequent watering considerably boosted grain yield by more than 120% compared to control treatment. However, the number of tillers m<sup>-2</sup>, number of grains spike<sup>-1</sup>, and 1000-grain weight all increased significantly along with the increase in grain yield. Additionally, results demonstrated that applying organic manure and high N levels had a favourable impact on the harvest index. The positive effects of nitrogen on the characteristics of the yield components could be primarily responsible for the grain yield enhancement caused by the application of high N levels. In this respect, Ayoub et al. [37]; [38].

Shahid et al. [39] reported that the considerable increase in grain yield and yield components following the application of poultry manure. Each increase in nitrogen application rate resulted in an increase in grain yield, grain crude protein content, the number of fertile tillers, spike length, flag leaf area, fresh weight, dry weight, plant height, and the number of grains.

Sheoran et al. [40] reported that the use of organic manures along with N and P fertilizers considerably improved all yield characteristics, including plant height, number of tillers/m of row length, spike length, and number of grains/spikes. When FYM, chicken manure, and press mud were combined with the appropriate doses of N and P, higher grain yields of wheat and were found.

Deo et al. [41] reported that the growth characteristics such plant height, number of shoots per square metre and Leaf area index were shown to be significantly higher under the I<sub>1</sub> (critical stages) treatment. The I<sub>1</sub> (critical phases) and I<sub>4</sub> (1.0 IW: CPE ratio) treatments were shown to have higher total effective tillers m<sup>-2</sup>, length of spike, harvest index (%), and test weight over both years. The highest grain and straw yields under I<sub>1</sub> (critical phases) treatment were observed throughout the period of the two years. This finding was supported the finding of Maliwal et al. [42].

Verma et al. [43] reported that irrigation applied at 0.9 IW/CPE ratio (I<sub>2</sub>) recorded the maximum values of growth indices, matter accumulation per metre row length, leaf area index, leaf area duration, crop growth rate, relative growth rate, and net assimilation rate and yield - grain, straw, biological, and Harvest Index proved significantly superior over I<sub>1</sub>, I<sub>4</sub>, and I<sub>3</sub> with treatment I<sub>5</sub> being the exception. Finding support the result of Bastia and Rout [44].

Pal et al. [45] reported that the number of shoots, plant height, dry matter accumulation, and yield-attributing characters such as number of grains spike<sup>-1</sup>, number of spike m<sup>-2</sup>, length of spike, test weight, grain, and straw yield by crop were significantly higher at the CRI stage (I<sub>1</sub>), 0.6 IW/CPE ratio (I<sub>2</sub>), and 0.8 IW/CPE ratio (I<sub>3</sub>) than at the 1.0 IW/CPE ratio (I<sub>4</sub>) moisture regime. Similar result find by Karim [46] Brahma et al. [47-50].

## 5. CONCLUSION

The study's findings showed that the wheat crop produced considerably higher levels of plant height, dry matter, number of shoots, leaf area index, effective teller, grain production, straw yield, and harvest index when irrigation timing was appropriate. The application of both organic and inorganic nitrogen sources promotes wheat crop development and production.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Kumar S, Agrawal S, Jilani N, Kole P, Kaur G, Mishra A, Tiwari H. Effect of integrated nutrient management practices on growth and productivity of rice: A review. *The Pharma Innovation Journal*. 2023;12(5):2648-2662.
2. Kumar S, Kumar S, Kumar R, Pathak D, Kumar D, Kumar A, Singh S, Tiwari H. Assessment of physico-chemical properties of soil as influenced by different moisture regimes and nitrogen sources in wheat crop. *International Journal of Plant and Soil Science*. 2023;35(19):765-772.
3. Gupta AK, Nainwal RC, Singh D, Singh AK. Response of irrigation scheduling on growth and yield of late sown wheat. *Progressive Research-An International Journal*. 2016;11:5307-5310.
4. Kumar P, Pannu RK, Khokhar SK. Effect of organic sources of nutrition and irrigation levels on growth and yield of wheat (*Triticum aestivum* L.) *International Journal of Life Sciences Biotechnology and Pharma Research*. 2012;1(4):178-186.
5. Kumar R, Singh BN, Sharma JD, Kumar P, Mohan B. Effect of wheat varieties, moisture regimes and nutrient supply system on yield, quality and nutrient uptake of late sown wheat (*Triticum aestivum*). *International Journal of Chemical Studies*. 2018;6(4):72-74.
6. Kutman UB, Yildiz B, Cakmak I. Effect of nitrogen on uptake, remobilization and partitioning of zinc and iron throughout the development of durum wheat. *Plant and Soil*. 2011;342(1-2):149-164.
7. Pandey PR, Singh SP, Dhyani BP, Kumar Y, Singh A, Kumar A, Kumar S. Impact of different nutrient management practices on the nutrient dynamics of wheat crop in Western Uttar Pradesh, India. *International Journal of Plant & Soil Science*. 2023;35(19):560-571.
8. Shaaban SM. Effect of organic and inorganic nitrogen fertilizer on wheat plant under water regime. *Journal of Applied Sciences Research*. 2006;2(10):650-656.
9. Agbede TM, Ojeniyi SO, Adeyemo AJ. Effect of Poultry Manure on Soil Physical and Chemical Properties, Growth and Grain Yield of Sorghum in Southwest, Nigeria. *American-Eurasian Journal of Sustainable Agriculture*. 2008;2(1):72-77.
10. Akanni DI. Response of nutrient composition and yield components of tomato (*Lycopersicon esculentum* Mill) to livestock manure. Ph.D. Thesis, Department of Crop, Soil and Pest Management, Federal university of Technology, Akure. 2005;120.
11. Jat RA, Wani SP, Sahrawat KL, Singh P, Dhaka PL. Fertigation in vegetable crops for higher productivity and resource use efficiency. *Indian Journal of Fertilizers*. 2011;7(3):22-37.
12. Saren BK, Dey S, Mandal D. Effect of irrigation and sulphur on yield attributes, productivity, consumptive use, and consumptive use efficiency of wheat (*Triticum aestivum*). *Indian J. Agri. Sci*. 2004;74(5):257-261.
13. Shah Z, Ahmad IM. Effect of integrated use of farm yard manure and urea on yield and nitrogen uptake of wheat. *journal of agricultural and biological science*. 2006;1(1):60-65.
14. Channabasanagowda, Patil NKB, Patil BN, Awaknavar JS, Ninganur BT, Hunje R. Effect of Organic Manures on Growth, Seed Yield and Quality of Wheat, Karnataka J. Agric. Sci. 2007;21(3):366-368.
15. Rehman S, Khalil SK, Rehman A, Saljoqi AUR. Organic and inorganic fertilizers increase wheat yield components and biomass under rainfed condition. *Sarhad J. Agric*. 2008;24(1):11-20.
16. Atikullah MN, Sikder RK, Asif MI, Mehraj H, Uddin AFMJ. Effect of Irrigation Levels on Growth, Yield Attributes and Yield of Wheat, *Journal of Bioscience and Agriculture Research*. 2014;02(02):83-89.
17. Laghari GM, Oad FC, Gandahi STAW, Siddiqui MH, Jagirani AW, Oad SM. Growth, yield and nutrient uptake of various wheat cultivars under different fertilizer regimes. *J. of Agri*. 2010;26(4):489-497.
18. Kale ST, Kadam SR, Gokhle DN, Waghmare PK. Response of wheat varieties to different levels of fertilizer on growth and yield under late sown condition. *Int. J. of Agri. Sci*. 2015;11(1):77-80.
19. Pandey N, Kumar S, Singh G, Singh RA. Effect of sulphur nutrient and moisture regimes on growth and productivity of

- wheat (*Triticum aestivum* L.) varieties international. Journal of Current Microbiology and Applied Science. 2017;6(9):3623-3628.
20. Singh SV, Raghuvanshi N, Sharma A, Singh RA. Effect of Nitrogen Management Practices on Growth and Yield of Late Sown Wheat (*Triticum Aestivum* L.) Varieties. The Bioscan. 2017;12(3):1813-1817.
  21. Rahman MA, Karim AJMS, Haque MM, Eqashira K. Effect of irrigation and nitrogen fertilization on plant growth and root characteristics of wheat on a clay terrace soil of Bangladesh. Journal of the Faculty of Agriculture. 2006;45(1):301-308.
  22. Kumar V, Raghuvanshi N, Singh RA. Effect of different combination of FYM and urea on growth and yield of wheat (*Triticum aestivum* L.). Bulletin of Environment, Pharmacology and Life Sciences. 2017;6(2):395-398.
  23. Ullah I, Ali N, Durrani S, Shabaz MA, Hafeez A, Ameer H, Ishfaq M, Fayyaz MR, Rehman A, Waheed A. Effect of Different Nitrogen Levels on Growth, Yield and Yield Contributing Attributes of Wheat. International Journal of Scientific & Engineering Research. 2018;9:595-602
  24. Prashar A, Thaman S. Studies on irrigation requirements of late sown wheat (*Triticum aestivum* L.) Progressive Agri. 2005;5(1/2):10-12.
  25. Ibrahim M, Hassan AU, Iqbal M, Valeem EE. Response of wheat growth and yield to various levels of compost and organic manure. Pak. J. Bot. 2008;40(5):2135-2141.
  26. Nawab K, Shah P, Arif M, Ullah A, Khan MA, Mateen A, Ali K. Effect of cropping patterns, farm yard manure, K and ZN on wheat growth and grain yield. Sarhad J. Agric. 2011;27(3):234-239.
  27. Shah SA, Shah SM, Mohammad W, Shafi M, Nawaz H, Shehzadi S, Amir M. Effect of Integrated Use of Organic and Inorganic Nitrogen Sources on Wheat Yield. Sarhad.J.Agric. 2010;26:559-564.
  28. Iqbal AS, Abbasi MK, Rasool G. Integrated plant nutrition system (IPNS) in wheat under rainfed condition of Rawalkot Azad Jammu and Kashmir Pak. J. Soil Sci. 2002;21:79-86.
  29. Arif M, Ali S, Khan A, Jan T, Akbar M. Influence of farm yard manure application on various wheat cultivars. Sarhad J. Agric. 2006;22(1):27-29.
  30. Ali A, Ahmad A, Syed WH, Khaliq T, Asif M, Aziz M, Mubeen M. Effects of Nitrogen on Growth and Yield Components of Wheat. Sci.Int. (Lahore). 2011;23(4):331-332.
  31. Singh CM, Sharma PK, Kishore P, Mishra PK, Singh AP, Verma R, Raha P. Impact of integrated nutrient management on growth, yield and nutrient uptake by wheat (*Triticum aestivum* L.). Asian Journal of Agricultural Research. 2011;5:76-82.
  32. Brar AS, Mahal SS, Deol JS, Buttar GS. Productivity, economics and energetics of wheat under different methods of crop establishment and irrigation schedules. Annual Agricultural Research New Series. 2012;33(3):174-177.
  33. Tripathi SC, Singh RP. Effect of Chiseling, green manuring and plating methods on productivity and profitability of rice- wheat cropping system. Indian Journal of Agronomy. 2007;52(4):279-282.
  34. Dhindwal As-Hooda IS, Malik RK, Kumar S. Water productivity of furrow-irrigated raing season pulses plorted on raised bed. Indian Journal of Agronomy. 2006;51(1):49-53.
  35. Yousaf M, Fahad S, Shah AN, Shaaban M, Khan MJ, Sabiel SAI, Osman KA. The effect of nitrogen application rates and timings of first irrigation on wheat growth and yield. International Journal of Agriculture Innovations and Research. 2014;2(4):645-65.
  36. Ahmed BEAM, Ahmed FE, Dessougi HI, Interactive Effect of Chicken Manure with Nitrogen Fertilizer and Watering Regimes on Yield and its Components of Bread Wheat. Scholars Journal of Agriculture and Veterinary Sciences. 2016;3(3):234-237.
  37. Ayoub MS, Guertin S, Lussier S, Smith D. Timing and levels of nitrogen fertilizer effect on spring wheat yield in Eastern Canada. Crop Sci. 1994;34:748-756.
  38. Jan MT, Khan S. Response of wheat yield components to type of nitrogen fertilizer their levels and application time. Pak. J. Biosci. 2000;3(8):1277-1280.
  39. Shahid M, Saleem MF, Khan HZ, Wahid MA, Sarwar M. Improving Wheat (*Triticum Aestivum* L.) Yield and Quality by Integration of Urea with Poultry Manure, Soil Environ. 2015;34(2):148-155.
  40. Sheoran S, Raj D, Antil RS, Mor VS, Dahiyal DS. Productivity, seed quality and nutrient use efficiency of wheat (*Triticum aestivum*) under organic, inorganic and

- integrated nutrient management practices after twenty years of fertilization. *Cereal Research Communications*. 2017;45(2):315–325
41. Deo K, Mishra SR, Singh AK, Mishra AN, Singh A, Kumar S. Determination of suitable irrigation schedule for optimum water use efficiency of wheat crop. *Journal of Medicinal Plants Studies*. 2017;5(3):343-347.
  42. Maliwal GL, Patel JK, Kaswala RR, Patel ML, Bhatnagar R, Patel JC. Scheduling of irrigation for wheat (*Triticum aestivum*) under restricted water supply in Narmada region. *Indian Journal of Agricultural Sciences*. 2000;70(2):90-92.
  43. Verma HP, Sharma OP, Yadav LR, Yadav SS, Shivran AC, Kumar R, Balwan. Growth indices and yield of wheat (*Triticum aestivum* L.) as influenced by irrigation scheduling and organic manures. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(1):908-912.
  44. Bastia DK, Rout AK. Response of wheat (*Triticum aestivum*) cultivars to levels of fertilizers under limited irrigation. *Indian journal of Agronomy*. 2001;46(4):670-673.
  45. Pal S, Kumar S, Gangwar HK, Singh A, Kumar P. Effect of scheduling irrigation based on, IW/CPE ratio on dry matter accumulation, yield attributes, yield and economics of wheat (*Triticum aestivum* L.) *Journal of Pharmacognosy and Phytochemistry*. 2020;9(4):1946-1949
  46. Karim M. Growth and yield components of wheat under water stress of different growth stage. *Bangladesh J. Agril. Res*. 2011;36(3):455-468.
  47. Brahma R, Janawade AD, Palled YB. Effect of irrigation schedules, mulch and antitranspirant on growth, yield and economics of wheat (cv. DWD-1006). *Karnataka Journal of Agricultural Sciences*. 2010;20(1).
  48. Gill MS, Kulwinderjit Singh, Singh K. Effect of irrigation regimes and rates of nitrogen on yield and quality of durum wheat. *Res. (Punjab Agri University, Ludhiana, India)*. 1999;36(3-4):180-186.
  49. Jat ML, Shivran AC, Puniya MM, Boori PK, Ola BL, Verma HP. Effect of drip irrigation scheduling on growth and seed production of fennel (*Foeniculum vulgare Mill.*) under semi-arid agro-climatic condition. *International J Seed Spices*. 2015;5(2):67-73.
  50. Maqsood M, Ali A, Aslam Z, Saeed M, Ahmad S. Effect of irrigation and nitrogen levels on grain yield and quality of wheat. *International Journal of Agriculture and Biology*. 2002;4(1):164-165.

© 2023 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/106459>