



Prevalence of Malaria in University of Uyo Hostel in Uyo, Akwa Ibom State, Nigeria

**Ikpeida, N. W. ^{a*}, Nnanna-Chigozie, E. P. ^b,
Udoewah, S. A. ^c, Ndubuka, C. E. ^d and Inyang, S. X. ^a**

^a *Department of Animal and Environmental Biology, University of Uyo, Akwa Ibom State, Nigeria.*

^b *Department of Zoology and Environmental Biology, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria.*

^c *Department of Nursing Sciences, University of Port Harcourt, Nigeria.*

^d *Department of Optometry, Abia State University, Uturu, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121466>

Original Research Article

Received: 09/06/2024

Accepted: 11/08/2024

Published: 14/08/2024

ABSTRACT

Malaria has so far been a life-threatening parasitic disease transmitted by the bite of infected female anopheles mosquitoes in Nigeria at large and among students of University of Uyo hostels as a case study. Despite the massive increase in public and private efforts, research, symposiums and conferences to curb malaria over the years, it remains one of the most salient global health concerns. This study was aimed to investigate the prevalence of malaria in University of Uyo hostel. Questionnaires were given to the participant for their consent and standard methods of blood analysis for malaria test were employed for this research. A total number of participant for

*Corresponding author: E-mail: ikpeidannamso@gmail.com;

Cite as: N. W., Ikpeida, Nnanna-Chigozie, E. P., Udoewah, S. A., Ndubuka, C. E., and Inyang, S. X. 2024. "Prevalence of Malaria in University of Uyo Hostel in Uyo, Akwa Ibom State, Nigeria". *Asian Journal of Medical Principles and Clinical Practice* 7 (2):340-50. <https://journalajmpcp.com/index.php/AJMPCP/article/view/242>.

this research were 105 students who resides in the school hostel. The result from this study revealed malaria disease to be 64%. Females recorded highest of 58.38% against male with 40.62%. Age group 23-26 recorded highest while lowest was recorded for 15-18. The percentages of students using and not using LLINs were 48.57% and 51.43% respectively. However, positive results recorded for both cases were 41.12% and 62.50% respectively. 96.19% Of the students were knowledgeable of the control of malaria with the use of LLINs. Conclusively, based on this research; malaria is still a threat to the University of Uyo hostel residents. Based on these results, it is therefore pertinent for Students to be properly and regularly sensitized on malaria, its prevalence, control, treatment and preventive measures.

Keywords: Prevalence; malaria; University of Uyo Hostel; Akwa Ibom State; Nigeria.

1. INTRODUCTION

“Malaria has so far been a life-threatening parasitic disease transmitted by the bite of female anopheles mosquitoes” [1]. “Despite the massive increase in public and private efforts over the years, malaria remains one of the most salient global health concerns” [2]. “About half of the world population is at risk of malaria infection, and around 250 million cases occur annually, leading to approximately 1 million deaths each year” [1]. “The majority of the burden of disease caused by malaria is borne by the populations living in the highly endemic areas of sub-Saharan Africa. Within these areas, the populations at highest risk are pregnant women and infants” [3]. “Malaria mortality in Nigeria accounted for about 30% of the world total, and its associated burden relates to approximately 60% of outpatient visits to health facilities” [2]. “It is the most common Protozoan parasitic disease in the tropical and subtropical regions of the world” [4]. “Over the years, a lot of efforts have gone into controlling malaria in Nigeria and other African countries, the most affected by the disease, but the problem has not shown any sign of abating” [5]. “The reasons for the unsuccessful efforts to eradicate malaria disease in Nigeria includes lack of political will and commitment, low awareness of the magnitude of malaria problem, poor health practices by individuals and communities and resistance to drugs” [6]. “Malaria is the most common occurring parasitic infectious disease in school hostels” [7]. In general, ignorance to controlling and preventing malaria is another major problem of Nigerians. Therefore, this study aimed to investigate the prevalence of malaria in University of Uyo hostel.

2. MATERIALS AND METHODOLOGY

2.1 Materials

“Microscopy is the main tool for laboratory diagnosis of malaria. The thick blood film and the

thin blood film methods are employed. Field’s stain A (A polychrome methylene blue, disodium hydrogen phosphate and potassium hydrogen phosphate) and Field’s stain B (Eosin, disodium hydrogen phosphate and potassium hydrogen phosphate). Thin film is carried out following the examination of thick film to identify the particular species of the Plasmodium responsible for the infection; it is stained with commercially prepared Leishman stain. The materials employed in the study included a Leitz light microscope, EDTA (ethylene diamine tetra acetic acid) bottles, methylated spirit (methanol), cotton wool, tourniquet, syringes (5 mL) and needles (21 G)” (Epidi et al., 2008).

2.2 The Study Area

This study was conducted at the hostel of the University of Uyo, Nigeria. Uyo is the capital city of Akwa Ibom State, Nigeria. It lies in the coastal zone of the tropical rainforest of Nigeria, within latitude 4°32' N and 5°33' N and longitude 7°25' E and 8°25' E. The area has two rainfall peaks in June and September. The relative humidity is about 77.20%. Based on reports of the last population census, Uyo has an estimated population of 222 841 [8].

2.3 Study Population and Period of Study

The population which served as models for this study comprised One hundred and five (105) healthy looking students (41 males and 64 females) residing in the Uniuyo hostel. This study was carried out for duration of three (3) weeks (May 28 –June 20). At least 5 samples and filled questionnaires were collected per day and were taken to the laboratory for tests.

2.4 Inclusion Criteria

The criterion for participation in this study was strictly to the residents of the University of Uyo Hostel.

2.5 Exclusion Criteria

Residents who had treated malaria at least two weeks before the commencement of the research.

2.6 Sample Collection

The method of sample collection employed was venepuncture technique (Okocha et al., 2005; Epidiet al., 2008). Soft tubing tourniquet was fastened to the upper arm of the patient to enable the index finger feel a suitable vein. The puncture site was then cleansed with methylated spirit (methanol) and venepuncture made with the aid of a 21G needle attached to a 5mL syringe. When sufficient blood had been collected, the tourniquet was released and the needle removed immediately while the blood was transferred into an EDTA bottle (Epidiet al., 2008).

2.7 Data Collection

A structured questionnaire was designed and administered to the participants to obtain relevant demographic information and clinical characteristics.

2.8 Administration of Questionnaires

Well structured and approved questionnaires were given to randomly consented students that reside in Uniuyo hostels to provide information on their age, level (class), sex and their usage of ITNs. Confidentiality was maintained as no details related to participant's identity were used.

2.9 Laboratory Analysis

The collected blood samples were analyzed within 1 to 2 h of collection. Thick and thin blood films were prepared according to the technique outlined by Cheesebrough [9] and described by Epidiet al., (2008). A drop of each blood sample was placed in the center of a greasefree clean glass slide. Thereafter, the reverse side of the slide was cleaned with cotton wool and kept for air-drying and staining with field's stain. The slide was held with the dried thick film side facing downward and dipped in field's stain A (eosin) for 5 secs. It was washed off gently in clean water and then dipped in field's stain B (methyl azure) for 5 s and washed again in clean water. The back of the slide was cleaned with cotton wool and kept in the draining rack to air-dry for

eventual examination under the microscope, using oil immersion at 100X magnification to observe for *Plasmodium* parasites. Presence of ring forms.

2.10 Identification

Positive specimens were identified on the basis of microscopy. Using standard [10], a trained laboratory technician at health centre interpreted the malaria blood slides. Prevalence of *Plasmodium* was calculated as the proportion of sampled persons with a positive result divided by the number of persons who provided blood samples. All point estimates were weighted, with empirically estimated standard errors used to account for prevalence.

2.11 Data Analysis

Data obtained from this research were subjected to descriptive analysis. Chi-square test was also employed to ascertain the pattern of distribution of infection among the pregnant women examined, taking age and occupation into consideration. The statistical software used was SPSS version 20.

3. RESULTS AND DISCUSSION

3.1 Results

A total number of students random consented and selected for this study were 105 students. All the participated students reside in the Uniuyo hostel and had not treated malaria two weeks earlier before the commencement of this research. Microscopy results from the Uniuyo Health Services Department, were I carried out the study identified 64 (61.0%) malaria infections among the 105 (100.0%) students who had provided a blood sample. Presence of ring forms of *Plasmodium* and trophozoites of *Plasmodium* indicates positive results. The overall result is shown in Table 1.

Table 2 shows the sex related prevalence of parasites among students as the participation ratio of male (M) to female (F) were 41 (39.05%) and 64 (60.95%) respectively. Out of 41 males who participated in this study, 26 (63.41%) were positive and 15 (36.59%) were negative of *Plasmodium*. 38 (59.48%) positive and 26 (40.62) negative result was revealed for the total female students 64 (61.0%) that participated in this study.

Table 1. Showing the overall prevalence of malaria among students of the Uniuyo residential hostel

S/N	Sex	Age	Status	Use of LLINs	Knowledge on the Use of LLINs to Control Malaria
1	M	16	+	N	N
2	F	20	-	Y	Y
3	F	19	+	Y	Y
4	F	20	+	N	Y
5	F	17	-	Y	Y
6	F	25	-	Y	Y
7	M	22	+	Y	Y
8	M	23	+	N	Y
9	F	19	-	N	Y
10	F	18	+	N	Y
11	M	20	+	Y	Y
12	F	23	+	N	Y
13	M	22	-	N	Y
14	M	22	+	Y	Y
15	F	26	-	N	Y
16	F	26	+	N	Y
17	F	24	+	Y	Y
18	F	24	+	Y	Y
19	F	25	-	N	Y
20	F	22	+	Y	Y
21	M	22	+	N	Y
22	M	26	-	N	Y
23	F	26	+	N	Y
24	F	24	-	Y	Y
25	M	24	-	Y	Y
26	M	25	+	Y	Y
27	F	18	+	N	Y
28	F	19	+	Y	Y
29	F	20	+	Y	Y
30	F	26	+	N	Y
31	F	21	-	Y	Y
32	F	20	+	Y	Y
33	M	23	+	N	Y
34	M	24	-	Y	Y
35	F	23	+	N	Y
36	F	19	-	N	Y
37	M	15	+	Y	N
38	F	18	-	Y	Y
39	M	20	+	Y	Y
40	F	23	-	N	Y
41	F	24	+	Y	Y
42	F	23	+	N	Y
43	F	22	+	N	Y
44	F	21	-	Y	Y
45	M	19	+	N	Y
46	M	25	+	N	Y
47	F	23	+	Y	Y
48	M	15	-	N	Y
49	F	16	+	Y	Y
50	F	18	+	Y	Y
51	F	18	+	N	Y
52	M	17	+	Y	Y

S/N	Sex	Age	Status	Use of LLINs	Knowledge on the Use of LLINs to Control Malaria
53	F	16	-	Y	Y
54	F	15	+	Y	Y
55	M	20	+	N	Y
56	F	23	+	N	Y
57	M	22	+	N	Y
58	F	22	+	N	Y
59	F	26	+	Y	Y
60	F	26	-	N	Y
61	M	24	+	Y	Y
62	F	24	+	N	Y
63	F	25	+	Y	Y
64	M	18	-	Y	Y
65	F	19	-	N	Y
66	F	20	-	Y	Y
67	F	26	+	Y	Y
68	M	21	+	N	Y
69	F	20	-	Y	Y
70	F	23	-	Y	Y
71	M	24	+	N	Y
72	M	23	+	Y	Y
73	F	23	-	Y	Y
74	M	24	-	Y	Y
75	M	25	-	N	Y
76	M	18	+	Y	Y
77	F	19	+	N	Y
78	F	17	+	N	Y
79	M	16	+	N	N
80	F	16	+	N	Y
81	M	17	-	Y	N
82	F	18	-	Y	Y
83	M	17	-	Y	Y
84	F	18	+	N	Y
85	M	19	-	N	Y
86	F	20	+	N	Y
87	F	26	-	Y	Y
88	M	17	-	N	Y
89	F	16	+	N	Y
90	M	15	+	N	N
91	M	20	-	Y	Y
92	F	23	+	N	Y
93	M	22	-	Y	Y
94	F	22	-	Y	Y
95	M	26	+	N	Y
96	F	26	-	Y	Y
97	F	24	-	Y	Y
98	M	24	+	N	Y
99	M	25	+	N	Y
100	F	20	+	N	Y
101	F	21	-	Y	Y
102	F	25	-	Y	Y
103	M	19	+	N	Y
104	F	18	+	N	Y
105	M	20	+	N	Y

Keys: "F" = Female, "M" = Male, "Y" = Yes, "N" = No, "LLINs" = Long lasting insecticide nets, "+" = Positive, "-" = Negative

Table 3 shows age related prevalence of Plasmodium among the students. It revealed that the age group 23-26 [43 (40.95%)] which had highest number of participant also had the highest prevalence as positive results of 26 (60.47%) and negative results of 15 (36.59%) students were recorded. 15-18 {26 (24.77%)} age group had the lowest in both positive and negative results of 15 (57.70%) and 11 (43.30%) respectively.

Table 4 indicates the sex related prevalence of malaria infection among students using Long Lasting Nets (LLINs) of which 7 (41.12%) and 10 (58.88%) of the male subjects tested positive and negative respectively. Out of 34 (56.63%) female subjects that were using LLINs, 15 (44.11%) and

19 (55.88%) students tested positive and negative respectively.

Table 5 shows the sex related prevalence of malaria infection among students not using Long lasting nets (ITN). The result reveals that, out of 24 (58.55%) male students that were not using the LLINs, 15 (62.50%) were found positive and 9 (37.50%) were found negative and 21 (70.00%) and 9 (30.0%) of female students that did not use the LLINs were tested positive and negative respectively.

Table 6 reveals their level of knowledge on the use of LLINs, though did not justify their understanding on its effectiveness in the control of malaria.

Table 2. Sex related prevalence of Malaria parasites among students

Sex	No. of students (%)	Positive (+) (%)	Negative (-) (%)
M	41 (39.05)	26 (40.62)	15 (36.59)
F	64 (60.95)	38 (59.38)	26 (63.41)
Total	105	64 (60.95)	41 (39.05)

Table 3. Age related prevalence of Malaria parasites among students

Age group	No. of students (%)	Positive (+) (%)	Negative (-) (%)
15-18	26 (24.77)	15 (57.70)	11 (43.30)
19-22	36 (34.28)	22 (61.11)	14 (38.89)
23-26	43 (40.95)	26 (60.47)	17 (39.53)

Table 4. Sex related prevalence of malaria infection among students using insecticide treated nets (LLINs)

Sex	No. of students using (LLINs)	Positive (+) (%)	Negative (-) (%)
Male	17 (41.45)	7 (41.12)	10 (58.88)
Female	34 (56.63)	15 (44.11)	19 (55.88)
Total	51 (48.57)	22 (20.95)	29 (27.61)

Table 5. Sex related prevalence of malaria infection among students not using Long Lasting Insecticide Nets (LLINs)

Sex	No. of students NOT using (ITN)	Positive (+) (%)	Negative (-) (%)
Male	24 (58.55)	15 (62.50)	9 (37.50)
Female	30 (46.87)	21 (70.00)	9 (30.00)
Total	54 (51.43)	36 (34.28)	18 (17.14)

Table 6. Knowledge of the use of LLINs in the control of mosquitoes

Sex	Yes	No
Male	41 (39.05)	37 (62.50)
Female	64 (60.95)	64 (100)
Total	105	101 (96.19)

3.2 Discussion

Malaria occurs almost exclusively in the tropics and subtropics WHO, [11] and approximately, 40% of the world's population, mostly those living in the world's poorest countries, are at risk of malaria [12]. "Every year, more than 500 million people become severely ill with malaria in most cases and deaths occur within sub-Saharan Africa" [13].

In this study, the presence of ring forms of *Plasmodium* and Trophozoites of *Plasmodium* indicate positive results and its absence indicate negative results. Results from this study showed the overall prevalence of *Plasmodium* to be 61.0% in the University of Uyo hostel. This differs from the overall prevalence of 81.5% reported in a study by Okonko et al., (2005) among participants in Abeokuta, Ogun State, Nigeria, 51.5% reported by Epiidi et al., (2008) among blood donors in Abakaliki, Ebonyi State, Nigeria and 80.6% prevalence amongst students of a southwest Nigerian Federal University [7].

Malaria can affect all the ages, human groups and both male and female sexes. Studies have also shown seasonal variations in the rate of infections and differences in the types of malaria parasite in sex depending on the geographical conditions [14]. This study shows that malaria parasite was higher in females 38 (59.38%) than in male 26 (40.62%) of the total number [64 (60.95%)] of infected students. This is in agreement with the findings Okonko et al., [15] in the Southwestern, Nigeria, but disagrees with Autino et al. [16] and Abubakar et al., [17] who reported infection rate to be higher among adult male students [176 (73.64%)] than female students [63 (29.36%)] in Pakistan and Bauchi State University Gadau, Bauchi State, Nigeria which recorded 283 (70.7%) and 177 (29.3%) among male and female students respectively.

The age group 15-18 shows the lowest prevalence of malaria parasites among students. This result is attributed to their levels; students of ages 15-18 were mostly the new students (level 100) that newly resumed and moved in to the hostel with LLINs from their homes as a requirement for residents, while age group 23-26 showed the highest prevalence of malaria. Some students in this age group reported that they stopped using their LLINs because they were worn out, while others gave theirs to younger students. This, agrees with Olusegun-Joseph et al., [18] that reported that age group 15-20 had

80 (20%) and 21-25 had 140 (35%) prevalence of malaria in University of Lagos, Nigeria but disagrees with Mgbemena et al., [19] that reported that; out of 40 students, ages 16-22 had higher prevalence of 28 (70%) over ages 23-29 that recorded 12 (30%).

In the age comparison of malaria parasites prevalence, the study reveals that the students between ages 19-22 (61.11%) is almost comparable to that of the students between the ages of 23 – 26 (60.47%) and also comparable to a study which had 70% for students of FUTO, Imo State, Nigeria of ages 16 – 22 [19].

This study also reveals the prevalence of malaria parasites among students that uses and does not use LLINs amidst this research. Out of 41 male students, 17 (41.45%) were using LLINs and 7 (41.12%) recorded positive for malaria. 24 male students were not using the LLINs and 15 (62.50%) were recorded positive for malaria. The ratio of female students (64) that did and did not use the LLINs was 34 (56.63%) and 30 (46.87%) respectively. Among the 34 female students that used the LLINs, 15 (44.11%) recorded positive and 19 (55.88%) recorded negative. The 30 female students that did not use the LLINs recorded 21 (70.00%) and 9 (30.00%). This was as a result of their inconsistency in using the LLINs, mostly in the day. Yhdego et al., [20] reported that out of 76 people, the prevalence of malaria with the use of LLINs in Maygaba town, Ethiopia was 24 (31.58%) positive results. This research also reveals that a good number of them are knowledgeable of the use of LLINs, though does not reflect their understanding on the importance of its usage as it is one of the cheapest measure to fight malaria by avoiding direct contact to the body especially while at night as they are nocturnal in nature.

The transmission of malaria parasites in this study could be influenced by socio-economic and cultural factors when considering the hostel location and its structures which possibly play a vital role in influencing susceptibility to infection. The prevalence of the malaria among these subjects could also be attributed to the effect of climatic features on vector breeding and transmission as wet season usually promote mosquito breeding, hence the justification, because this study was carried out during the rainy season (May-June).

The fact that females were more infected than males in this study could be due to the sample

size difference. Also, female students in the southern states tends to expose their bodies with short skirts/shorts and even sleeveless shirt thus the contradiction in Abubakar et al., [17] result that reported higher prevalence in female student in Bauchi (a Muslim state) were generally, their females wear their hijabs that covers almost all part of their body and even at home, therefore disallowing direct contact with mosquitoes. This is also supported by Garley et al., [6] that reported the general gender ratio of LLINs usage is 57.2% and 48.8% in female and males respectfully. However, male students generally, prefer to sleep outside LLINs, hence the chances of being bitten by the mosquito vector increases. Males in some settings use health care services less than females because they pay less priorities to their health, making them hesitant to go long distances to health centers even when malaria is suspected Murray et al., [21], thus the higher prevalence in male [20,7].

From this study, as also supported by Natama et al., [22] malaria transmission is not homogenous through an endemic area but spotty and depends on two primary factors; location of the breeding sites and clustering of human habitations where people are serving as reservoirs of malaria parasites. It could also be as a result of poor or inadequate drainage system. Also in the school hostels, there is overpopulation of over 10 students per room thus making use of mosquito net as preventive measure cumbersome, which is not the standard of room allocation in Univeristy of Uyo hostel.

Additionally, a substantial number (51.43%) of the respondents did not possess LLINs due to two main reasons; some of theirs were worn-out while other gave theirs out to junior colleagues. This result indicated that there was an imbalanced and unsustainable possession of the use of LLINs of the students in the study area. Therefore, timely replacements of the worn out LLINs via malaria campaigns are important for effective prevention and control of malaria in this study area. On other hand, the effectiveness of LLINs depends on regular and consistent use, and people's perception towards its utilization [23,24]. In our study 101 (96.19%) students had the knowledge towards the use of LLINs. Nevertheless, only 51 (48.57%) of them were using LLINs. This finding is higher than similar studies conducted in Raya Azebo and Arsi zone, Ethiopia Tesfay et al., 2019; Ragasa et al., [25] but lower than the reports of other settings in Ethiopia [26]. The rest of the respondents did not

use LLINs while sleeping; this might be due to a negative attitude towards LLINs because of various reasons as mentioned by Taremwa et al [27] Furthermore, the use of LLINs were associated with several factors such as being carefree about their health, knowledge of malaria, insignificant level of campaign by public and private bodies and the socio-economic status of the school [28].

“The high prevalence of malaria parasites obtained in this study is worrisome because high-density urban African populations are not often considered particularly vulnerable to malaria infection. In other West African urban areas, malaria prevalence rates from 2% to 16% have been reported with large variation between communities” (Sabot et al., 2010). According to Raghavendra et al. [29], the observed low level of sensitivity of microscopy for identifying *Plasmodium* infections is similar to findings observed elsewhere [30]. This could be attributable to 2 factors: (1) many of the infections likely occurred at low parasite densities, and (2) the laboratory technician was responsible for reading a large number of slides with low parasite prevalence over a relatively short period [31].

In general, the findings of this study may help to suggest possible solutions to most African countries and other malaria endemic regions of the world particularly to school hostels by providing first-hand information about malaria prevalence and community awareness towards the cause, transmission, prevention and control of malaria and use of LLINs [32,33].

4. CONCLUSION AND RECOMMENDATIONS

4.1 Conclusion

The result of this study has shown that a large proportion of the targeted population do not have the LLINs and usage is poor among studied population unless the sociocultural issues as well as individual factors surrounding knowledge and correct use of the LLINs are resolved malaria would continue to threaten students and the entire Nigerian society. There is therefore the need to intensify awareness on how to use LLINs and making them available to the population most at risk in the endemic rural areas.

4.2 Recommendations

To curb the prevalence of malaria in University of Uyo hostel, the following are recommended:

- Students should be properly and regularly sensitized on malaria, its prevalence, control, treatment and prevention.
- Students should be properly and regularly sensitized on the effectiveness on the use of LLINs; as it is the cheapest preventive and control measure.
- School management should join the fight against malaria in Africa by providing LLINs to new and returning students and also checkmating their usage through the welfare committee of the school.
- Schools should partner with interested government and private bodies to provide sufficient LLINs for students.
- School management should impose the necessity of LLINs as a requirement for procurement of hostel space.
- Furthermore, it would be important for policy makers and malaria supporting partners to take appropriate measures for the effective prevention and control of malaria in developing nations like Nigeria.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Ethical approval was obtained from the Department of Animal and Environmental Biology, University of Uyo.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. WHO (World Health Organization) World Malaria Report; 2014.
Available:http://www.who.int/malaria/publications/world_malaria_report_2014/en/ Last accessed March 14th, 2016.

2. WHO (World Health Organization). Global technical strategy for malaria 2016-2030; 20 Avenue Appia, 1211 Geneva 27, Switzerland; 2017.
Available: <www.who.int/about/licensing/copyright_form/en/index.html> Accessed on 7 May 2017.
3. Walter K, John CC. Malaria JAMA. 2021; 327 (6):597.
Available:<https://doi.org/10.1001%2Fjama.2021.21468>
4. Dawaki S, Al-Mekhlafi M, Ithoi I, Ibrahim J, Atroosh WM, Abdulsalam AM, Sady H, Elyana FM, Adamu AU, Yelwa SI, Ahmed A, Al-Areeqi MA, Subramaniam LR, Nasr NA, Lau Y. Is Nigeria winning the battle against malaria? Prevalence, risk factors and KAP assessment among Hausa communities in Kano State. Malar J. 2016; 15:351-365.
5. Fox T, Furnival-Adams J, Chaplin M, Napier M, Olanga EA. House modifications for preventing malaria. The Cochrane Database of Systematic Reviews. 2022 (10).
6. Garley AE, Ivanovich E, Negroustoueva S, Ye Y. Gender differences in the use of insecticide-treated nets after a universal free distribution in Kano State, Nigeria: Post-campaign survey results. Malar J. 2013;12:119-126.
7. Erinle BA, Bada EO. Prevalence of malaria infection amongst students of a southwest Nigerian Federal University. GSC Advanced Research and Reviews. 2023;15(01):110–114.
8. Atser J, Udoh PU. Dimensions in rural water coverage and access in Akwa Ibom State, Nigeria. Afr J Environ Sci Technol; 2015.
DOI: 10.5897/ AJEST2014.1805
9. Cheesebrough M. District laboratory practice in tropical countries. Part 2. Cambridge University Press. 2004;357.
10. CDC (Centers for Disease Control and Prevention) Laboratory identification of parasites of Public Health Concern; 2003.
Available:<http://www.dpd.cdc.gov/dpdx/html/diagnosticprocedures.htm> [Accessed on 19th April, 2009].
11. WHO (World Health Organization). World malaria report. Geneva, Switzerland; 2011.

12. Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U. The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature*. 2015;526 (7572):207–211.
13. WHO (World Health Organization) World Malaria Report. Switzerland: World Health Organization; 2021. ISBN 978-92-4-004049-6.
14. Gething PW, Van B, Thomas P, Smith DL, Guerra CA, Patil AP, Snow RW, Hay S. Modelling the global constraints of temperature on transmission of *Plasmodium falciparum* and *P. vivax*; 2011.
15. Okonko IO, Soley FA, Amusan TA, Ogun AA, Udeze AO, Nkang AO, Ejembi J, Faleye TOC. Prevalence of malaria plasmodium in Abeokuta, Nigeria. *Malaysian Journal of Microbiology*. 2009;5 (2):113-118
16. Autino B, Noris A, Russo R, Castelli F. Epidemiology of Malaria in endemic areas. *Mediterranean Journal of Hematology and Infectious Diseases*. 2012;4:12.
17. Abubakar BM, Haruna A, Moi IM, Katagum YM. Prevalence of malaria infection and Associated Risk Factors among students of Bauchi State University Gadau, Bauchi State, Nigeria. *Gadau Journal of Pure and Allied Sciences*. 2012;1(2):95-102.
18. Olusegun-Joseph TS, Oboh MA, Uduak MU. A Survey of malaria prevalence and antimalarial preventive measures amongst students of University of Lagos, Nigeria. *Afr. J. Clin. Exper. Microbiol*. 2016;17(4): 267-273
19. Mgbemena IC, Ezea CO, Ebe TE, Udensi UJ, Nwachukwu AA, Nzenwa DC, Nwannah AL. Asymptomatic malaria among students of Federal University of Technology, Owerri (FUTO), Imo State, Nigeria. *Biological Sciences and Pharmaceutical Research*. 2016;4(6)50-57.
20. Yhdego TG, Gardew AD, Yifat FT. Malaria prevalence, knowledge and associated factors among household heads in Maygaba town, Ethiopia. *Plos Glob Public Health*. 2022;2(3):15.
21. Murray CJ, Rosenfeld LC, Lim SS, Andrews KG, Foreman KJ, Haring D, Fullman N, Naghavi M, Lozano R, Lopez AD. Global malaria mortality between 1980 and 2010: A systematic analysis. *Lancet*. 2012;379 (9814):413–431.
22. Natama HM, Rovira-Vallbona E, Somé MA, Zango SH, Sorgho H, Guetens P, Coulibaly-Traoré, M, Valea I, Mens PF, Henk DF, Schallig H, Kestens L, Tinto H, Rosanas-Urgell A. Malaria incidence and prevalence during the first year of life in Nanoro, Burkina Faso: a birth-cohort study. *Malar J*. 2018;17:16311.
23. WHO (World Health Organization). Malaria prevention through Long lasting nets. Geneva: WHO. 2016;3
24. Biadgilign S, Reda A, Kedir H. Determinants of ownership and utilization of insecticide-treated bed nets for malaria control in Eastern Ethiopia. *J Trop Med*. 2012;12:235-245.
25. Ragasa AS, Wodajo LT, Demissie HF. Proper utilization of long lasting treated net and associated factors at ZuwayDugda district, Arsi zone, Ethiopia. *J Public Health Epidemiol*. 2019;11:13–24.
26. Shiferaw M, Alemu M, Tedla K, Tadesse D, Bayissa S, Bugssa G. The Prevalence of Malaria in TselemtiWereda, North Ethiopia: A Retrospective Study. *Ethiop J Health Sci*. 2018;1;28:539–546.
27. Taremwa IM, Ashaba S, Adrama HO, Ayebazibwe C, Omoding D, Kemeza I. Knowledge, attitude and behaviour towards the use of insecticide treated mosquito nets among pregnant women and children in rural Southwestern Uganda. *BMC Public Health*. 2017;17(1): 794-805.
28. Singh M, Brown G, Rogerson SJ. Ownership and use of insecticide-treated nets during pregnancy in sub-Saharan Africa: A review. *Malar J*. 2013;1(12):268-274.
29. Raghavendra K, Barik TK, Reddy BP, Sharma P, Dash AP. Malaria vector control: From past to future. *Parasitology Research*. 2011;108(4):757–779.
30. Jimenez-Ruiz E, Morlon-Guyot J, Daher W, Meissner M. Vacuolar protein sorting mechanisms in apicomplexan parasites. *Molecular and Biochemical Parasitology*. 2016;209(1–2):18–25.
31. Carter R, Mendis KN, Roberts D. Spatial targeting of interventions against malaria. *WHO Bull*. 2000;78: 1401-1411.

32. Westling J, Yowell CA, Majer P, Erickson JW, Dame JB, Dunn BM. Plasmodium falciparum, P. vivax, P. malariae: A comparison of the active site properties of plasmepsins cloned and expressed from three different species of the malaria parasite. *Experimental Parasitology*. 1997;87(3):185–193.
33. White NJ. Plasmodium knowlesi: The Fifth Human Malaria Parasite. *Clinical Infectious Diseases*. 2008;46(2):172–173.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/121466>