



The Effect of Using Compost on Cowpea Productivity in the Semi-Arid Region of Brazil

Debora Viviane Alves ^{a++}, Paulo César Ferreira Linhares ^{b#*},
Maria Eduarda Pereira dos Santos ^a,
Leonardo Angelo Mendonça ^a,
Rita de Cássia dos Santos Dantas ^a,
Mateus de Freitas Almeida dos Santos ^{ct},
Bruno Lucas Medeiros de Freitas ^a,
Thiffany Louhane Alves Dantas ^a,
Maria Cecília Costa Godeiro ^a, Lunara de Sousa Alves ^{d‡},
Jezimiel Oliveira da Silva ^a, Uilma Laurentino da Silva ^c,
Walter Martins Rodrigues ^{c^} and Bianca Alves Souza ^{a##}

^a Researcher and Member of the Jitirana Research group of the Federal Rural University of the Semi-Arid, Mossoró, RN, Brazil.

^b Federal Rural University of the Semi-Arid, leader of the Jitirana Research Group, Mossoró, RN, Brazil.

^c Federal Rural University of the Semi-arid, Mossoró, RN, Brazil.

^d Jitirana Research Group (National Council for Scientific and Technological Development-CNPQ), Mossoró, RN, Brazil.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajrcs/2024/v9i3291>

Open Peer Review History:

⁺⁺ Agronomy Engineering Student,

[#] PhD Researcher;

[†] Master's Student in Phytotechnics;

[‡] Researcher;

[^] PhD Researcher of Phytotechnics;

^{##} Professor and PhD;

*Corresponding author: E-mail: paulolinhares@ufersa.edu.br;

Cite as: Alves, Debora Viviane, Paulo César Ferreira Linhares, Maria Eduarda Pereira dos Santos, Leonardo Angelo Mendonça, Rita de Cássia dos Santos Dantas, Mateus de Freitas Almeida dos Santos, Bruno Lucas Medeiros de Freitas, Thiffany Louhane Alves Dantas, Maria Cecília Costa Godeiro, Lunara de Sousa Alves, Jezimiel Oliveira da Silva, Uilma Laurentino da Silva, Walter Martins Rodrigues, and Bianca Alves Souza. 2024. "The Effect of Using Compost on Cowpea Productivity in the Semi-Arid Region of Brazil". *Asian Journal of Research in Crop Science* 9 (3):76-85. <https://doi.org/10.9734/ajrcs/2024/v9i3291>.

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121594>

Original Research Article

Received: 01/07/2024

Accepted: 03/09/2024

Published: 07/09/2024

ABSTRACT

The production of cowpea is part of the agricultural activities of farmers in the semi-arid region of Brazil, being important in the aspect of acquiring income and source of food, in this sense, the use of rooster tree (*Calotropis procera*) mixed with cattle manure in cowpea productivity in the semi-arid region of Brazil. The experiment was conducted from August to November 2023 at the Rafael Fernandes experimental farm, in the district of Alagoinha (5°03'37 "S, 37°23'50" W), northwest of Mossoró, State of Rio Grande do Norte, Brazil, with an area of 400 hectares. The experimental design used was randomized complete blocks in a 4 x 2 factorial scheme, with three replications. The first factor consisted of four quantities of cattle manure (0.0; 1.1; 2.2 and 4.4 kg m⁻²), the second factor being the presence and absence of rooster tree. Each plot consisted of dimensions of 2.2 m x 1.2 m, with a total area of 2.64 m². Each plot had eight holes with three plants, making a population of 90,900 plants ha⁻¹. After harvesting the dried pods, they were stored in plastic bags, identified and taken to the DCAF/UFERSA Post-Harvest laboratory, where the following characteristics were measured: pod length, expressed in cm, number of pods plant⁻¹, expressed in plant⁻¹ units, weight of five pods, expressed in grams, weight of 100 grains, expressed in grams and dry grain productivity, expressed in kg ha⁻¹. The highest cowpea productivity was observed at a dose of 4.4 kg m⁻², with a maximum value of 2,858.85 kg ha⁻¹. The presence of rooster tree did not influence any of the characteristics evaluated. The use of cattle manure is an option for farmers in the semi-arid region.

Keywords: Food production; agroecological agriculture and family farming.

1. INTRODUCTION

Cowpea (*Vigna unguiculata* (L.) Walp.) is an annual legume belonging to the Fabaceae family and native to Central Africa [1], it is cultivated mainly in the dry areas of the tropics in Latin America, Africa and south from Asia [2]. Its grain production worldwide is more than 6.0 million tons in an area of more than 12 million hectares [3]. Its grain production worldwide is more than 6.0 million tons in an area of more than 12 million hectares [3].

This crop is a source of protein for a large number of people living in semi-arid regions, with a chemical composition of 23.4% protein, 1.8% fat and 60.3% carbohydrates, in addition to being an important source of calcium and iron [1]. In Brazil, cowpea cultivation is of paramount importance for agricultural systems, especially family agricultural systems, occupying more than 1.2 million hectares annually [4].

In the Northeast, where agriculture is largely family-based, the crop stands out for its rusticity in the semi-arid climate, cultivated mainly for the production of dry or green grains for human consumption [5]. The use of local varieties has several advantages linked to production sustainability, such as resistance to diseases, pests and climate imbalances, and seeds can be stored for subsequent harvests, which reduces production costs [6].

According to Silva et al.[7] manure is a viable option for cowpea production among farmers in the semi-arid region, as it allows for an increase in organic matter, enabling an increase in productivity. Some research has shown the cultivation of landraces for grain production. Souza et al.[8] obtained values between 10.75 (BRS Itaim) and 14.85 grains per pod (BRS Tumucumaque), with an overall average of 13.04 grains per pod [9] highlights the BRS xique xique cultivar (13.0 grains/pod) as an option for farmers in rural Paraiba.

Social technologies that promote the obtaining of renewable sources are attractive not only for their environmental advantages, but also for improving social and economic aspects. One of the widespread technologies is the use of biofertilizer, which can alleviate the problems of family farmers in agroecological systems or in conversion, and can be used both as a preventative against insect pests and diseases and as a source of the main nutrients for plants [10].

Given the importance of mixing organic fertilizers in the production of large crops, the objective was to study the effect of using compost on cowpea productivity in the semi-arid region of Brazil.

2. MATERIALS AND METHODS

2.1 Experimental Area

The experiment was conducted from August to November 2023 at the Rafael Fernandes experimental farm, in the district of Alagoinha (5°03'37 "S, 37°23'50" W), northwest of Mossoró, State of Rio Grande do Norte, Brazil, with an area of 400 hectares [11]. According to Carmo et al. [12] and the Köppen classification, the local climate is BSwH', dry and very hot, with a dry season, often from June to January, and a rainy season from February to May, average annual precipitation of 673.9 mm and average relative humidity of 68.9%.

The region's climate, according to the Köppen Geiger classification, is dry and very hot with two seasons: a dry season starting in June and ending in January and a rainy season starting in February and ending in May [13]. The soil in the experimental area was classified as a typical dystrophic red yellow argisol, with a sandy loam texture [14].

Before setting up the experiment, soil samples were collected from the 0-20 cm arable layer, homogenized, and sent to the soil chemistry and fertility laboratory for the following analyses: pH (water) = 6.6; EC = 0.20 dS m⁻¹; organic matter = 1.2 g kg⁻¹; nitrogen= 0.15 g kg⁻¹; phosphorus = 28.92 mg dm⁻³; potassium= 38.17 mg dm⁻³; calcium= 19.24 cmolc dm⁻³; magnesium = 8.74 cmolc dm⁻³; sodium = 3.27 mg dm⁻³; copper= 0.47 mg dm⁻³; iron= 2.35 mg dm⁻³; molybdenum = 9.41 mg dm⁻³; and zinc= 2.48 mg dm⁻³.

2.2 Experimental Design

The experimental design used was randomized complete blocks in a 4 x 2 factorial scheme, with three replications. The first factor consisted of four quantities of cattle manure (0.0; 1.1; 2.2 and 4.4 kg m⁻²), the second factor being the presence and absence of rooster tree.

Each plot consisted of dimensions of 2.2 m x 1.2 m, with a total area of 2.64 m². Each plot had eight holes with three plants, making a population of 90,900 plants ha⁻¹ (Fig. 1).



Fig. 1. Representation of the experimental area with cultivation of cowpea (*Vigna uniculata* L.), a Creole cultivar (Canapum), at different stages of development

The preparation of the area was carried out through manual weeding, followed by marking the experimental plots, opening the pits and adding the irrigation system with 14 mm hoses. Creole cowpea seed (*Vigna uniculata* L.) was used, coming from producer Francisco Vicente Filho from the city of Martins, RN, Brazil. Planting was carried out on August 9, 2023 at the Rafael Fernandes Experimental Farm, belonging to the Universidade Federal Rural do Semi-Árido, Brazil.

Manual weeding was used to remove spontaneous plants that competed with the cowpea crop for water and nutrients at all stages of crop development. Irrigation was carried out by sprinkler, with a daily irrigation shift divided into two applications (morning and afternoon).

2.3 Chemical Composition of Rooster Tree (*Calotropis procera*) and Cattle Manure

The rooster tree (*Calotropis procera*) was harvested within the campus of the Federal Rural University of the Semi-arid (UFERSA), being cut from the apex of the plant to the green insertion

of the stem (Fig. 2). Samples were taken from the rooster tree for carbon (C) analysis; nitrogen (N); phosphorus (P); potassium (K⁺); calcium (Ca²⁺); magnesium (Mg²⁺) and carbon/nitrogen ratio. The values observed were: 550 g kg⁻¹ C, 20.0 g kg⁻¹ N, 11.1 g kg⁻¹ P, 15.5 g kg⁻¹ K, 10.6 g kg⁻¹ Ca, 13.7 g kg⁻¹ Mg and a nitrogen/carbon ratio of 28/1.

The cattle manure was collected in the cattle sector at UFERSA, from dairy cows, where they feed roughage and concentrate, whose concentration presented the following values: 7.45 g kg⁻¹ of nitrogen (N); 191.15 g kg⁻¹ of organic matter (MO); 865.4 mg dm⁻³ of phosphorus (P); 2547.7 mg dm⁻³ of potassium (K⁺); 465.9 mg dm⁻³ of sodium (Na⁺); 7.14 cmolc dm⁻³ of Ca²⁺; 0.40 cmolc dm⁻³ of Mg²⁺ and 13.02 cmolc dm⁻³ of cation exchange ability (CTC).

2.4 Evaluated Characteristics of Cowpea

Four harvests were carried out in the experimental area with cowpea, starting on October 25, 2023 and ending on November 30, 2023, with the objective of harvesting dry grains (Fig. 3).



Fig. 2. Rooster tree (*Calotropis procera*) in the vegetative (a) and reproductive (b) phase in a vegetation area within UFERSA



Fig. 3. Cowpea plants in the flowering period and producing dry pods in the semi-arid region of Brazil

After harvesting the dried pods, they were stored in plastic bags, identified and taken to the DCAF/UFERSA Post-Harvest laboratory, where the following characteristics were measured: Pod length [twenty pods were measured per treatment, expressed in cm (CV)], Number of pods plant⁻¹ [the total number of pods in the experimental plot were counted, divided by the number of plants, expressed in plant⁻¹ units (NVP)], weight of five pods [In each experimental plot, five representative pods were removed, weighed on an electronic scale with a precision of 0.001g, expressed in grams (P05V)], weight of 100 grains [It consisted of weighing 100 grains of each experimental plot on a 0.001g precision scale, expressed in grams (P100G)] and dry grain productivity [It consisted of measuring the grain weight of each experimental plot multiplied by ha, expressed in kg ha⁻¹].

2.5 Statistical Analysis

Statistical analysis was performed according to conventional methods of analysis of variance [15], using ESTAT statistical software [16]. The response curve fitting procedure was performed using the ESTAT Software [16], applying regression analysis and conducting hypothesis testing that helps the researcher accept or reject a statistical hypothesis based on experimental results [17,18].

3. RESULTS AND DISCUSSION

No significant interaction was observed between the doses of cattle manure and the absence and presence of rooster tree on the agronomic

characteristics of cowpea, demonstrating that the factors behaved independently. However, significance was observed at the level of P<0.01 probability in the factor cattle manure doses (DE) in the characteristics pod length, number of pod plant⁻¹, weight of 100 grains, weight of 05 pods and productivity of grains.

However, significance was observed at the level of P<0.01 probability in the factor cattle manure doses (DE) in the characteristics pod length, number of pod plant⁻¹, weight of 100 grains, weight of 05 pods and productivity of grains. For the factor absence and presence of rooster tree, no statistical difference was observed (Table 1).

The different doses of cattle manure contributed positively to the increase in all characteristics, which is probably due to the increase in organic matter in the soil, with nutrient availability and greater water retention in the 0-20 cm layer.

For the pod length characteristic, there was an increase of 1.6 cm between the highest dose 4.4 kg m⁻² (19.2 cm) due to the absence of fertilization, dose 0 kg m⁻² (17.6 cm). (Fig. 4). Regarding the presence and absence of rooster tree (*Calotropis procera*) there was no statistical difference with values of 18.5 and 18.0 cm per plant respectively (Table 2). Zumba [19], studying the cultivation of cowpea using inoculants with organic and mineral fertilizers, found a maximum pod length of 20.5 cm, higher than the aforementioned research. According to Pereira et al.[20], agronomic characteristics of beans are affected by increasing amounts of cattle manure.

Table 1. F values for pod length, expressed in cm (CV), number of pods plant⁻¹ (NVP), weight of 100 grains, expressed in g (P100G), weight of five pods, expressed in g (P05V), weight of 100 grains (W100G) and dry grain productivity, expressed in kg ha⁻¹ (DGP) of cowpea (*Vigna unguiculata* L.) in the semi-arid region of Brazil

Causes of variation	GL	CV	NVP	P100G	P05V	W100G	DGP
Doses of manure (DE)	3	6.23**	7.34**	10.52**	9.57**	13.65**	15.70**
Absence and presence (AP)	1	2.23 ^{ns}	1.81 ^{ns}	0.12 ^{ns}	1.21 ^{ns}	0.89 ^{ns}	3.28 ^{ns}
DE X AP	3	2.81 ^{ns}	1.74 ^{ns}	2.93 ^{ns}	1.37 ^{ns}	1.21 ^{ns}	0.70 ^{ns}
Treatments	7	-----	-----	-----	-----	-----	-----
Block	2	6.85**	8.34**	8.60**	5.24*	4.56**	9.34**
Residue	14	-----	-----	-----	-----	-----	-----
Overall average	----	18.27	6.76	26.69	28.66	26.6	2,291.3
CV (%)	---	3.68	10.46	6.29	7.32	8.14	17.0

Non-significant effect by F test (ns); ** Significant effect by F test at 1% probability level; *Significant effect by F test at 5% probability level




Fig. 4. Pod length as a function of different doses of cattle manure incorporated into the soil in the semi-arid region of Brazil

In the characteristic number of pods per plant there was an increase of 60.6% due to the doses of 4.4 kg m⁻² (8.5 pods plant⁻¹) due to the absence of fertilization (5.3 pods plant⁻¹) with a maximum value of 8.5 pods plant⁻¹ (Fig. 5). For the presence and absence of rooster tree (*Calotropis procera*) factor, no statistical difference was observed with maximum values of 7.3 and 6.2 respectively (Table 2). Alves et al. [21], studying the production of cowpea depending on different dosages and

concentrations of biofertilizers, found a number of plant⁻¹ pods of 20.75 units, higher than the aforementioned research.

For the characteristic weight of five pods, there was an increase of 2.9g due to the higher dose of 4.4 kg m⁻² (31.5g) and the absence of fertilization (28.6g). (Fig. 6). Regarding the presence and absence of silk flower, no statistical difference was observed with values of 30.3 and 29.3g respectively. (Table 2).




Fig. 5. Number of plant⁻¹ pods as a function of different doses of cattle manure incorporated into the soil in the semi-arid region of Brazil.




Fig. 6. Weight of five pods depending on different doses of cattle manure incorporated into the soil in the semi-arid region of Brazil

For the weight characteristic of 100 grains, there was an increase of 2.0g due to the dose of 4.4 kg m⁻² (28.2g) and the absence of fertilization (26.2g) (Fig. 7). For the factor presence and absence of rooster tree there was no statistical difference with maximum values of 26.8 and 26.5g, respectively (Table 2). The time of incorporation of the fertilizers probably influenced so that there was no statistical difference. Zumba [19], studying the cultivation of cowpea using inoculants with organic and mineral fertilizers, with a weight of 100 g of seeds of 20.88g, lower than the aforementioned research. Alves et al.[21], studying the production of cowpea depending on different dosages and concentrations of biofertilizers, found the weight of 100 seeds to be 17.15g, lower than the aforementioned research. Probably the quality of the biofertilizer contributed greatly to a lower value.

For productivity, there was an increase of 1.1 kg ha⁻¹ due to the higher dose of 4.4 kg m⁻²

(2,858.85 kg ha⁻¹) and the absence of fertilization (1,712.78 kg ha⁻¹) (Fig. 8). For the factor presence and absence of rooster tree, no statistical difference was observed with values of 2,288.7 kg ha⁻¹ and 2,193.9 kg ha⁻¹ respectively (Table 2). [19], studying the cultivation of cowpea using inoculants with organic and mineral fertilizers found productivity of 840.49 kg ha⁻¹, lower than the aforementioned research. Galbiatti et al. [22] obtained pod bean productivity of 926.3 kg ha⁻¹, with the application of cattle manure biofertilizer, a value below that of the aforementioned research. Probably the quality of the mixture of biofertilizer with cattle manure compromised productivity [23], studying the yield of cowpea cultivated with cattle manure and mineral fertilizer, found a dry grain yield of 3.0 t ha⁻¹, equivalent to 3,000 kg ha⁻¹, a value higher than the aforementioned research. Probably, during the growth and development of bean plants, the mixture of cattle manure with mineral fertilizer greatly contributed to a balanced form of the crop's nutritional needs.




Fig. 7. Weight of 100 grains depending on different amounts of cattle manure incorporated into the soil in the semi-arid region of Brazil




Fig. 8. Productivity of dry cowpea grains as a function of different doses of cattle manure incorporated into the soil in the semi-arid region of Brazil

Table 2. Average values by F test, for pod length, expressed in cm (CV), number of pods plant⁻¹ (NVP), weight of 100 grains, expressed in g (P100G), weight of five pods, expressed in g (P05V), weight of 100 grains (W100G) and dry grain productivity, expressed in kg ha⁻¹ (DGP) of cowpea (*Vigna unguiculata* L.) in the semi-arid region of Brazil

Forms of application	CV	NVP	PCV	P100G	PG
Presence of the rooster tree	18.5 a	7.3 a	30.3 a	26.8 a	2,388.7 a
Absence of the rooster tree	18.0 a	6.2 a	29.3 a	26.5 a	2,193.9 a
Average	18.3	6.8	29.8	26.6	2,291.3

Averages followed by the same letter do not differ statistically at the 5% probability level.

4. CONCLUSION

The highest cowpea productivity was observed at a dose of 4.4 kg m⁻², with a maximum value of 2,858.85 kg ha⁻¹. The presence of rooster tree did not influence any of the characteristics evaluated. The use of cattle manure is an option for farmers in the semi-arid region.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

The authors of this scientific article readily declare that there is no commitment to generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.). There is no relationship.

ACKNOWLEDGMENTS

To the Hairy woodrose (*Merremia aegyptia* L.) research group (CNPQ-National Council for Scientific and Technological Development, Brazil), a pioneer in Brazil in the use of spontaneous species from the semi-arid region as green manure in the agroecological production of leafy and root vegetables, and the Universidade Federal Rural do Semi-Árido

UFERSA, for the physical structure and granting of scientific initiation scholarships for the development of research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Gupta RK, Arya M, Kumar A, Kumari P. Study on Genetic Variability in Cowpea [*Vigna unguiculata* (L.) Walp]. Current Journal of Applied Science and Technology. 2019;33(2):1-8.
- Boukar O, Belko N, Chamarthi S, Togola A, Batiemo J, Owusu E. Cowpea (*Vigna unguiculata*): Genetics, genomics and breeding. Plant Breeding. 2018;137(3):1-10.
- Durojaye HA, Moukoubi YD, Dania VO, Boukar O, Bandyopadhyay R, Ortega-Beltrana Evaluation of cowpea (*Vigna unguiculata* (L.) Walp.) landraces to bacterial blight caused by *Xanthomonas axonopodis* pv. *vinicola*. Crop Protection. 2019;116: 77-81.

4. Marinho RDCN, Ferreira LDVM, Silva AFD, Martins LMV, Nóbrega RSA, Fernandes-Júnior PI. Symbiotic and agronomic efficiency of new cowpea rhizobia from Brazilian Semi-Arid, 2017;76(2):273-281.
5. Lima LKS. Development of cowpea due to the use of waste from the coffee industry as a source of potassium (80 f., Dissertation (Masters in Agronomy /Phytotechnics), Center for Agricultural Sciences, Federal University of Ceará, Fortaleza); 2014.
Available:<https://www.repositorio.ufc.br/>
6. Carpentieri-Pípolo V, Souza A, Silva DA, Barreto, TP, Garbuglio DD, Ferreira JM. Evaluation of Creole corn cultivars in a low-tech system. Acta Scientiarum. Agronomy. 2010;32(2): 229-233.
7. Silva RASda, Linhares PCF, Souza ASdos, Pereira MFS, Assis JPde, Sousa RPde, Neves AM, Alves LSde, Silva RIGda. Yield of Pingo de Ouro cowpea bean under different sowing densities and goat manure doses in the region of Mossoró, Brazil. Journal of Agricultural Science. 2018;10(12): 329-335.
Available:<https://ccsenet.org/journal/index.php/jas/article/view/0/37346>
8. Souza AC, Ribeiro RP, Jacinto JTD, Cintra ADAR, Amaral RS, Santos AC, Matos FS. Jatropha and bean consortium: alternative for family farming. Agrarian. 2013;6(19):36-42.
9. Santos DP, Lima LKS. Agronomic evaluation of cowpea varieties in rainfed cultivation in the municipality of Coremas-PB. Green Journal of Agroecology and Sustainable Development. 2015;10(1):218-222.
Available:<https://doi.org/10.18378/rvads.v10i1.2950>
10. Silva ME, Morilha RF, Escobar M, Quevedo L. Evaluation of organic cowpea cultivation with application of different doses of biofertilizer, Cadernos de Agroecologia. 2020;5(2):1-10.
11. Rêgo LGS, Martins CM, Silva EF, Silva JJA, Lima RNS. Pedogenesis and soil classification on an experimental farm in Mossoró, Rio Grande do Norte, Brazil. Caatinga Magazine. 2016;29(4):1036-1042.
12. Carmo Filho F, Oliveira OF. Mossoró: a municipality in the semi-arid northeast, climatic characterization and floristic aspect. Mossoró: ESAM. (Mossoroense Collection, Series B); 1995;62.
13. Beck HE, Zimmermann NE, Mcvigar TR, Vergopolan N, Berg A, Wood EF. Present and future Köppen-Geiger climate classification maps at 1-km resolution. Scientific Data. 2018;5:180214.
14. Santos JAS, Soares CMG, Corrêa AM, Teodoro PE, Ribeiro LPE, Abreu HKA (2014). Agronomic performance and genetic dissimilarity among cowpea (*Vigna unguiculata* (L.) Walp.) genotypes. Global Advanced Research Journal of Agricultural Science, 3(1):271-277.
15. Banzatto DA, Kronka SN. Agricultural experimentation. 3rd ed. Jaboticabal: FUNEP. 1995;245.
16. Barbosa JC, Malheiros EB, Banzatto D. A. ESTAT: A system for statistical analysis of agronomic trials. Jaboticabal: Unesp, Version 2.0; 1992.
17. Assis JP, Sousa RP, Linhares PCF. Statistical hypothesis testing. EdUFERSA; 2020.
Available:<https://livraria.ufersa.edu.br/wp-content/uploads/sites/165/2020/08/testes-de-hipoteses-estatisticas-edufersa.pdf>
18. Assis JP. Simple linear regression, simple linear correlation, multiple linear regression and multiple linear correlation. EdUFERSA. 2013;310.
Available:<https://livraria.ufersa.edu.br/regressao-e-correlacao-linear-simples-e-multipla/>
19. Zumba JS. Cowpea cultivation using inoculant, organic and mineral fertilizers. 2016. 49 f. Dissertation (Master's in Agricultural Production) – Garanhuns Academic Unit, Federal Rural University of Pernambuco, Garanhuns; 2016.
20. Pereira VGC, Gris DJ, Marangoni T, Frigo JP, Azevedo KD, Grzesiuck AE. Agroclimatic requirements for bean (*Phaseolus vulgaris* L.) cultivation. Brazilian Journal of Renewable Energies. 2014;3(1):32-42.
21. Alves JMA, Araújo NP, Uchoa SCP, Albuquerque JAA, Silva AJ, Rodrigues GS, Silva DCO. Agroeconomic evaluation of the production of cowpea cultivars in intercropping with cassava cultivars in the cerrado of Roraima. Revista Agro@ambiente On-line. 2009;3(1): 15-30.
22. Galbiatti JA, Silva FG, Franco CF, Caramelo AD. Development of the bean plant using biofertilizer and mineral

- fertilizer. Agricultural Engineering, Jaboticabal. 2011;31(1):167-177.
23. Oliveira AP, Araújo JS, Alves EU, Noronha MAS, Cassimiro CM, Mendonça, FG Yield of cowpea cultivated with cattle manure and mineral fertilizer. Horticulture Brasileira, Brasília. 2001;19 (1):81-84.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/121594>