



---

## Synthesis of 3-azoindoles

Harindran Suhana<sup>1\*</sup>, M. Arthanareeswari<sup>1</sup> and P. Kamaraj<sup>1</sup>

<sup>1</sup>Department of Chemistry, Faculty of Engineering and Technology, SRM University,  
Kattakulathur-603203, Tamil Nadu, India.

### Authors' contributions

*This work was carried out in collaboration between all authors. Authors MA and PK have done the editing of the paper. Author HS designed the scheme, the protocol for synthetic pathway and carried out the synthesis and characterization of the compounds. All authors read and approved the final manuscript.*

Original Research Article

Received 1<sup>st</sup> July 2013  
Accepted 14<sup>th</sup> September 2013  
Published 5<sup>th</sup> February 2014

---

### ABSTRACT

Azo compounds are an important class of organic colorants which consist of a conjugated chromophore azo (-N=N-) group in association with one or more aromatic or heterocyclic systems<sup>1</sup>. Azo dyes based on heterocyclic amines have higher tinctorial strength and give brighter dyeing than those derived from aniline based diazo components. Literature survey reveals that not much work has been done on N-alkyl 3-azoindoles and their pharmacological activity. It has been considered worthwhile to synthesise N-methyl-3-azoindoles as part of our ongoing research on the development of novel synthetic route for pyrrolo(3,2-b)indoles. We herein report a simple procedure for the synthesis of 3-azoindoles.

*Keywords: Azo-coupling; N-methyl-3-azoindoles; synthesis; electrophilic substitution.*

### 1. INTRODUCTION

It has been known for many years that azo compounds [1,2] are the most widely used class of dyes due to their versatility in various fields such as the dyeing of textile fibre, the coloring of different materials, coloured plastics, biological-medical studies and advanced applications in organic synthesis [3-11]. For the past two decades, heterocyclic coupling

---


\*Corresponding author: Email: [suhana.h@ktr.srmuniv.ac.in](mailto:suhana.h@ktr.srmuniv.ac.in);

agents [12] have received a great deal of attention because of their excellent properties which are lightfastness, good substantivity, good migration and brilliant shade. Some of them are pyrazolones, iminopyrazolones, pyridones, and pyrimidines.

Manuela et al. [13] reported that heterocyclic azo dyes derived from 5-N,N-dialkylamino 2,2'bithiophene couplers could be used as thermally stable, solvatochromic probes which therefore find applications in electro-optical devices and sensors. Lee et al. [14] attempted to synthesize a range of temporarily solubilized disperse dyes which have a wider variety of colour by using the aromatic heterocyclic compounds such as pyridone and indole as coupling components. They claimed that the light fastness of the dyes was moderate and all the other fastness results were good to excellent. Furthermore, it has been reported that addition-curable phenolic resins functionalized with phenylmaleimide-system has improved the thermal characteristics and give good mechanical performance to composites. They are desirable for application in composites for thermo-structural applications [15].


## 2. RESULTS AND DISCUSSION

Indole readily reacts with weak electrophilic reagents such as benzene diazonium ion [16] in neutral or alkaline media to give the azo coupled product at 3-position. Initially 1, 2-dimethylindole [17] was prepared in 77% yield by methylation of 2-methyl indole with methyl iodide using sodium hydride in DMF (Scheme 1). The structure of the product was confirmed by its melting point, IR and <sup>1</sup>H NMR spectra.




Scheme 1. Synthesis of 1, 2-dimethylindole

Then, 1,2-dimethylindole (1) was coupled with diazotized p-nitro aniline to afford 1,2-dimethyl-3-(p-nitrophenylazo)indole<sup>18</sup> (2) in 90% yield (Scheme 2). The melting point, IR and <sup>1</sup>H NMR spectra (Fig. 1) were in accordance with the product structure.



Scheme 2. Synthesis of 1,2-dimethyl-3-(p-nitrophenylazo)indole

The diazotization of 2-cyano-4-nitro aniline under the same conditions proved to be difficult. Finally it was found that the diazotiation of 2-cyano-4-nitro aniline proceeded with sodium nitrite and 50% sulphuric acid. The resulting diazonium salt was coupled with 1, 2 dimethylindole to give the expected product namely 3-(2'-cyano-4'nitrophenylazo)-1, 2-dimethyl indole (**3**) as a bright red solid in 85% yield (Scheme 3). The structure of this azo coupled product has been confirmed by IR, <sup>1</sup>H NMR and elemental analysis data.



**Scheme 3. Synthesis of 3-(2'-cyano-4'nitrophenylazo)-1, 2-dimethylindole**

The IR spectrum of this azo derivative showed bands at 2245 cm<sup>-1</sup> due to cyano group, at 1342 and 1519 cm<sup>-1</sup> due to nitro group, and at 1645 cm<sup>-1</sup> due to C=N group.

The <sup>1</sup>H NMR spectrum exhibited a 3H singlet at δ 2.9 due to C-methyl hydrogens, a 3H singlet at δ 3.8 due to N-methyl hydrogen's, and a 7H multiplet at δ 7.1-8.7 due to aromatic hydrogens (Fig. 2).

Further the mass spectrum exhibited a molecular ion peak at *m/z* = 318.6 and base peak at *m/z* = 143.9

### 3. EXPERIMENTAL

The IR spectra were recorded on FTIR -8300 Shimadzu Spectrophotometer (range 400cm<sup>-1</sup> -4000cm<sup>-1</sup>). <sup>1</sup>H NMR spectra were recorded on Bruker-DPX 200 (200MHz) NMR Spectrophotometer. TMS was chosen as the internal standard in DMSO-d<sub>6</sub> as solvent. The mass spectra were recorded on JOEL-JMS-DX 303 HF and Finnigan MAT-8230 GC-Mass spectrometers.

#### 3.1 N-methylation of 2-methylindole

##### 3.1.1 Synthesis of 1,2-dimethylindole (1)

To a well stirred suspension of NaH (2.4g, 100mmol) in dry DMF (80mL) at 0°C was added a solution of 2-methylindole (13.1g, 100mmol) in dry DMF and the reaction mixture was stirred at 0°C under nitrogen atmosphere. After 0.5 h, methyl iodide (6.5mL, 100mmol) was added slowly and then stirred again for 6h. The reaction mixture was poured over crushed ice and the solid obtained was filtered and dried. It was purified by chromatography over silica gel using hexane as eluent.

Yield: 10g, (77%)

Mp: 55°C (Lit<sup>17</sup>Mp 55-58°C)

### 3.2 Diazonium Coupling of 1,2-dimethylindole with 2-cyano-4-nitroaniline

#### 3.2.1 Synthesis of 1, 2-dimethyl-3-(p-nitrophenylazo)indole (2)

p-nitrophenyldiazonium salt was prepared from p-nitroaniline (0.759g, 5.5mmol), 30% hydrochloric acid (5mL) and sodium nitrite (0.380g, 5.5mmol). Its pH was adjusted by the addition of sodium acetate followed by a solution of 1, 2-dimethylindole (0.720g, 5mmol) in acetic acid (2mL) at 0-50°C. After stirring for 30 min at this temperature the reaction mixture was stirred for 2 h at room temperature. The precipitated compound was collected by filtration and washed thoroughly with cold water and desiccated over anhydrous calcium chloride. It was recrystallized from ethanol.

Yield: 1.3g, (88%)

Mp: 206-209°C dec (Lit<sup>18</sup>Mp: 209-210°C dec)

IR (KBr): 1368, 1512 (NO<sub>2</sub>)

<sup>1</sup>H NMR (CDCl<sub>3</sub>/TMS) (200MHz): 2.75(s, 3H, CH<sub>3</sub>), 3.8 (s, 3H, N-CH<sub>3</sub>), 7.3-8.6(m, 8H, Aromatic-H)

### 3.3 Diazonium coupling of 1, 2-dimethylindole with 2-cyano-4-nitroaniline in Acetic Acid

#### 3.3.1 Synthesis of 3-(2'-cyano-4'-nitrophenyl)azo-1,2-dimethylindole (3)

2-cyano-4-nitrophenyl diazonium salt prepared from 2-cyano-4-nitroaniline (0.902, 5.5mmol) 50% sulfuric acid (10mL) and sodium nitrite (0.380g, 5.5mmol) in water was neutralized with sodium acetate and added to a solution of 1,2-dimethylindole(0.725g, 5mmol) in acetic acid(5mL) at 0-50°C with stirring. After 30 min at this temperature the mixture was stirred for 2 h at room temperature. The red solid obtained was filtered and dried over anhydrous calcium chloride.

<sup>1</sup>H NMR (CDCl<sub>3</sub>/TMS): 2.9 (s, 3H,-CH<sub>3</sub>) 3.8 (s, 3H, -CH<sub>3</sub>) 7.1-8.7(m, 7H, Aromatic-H)

Mass: m/z (%): 318.6 (M+, 33.5), 171.9 (32.4), 143.9(100), 114.9 (29.8), 76.9(22.5), 57(9.3)

Analysis: C <sub>17</sub> H <sub>13</sub> N <sub>5</sub> O <sub>2</sub>	C	H	N
Calcd	63.95	4.10	21.93
Found	63.90	4.44	21.89

Yield: 1.24 g, (78%)

Mp: 260-264°C

IR (KBr): 2245(CN), 1342, 1519 (NO<sub>2</sub>)




Fig. 1. <sup>1</sup>H NMR spectrum of 1, 2-dimethyl-3-(p-nitrophenylazo)indole (2)




Fig. 2. H1 NMR spectrum of 3-(2'-130 cyano-4'-nitrophenyl)azo-1,2-dimethylindole (3)

#### 4. CONCLUSION

The above work presents a simple route for the synthesis of a novel 3-azo indole. We have synthesized and characterized a new heterocyclic azo dye 3-(2'-cyano-4'-nitrophenyl)azo-1,2-dimethylindole by the coupling reaction of 2-cyano-4-nitrophenyldiazonium ion with 1,2-dimethylindole.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Abrahart EN. Dyes and Their Intermediates. London: Pergamon Press; Oxford; 1968.
2. Bindu RL, Reghunadhan Nair CP, Ninan KN. Phenolic Resins Bearings Maleimide Groups: Synthesis and Characterization. *Journal of Applied Polymer Chemistry*. 2000;38:641-652.
3. Catino SC, Farris RE. In M. Grayson (ed.) *Azo Dyes: Concise Encyclopaedia of Chemical Technology*. New York: John Wiley and Sons; 1985.
4. Christie RM. *Color Chemistry*. United Kingdom: Royal Society of Chemistry; 2001.
5. Fadda AA, Etman HA, Amer FA, Barghout M, Mohammed KS. Azo Disperse Dyes for Synthetic Fibers. 1: 2-Methyl-and 2-phenylquinazolone Derivatives. *Journal of Chemical Technology and Biotechnology*. 1994;61:343-349.
6. Kocaokutgen H, Gumrukcuoglu IE. Synthesis and Spectral Characterization of Some Novel Azo Dyes Containing Tert-butyl Groups. *Turkish Journal of Chemistry* 995;19:219-22
7. Hartman H, Schulze M, Guenther R. Nucleophilic Substitution in Arylazo Phenols - A Simple Route for Preparing Chlorosubstituted Azobenzenes. *Dyes and Pigments*. 1991;15:255-262.
8. Law KY, Tarnawskyj IW, Lubberts, P. Azo Pigments and Their Intermediate: A Spectroscopic Assessment of Metal/Azo Complexes as Negative Charge Control Agents in Color Toners. *Journal of Imaging Science and Technology*. 1994;38:81-85.
9. Peters AT, Chisowa E. Colour-Constitution Relationships in 2-acylamino-4-N,N-diethylaminoazobenzene Disperse Dyes. *Dyes and Pigments*. 1993;22:223-238.
10. Tanaka K, Matsuo K, Nakanishi A, Jo M, Shiota H, Yamaguchi M, Yoshino S, Kawaguchi K *Chemical and Pharmaceutical Bulletin*, Synthesis and Antimicrobial Activities of Five-Membered Heterocycles Having a Phenylazo Substituent. 1984;32:3291-3298.
11. Venkataraman K. *The Chemistry of Synthetic Dyes*. Vol. 3. New York: Academic Press; 1970.
12. Schwander HR. Heterocyclic Azo Coupling Components *Dyes and Pigments*. 1982;3:133-160.
13. Manuela MRM, Ferreira AMFP, Pereira MMF, Belsley M, Moura JCVP. The Synthesis and Characterization of Heterocyclic Azo Dyes Derived from 5-N,N-dialkylamino-2,2'-bithiophene Couplers. *Dyes and pigments*. 2009;83:59-65.
14. Lee JJ, Lee W J, Kim JP. Dispersant-Free Dyeing of Polyester with Temporarily Solubilized Azo Disperse Dyes from Indole Derivatives. *Fibers and Polymers*. 2003;4:66-70.
15. Yang Y, Min Z, Yi L. A Novel Addition Curable Novolac Bearing Phthalonitrile Groups: Synthesis, Characterization and Thermal Properties. *Polymer Bulletin*, 2007;59:185-194.

16. Anthony H, Patrick PL. Electrophilic substitution in indoles. Part 14. Azo-coupling of indoles with *p*-nitrobenzenediazonium fluoroborate. *J. Chem. Soc. Perkin Trans. 2*, 1987;1483-1488.
17. Noland WE, Kuryla WC, Lange RF. The Synthesis of Carbazoles from 3-Vinylindoles with Tetracyanoethylene and Dimethyl Acetylenedicarboxylate, *J. Am. Chem. Soc.* 1959;81;6010.
18. Jackson AH, Prasitpan N, Patrick VR, Shannon, Tinker AC. Electrophilic substitution in indoles. Part 15. The reaction between methylenedi-indoles and *p*-nitrobenzenediazonium fluoroborate. *J. Chem. Soc. Perkin Trans. 1*. 1987;11:2543.

---

© 2014 Suhana et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<http://www.sciencedomain.org/review-history.php?iid=405&id=7&aid=3548>