



International Journal of Plant & Soil Science
3(9): 1055-1068, 2014; Article no. IJPSS.2014.9.003

SCIENCEDOMAIN *international*
www.sciencedomain.org



Genetic Control for Some Traits Using Generation Mean Analysis in Bread Wheat (*Triticum aestivum* L.)

Naheif E. M. Mohamed^{1*}

¹*Agronomy Department, Faculty of Agriculture, Sohag University, Egypt.*

Author's contribution

This whole work was carried out by the author NEMM.

Original Research Article

Received 7th April 2014
Accepted 13th May 2014
Published 16th June 2014

ABSTRACT

In order to study the inheritance and genetic analysis of drought tolerance indicators a six generations of P₁, P₂, F₁, F₂, Bc₁ and Bc₂ of two wheat crosses i.e., Sakha 94xTokwie (C₁) and Giza 168xTokwie (C₂) under normal irrigation (N) and drought stress (D) were studied using generation mean analysis at Faculty of Agriculture, Sohag University, Egypt. Genetic variation was found for No. of spikes/plant (NS), 100-seed weight (SW), grain yield (GY), biological yield (BY), relative water content (RWC) and chlorophyll content (CC) (N&D) in two crosses. High heterosis was observed for all studied characters (N&D) except CC in two crosses. Genetic analysis showed over dominance in the inheritance of all studied characters (N&D) in two crosses. High to moderate heritability values in broad sense were detected for all characters in both crosses. Narrow-sense heritability (C₁&C₂) ranged from 0.18 for CC (D) to 0.37 for RWC (D) in C₁. The genetic advance (C₁&C₂) was high (more than 40%) for GY (N&D), while NS, BY, RWC and CC (N&D) were moderate (14-40%), indicating the importance of direct selection for these characters. The genetic models fitted for all studied characters (N&D) in two crosses except RWC (D in C₁), indicated dominance and additive x additive gene effects. Both additive x additive and dominance x dominance effects were significant for all studied characters (N&D) in two crosses except RWC (D in C₁), supporting the presence of duplicate type of epistasis. Since several important characters are influenced by dominance and non-allelic gene interaction, it is advisable to delay selection to later generation with increased homozygosity.

*Corresponding author: E-mail: naheif1974@gmail.com;

Keywords: *Wheat (Triticum aestivum L.); drought stress; generation mean analysis; gene action.*

1. INTRODUCTION

In Egypt, wheat production is far below what is needed to meet the local consumption of the growing population resulting in increasing wheat imports. To formulate an efficient breeding program for developing drought-tolerance varieties, it is essential to understand the mode of inheritance, the magnitude of gene effects and their mode of action [1,2,3]. The plant breeder is interested in the estimation of gene effects in order to formulate the most advantageous breeding procedures for improvement of the attribute in question. Therefore, breeders need information about nature of gene action, heterosis, inbreeding depression, heritability and predicted genetic gain from selection for yield and yield components [4]. Listed three major factors that must be considered and which may limit progress in the analysis of quantitative genetic variation: the number of genes involved the type of gene action, and the genotype- environment interaction.

The genetical studies based on the means and variances of basic generations, is a simple method for estimating the gene effects for a polygenic trait and has been reviewed in many crop species. The greatest merit of generation means analysis lies in its ability to estimate the epistatic effects [5]. The possibility of epistasis accounting for a significant proportion of genetic variance of quantitative trait has been investigated extensively in previous studies in crop plants. Amount and type of epistasis can have a major consequence on both the reliability of predictions and the design of breeding program. Statistically, detection of epistasis using generation means analysis is more reliable and efficient than by the analysis of variance approach [6]. However, it has its own limitations and several assumptions. Triple test cross is a powerful method of genetic analysis, which provides unbiased estimates for epistasis. In addition, it also estimates the additive and dominance components of variation with high accuracy when epistasis is absent [7]. The variance estimates attributed to environment, total genetic, additive and dominance deviation effects were obtained from the phenotypic variances for populations P1, P2, F1, F2, BC1 and BC2. These estimates allowed the determination of heritabilities in the broad and narrow sense, mean degree of dominance and minimum number of genes that control each character, by using Burton's expression [8]. The objective of the present investigation was to investigate the genetic analysis of quantitative indicators of drought tolerance in wheat under drought condition using generation mean analysis.

2. MATERIALS AND METHODS

2.1 Plant Materials and Experiments

The two Egyptian cultivars, Sakha 94 and Giza 168 were more adapted in Egypt and proved high yielding. However, the introduced line (Tokwie) is characterized as a drought tolerant. Therefore, the line introduced was crossed with the Egyptian cultivars in order to enlarge the variability for selection in the breeding program for these characters.

The experiments reported herein were carried out during the three successive growing seasons of 2010/2011, 2011/2012 and 2012/2013. In 2010/2011, the parent genotypes of hexaploid wheat (*Triticum aestivum* L.) were sown to secure enough hybrid seed (Table 1). Two crosses namely Sakha 94 x Tokwie (Cross 1) and Giza 168xTokwie (Cross 2) were developed at Faculty of Agriculture, Sohag University, Egypt.

Table 1. Pedigree and origin of the genotypes used in the two bread wheat crosses

Cross	Parental name	Pedigree	Origin
Cross 1	Sakha 94 (P1)	Opata/Rayon//Kauz	Egypt
	Tokwie (P2)	-----	South Africa
Cross 2	Giza 168 (P1)	Mill/Buc//Seri	Egypt
	Tokwie (P2)	-----	South Africa

In 2011/2012 season, F₁ plants were selfed to produce F₂ seeds and backcrossed to the parents to produce BC₁ and BC₂ seeds. In 2012/2013 season, the parents (P₁ and P₂), the first (F₁) and second (F₂) generation hybrids and the first (P₁×F₁=BC₁) and second (P₂×F₁=BC₂) backcrosses were grown in two experiments in a randomized complete blocks design with two replicates for each one. Each replicate consisted of 20 grains in one row for each of the parents and F₁, 40 grains in two rows of each of back cross and 80 grains in four rows for the F₂ population. Rows were 2.0 m long and 30 cm apart and 10 cm between plants. The first experiment was under normal irrigation (N) (gave irrigation when ever required), the second experiment was under drought stress (D) (after the emergence of 50% of the spikes, the water stress treatment received no more water until harvesting). The soil was fertilized at the rate of 20 kg/fed (15% P₂O₅) and 80 kg/fed (33.5% ammonium nitrate) and weeds were controlled by hand.

Data were recorded on 5 competitive individual plants for non-segregate basis as (P₁, P₂ and F₁) and 10 plants for BC₁ and BC₂ and 60 plants for F₂ population for each replicate follows:

1. No. of spikes/plant (NS).
2. 100-seed weight (SW) in grams.
3. Grain yield/plant (GY) in grams.
4. Biological yield/plant (BY) in grams.
5. Relative water content (RWC): A 4 cm segment of the youngest leaf was taken and cut into 2 cm segments and weighed (fresh weight=FW). Then the segments were placed in distilled water for 4 hours and reweighed to obtain turgor weight (TW). Thereafter the leaf segments were oven dried and weighed (dried weight=DW). RWC was calculated using the formula of [9], $RWC\% = [(FW - DW) / (TW - DW)] \times 100$.
6. Chlorophyll content (CC). Chlorophyll content was measured using a SPAD-502 chlorophyll meter (Minolta, Japan). For this measurement the average of three leaves per plant per replication per treatment was taken.

2.2 Statistical Analysis

Analysis of variance and mean comparison of the characters was done using SAS Software. Generation mean analysis was performed using Mather and Jinks method [5]. In this method the mean of each character is indicated as follows:

$$Y = m + \alpha [d] + \beta [h] + \alpha^2 [i] + 2\alpha \beta [j] + \beta^2 [1]$$

Where:

- Y = The mean of one generation
- m = The mean of all generation
- d = The sum of additive effects

- h = The sum of dominance effects
- i = The sum of additive x additive interaction (complementary)
- l = The sum of dominance x dominance interaction (duplicate)
- j = Sum of additive x dominance and α , 2α β and β^2 are the coefficients of genetic parameters.

The genetic parameters (m, [d], [h], [l], [j], [1]) were tested for significance using a t-test.

To estimate the parameters and to select the most suitable model the least squares method and the joint scaling test of Mather and Jinks [5] were employed.

Potence ratio, was estimated by using the formula of Smith [10].

Stress Tolerance index (STI) for grain yield were computed as formula using by [1].

$$STI = (GY_N)(GY_D) / (GY_N)^2$$

Where GY_N is grain yield under normal irrigation and GY_D is grain yield under drought.

Broad-sense (H_b^2) and narrow-sense (H_n^2) heritability were estimated by [11]. Formulas:

$$H_b^2 = [V_{F2} - (V_{P1} + V_{P2} + V_{F1}) / 3] / V_{F2} \quad H_n^2 = [2V_{F2} - (V_{BC1} + V_{BC2})] / V_{F2}$$

Genetic advance was calculated [12] with a selection intensity of $i=5\%$ for all the characters as:

$$G_A = i.H_b.\sqrt{V_{F2}}$$

The components of variation for six generations were calculated by the formulae of F2 variance were obtained by the following formula of Mather and Jinks [5] as:

$$\begin{aligned} E &= 1/3 (V_{P1} + V_{P2} + V_{F1}) \\ D &= 4V_{F2} - 2 (V_{BC1} + V_{BC2}) \\ H &= 4(V_{F2} - 1/2V_D - V_E) \\ F &= V_{BC1} - V_{BC2} \end{aligned}$$

Where:

- D - Additive genetic variance
- H - Dominance variance
- E - Environmental component of variance
- F - Correlation between D and H over all loci

3. RESULTS AND DISCUSSION

The analysis of variance (Table 2) revealed significant differences for two environments and generations for No. of spikes/plant (NS), 100-seed weight (SW), grain yield (GY), biological yield (BY), relative water content (RWC) and chlorophyll content (CC) in two crosses, indicating the existence of genetic variation and possibility of selection for drought tolerance. The genotypes x environments interaction was also significant for all studied characters in C_2 , except for GY, displaying their similar response and different responses of other traits. While, the genotypes x environments interaction was non-significant for all studied

characters in C₁. Genetic variation was found in wheat for NS, SW, BY and GY by [12,13,14] and for RWC by [1,15].

Table 2. Pedigree and origin of the genotypes used in the two bread wheat crosses

SOV	df	Mean square					
		NS	SW	BY	GY	RWC	CC
Cross 1							
Environments (A)	1	9.61**	5.58**	10859.21**	715.70**	1234.65**	659.63**
Error	2	0.05	0.08	7.02	45.63	3.33	2.41
Generations (B)	5	8.94**	0.82**	283.17*	191.21**	120.94**	227.39**
A x B	5	0.35 ^{ns}	0.13 ^{ns}	25.55 ^{ns}	7.54 ^{ns}	12.54 ^{ns}	11.98 ^{ns}
Error	20	0.25	0.07	84.49	14.90	15.23	1.05
Cross 2							
Environments (A)	1	14.06**	9.06**	11600**	620.63**	1441.29**	1416.27*
Error	2	0.05	0.001	149.70	0.08	11.95	2.94*
Generations (B)	5	18.49**	0.53	269.52**	207.59**	532.92**	179.34**
A x B	5	1.03**	0.10**	20.87**	5.71 ^{ns}	53.83**	58.96**
Error	20	0.17	0.02	48.62	4.13	10.39	4.26

*Ns: No. of spikes/plant; sw: 100-seed weight in grams; BY: Biological yield/plant in grams; GY: Grain yield/plant in grams; RWC: Relative water content %; CC: Chlorophyll content * and** significant at 5% and 1% levels of probability, respectively*

The data six generations means (Table 3) showed that F₁ hybrids were higher than mid-parent and or best parent for all studied characters under both conditions in two crosses except CC. These results showed the presence of heterotic effects for these characters.

In fact the development of any plant breeding program is dependent upon the existence of genetic variability. The efficiency of selection and expression of heterosis also largely upon the magnitude of genetic variability present in the plant population [1,2,16,17,18]. The potence ratio presented in table (3), its values ranged from less than one (0.11) for CC (D in C₂) to more than one (36.91) for RWC (D in C₂), indicating the presence of over dominance for all studied characters in two Crosses under normal (N) and drought stress (D) except CC (D in C₁) was partial dominance. These results are in line with those obtained by [13,16,19,20].

The highest stress tolerance index (Table 4) was revealed by the F₁ hybrid (STI=0.85 in C₁ and 0.83 in C₂), displaying the presence of heterobeltiosis for drought resistance in the F₁ hybrid, followed by P₂ (0.81) in C₁ and P₂ (0.81) and P₁ (0.80) in C₂.

The degree of dominance (h/d), broad-sense (H_b) and narrow-sense (H_n) heritabilities, genetic advance (GA) and genetic components of variation are presented in Tables (5&6), which shows that the degree of dominance (h/d) for all studied characters was greater than one in two crosses (N&D) except NS (N in C₂), indicating the presence of the overdominance type of gene action in the inheritance of these traits. Selection of these characters must therefore be delayed until the F₃ or F₄ generation. This delay permits a loss of non-additive genetics variance through inbreeding, so that the additive genetics variance can be more clearly evaluated. these results are in harmony with those obtained by [21]. Whereas they revealed that, the complex genetic behavior particularly additive and dominance components could be successfully exploited in later generation.

NS (N in C_2) was controlled by the additive type of gene action; the pedigree method of selection can be used for improved of this trait, While for characters under control of the non-additive type of gene action, biparental mating offers good prospects for increasing the frequency of genetic recombination, hastening the rate of genetic improvement, through it may be necessary to resort to heterosis breeding [1,13,15,22,23,24,25].

Heritability estimate indicates the progress from selection for plant characters is relatively easy or difficult to make in breeding program. Plant breeders, through experience, can perhaps rate a series of their response to selection. Heritability gave a numerical description of this concept. Assessment of heritability of various traits is of considerable important in crop improvement program, for example, to predict response to selection [26]. High to moderate broad-sense heritability estimates for all studied characters in two Crosses (N&D) (Tables 5&6) showed that effective progress can be mad through selection. Moderate narrow-sense heritability (0.2-0.5) was show for all studied characters in two crosses (N&D) except CC (D) in Cross 1 and RWC (D) in Cross 2 indicated low heritability estimate (less than 0.2) [27]. The difference between H_n and H_b exhibits the involvement of the dominance effect in the genetic constitution of these characters.

The variation observed between the genotypes for the characters investigated exhibited that selection maybe effective for the improvement of drought tolerance [1,2,28,29], however, the selection efficiency is related to the magnitude of heritability and genetic advance [12,17]. Heritability estimates along with genetic advance are important selection parameters and normally more helpful in predicting the gain under selection than heritability estimates alone. However, heritability estimates are influenced by the type of genetic material, sample size, method of sampling, conduct of experiment, method of calculation and effect of linkage. Genetic advance which refers to the improvement in the mean genotypic value of selected individuals over the parental population is influenced by the genetic variability, heritability and selection intensity [30,31].

The rate of genetic advance is connected with heritability [5]. The genetic advance (C_1 & C_2) was high (more than 40%) for GY (N&D), while NS, BY, RWC and CC (N&D) were moderate (14-40%), indicating the importance of direct selection for these characters and the significance of indirect selection for SW (N&D) in two crosses with low genetic advance (less than 14%) through correlated response with characters having high heritability and genetic advance [1,14,32,33,34].

Degree of dominance and variance components are presented in Tables (5&6), Ew, D and H are environmental, additive and dominance components, respectively. F is an indicator of correlation between D and H over all loci. If F is zero it means that dominant genes are in the parent with high performance, while negative F exhibits that dominant genes are in the low performance parent. If the ratio of $F/\sqrt{D \times H}$ is equal to or near one confirms that the magnitude and sign of dominance for all the genes monitoring the character is equal, therefore, the ratio $\sqrt{H/D}$ is a good estimator of dominance. If $F/\sqrt{D \times H}$ is equal to zero or close to zero, the magnitude and sign of the genes controlling the character is not equal and hence $\sqrt{H/D}$ explains average dominance. The h/d ratio estimates the degree of dominance [1,15,18,35]. The ratio of $\sqrt{H/D}$ for all studied characters (N&D) in two crosses showed average dominance except NS (D), GY (D) and CC (N&D) in C_1 and GY (N), RWC (N) and CC (N&D) in C_2 showed over dominance.

Table 3. Mean comparison of the characters studied

Generations	Characters											
	NS		SW		BY		GY		RWC		CC	
	N	D	N	D	N	D	N	D	N	D	N	D
Cross 1												
Gemmeiza 9 (P ₁)	14.87	13.17	6.10	4.84	100.97	64.60	44.49	33.76	67.97	54.67	46.47	40.17
Inbred line 1 (P ₂)	10.67	10.17	5.63	4.80	96.98	60.91	37.70	30.58	71.21	61.73	59.70	49.37
F ₁ (P ₁ x P ₂)	16.07	14.67	6.30	5.56	109.32	68.29	48.41	41.03	74.02	66.91	52.57	45.40
F ₂	13.17	12.17	5.11	4.63	82.69	52.95	32.52	23.16	69.85	55.21	39.37	33.93
P ₁ x F ₁ (BC ₁)	14.67	13.67	5.74	4.91	96.16	65.05	40.66	30.49	67.14	53.35	46.20	37.43
P ₂ X F ₁ (BC ₂)	12.67	12.07	5.36	4.78	95.70	61.61	36.93	28.18	68.50	56.55	54.30	40.93
LSD _{0.05}	1.88	1.01	0.45	0.42	4.25	5.19	4.31	3.85	2.01	1.36	3.08	3.94
Potence ratio	-1.57	-2.00	1.83	4.27	5.19	3.00	1.89	4.32	-2.73	14.97	-1.10	1.41
Cross 2												
Sids 1 (P ₁)	14.67	12.67	5.75	4.62	101.76	67.85	42.40	34.07	66.97	52.73	53.70	39.33
Inbred line 2 (P ₂)	10.67	10.17	5.63	4.80	96.98	60.91	37.70	30.58	71.21	61.73	59.70	49.37
F ₁ (P ₁ x P ₂)	16.67	14.17	6.30	4.95	111.44	79.03	47.11	38.91	86.14	78.36	67.73	44.40
F ₂	14.17	13.67	5.13	4.52	96.58	59.19	30.39	22.14	56.74	51.66	43.87	39.33
P ₁ x F ₁ (BC ₁)	14.17	13.17	6.09	5.00	100.03	62.42	35.41	26.40	74.97	55.87	52.53	40.43
P ₂ X F ₁ (BC ₂)	12.67	11.67	5.75	4.73	98.63	60.62	39.72	28.80	81.87	61.60	55.37	44.77
LSD _{0.05}	1.90	1.20	0.51	0.46	3.52	2.59	2.52	2.32	1.49	2.41	2.93	2.89
Potence ratio	-2.00	-2.20	-10.22	2.70	5.05	4.22	-2.40	-3.78	8.04	36.91	19.46	0.11

Ns: No. of spikes/plant; sw: 100-seed weight in grams; BY: Biological yield/plant in grams; GY: Grain yield/plant in grams; RWC: Relative water content %; CC: Chlorophyll content

Table 4. Grain yield/plant under normal (GY_N) and drought stress (GY_D), and stress tolerance index (STI) for each generation

Generations	GY _N	GY _D	STI	Generations	GY _N	GY _D	STI
Cross 1				Cross 2			
Gemmeiza 9 (P ₁)	44.49	33.76	0.76	Sids 1 (P ₁)	42.40	34.07	0.80
Inbred line 1 (P ₂)	37.70	30.58	0.81	Inbred line 2 (P ₂)	37.70	30.58	0.81
F ₁ (P ₁ x P ₂)	48.41	41.03	0.85	F ₁ (P ₁ x P ₂)	47.11	38.91	0.83
F ₂	32.52	23.16	0.71	F ₂	30.39	22.14	0.73
P ₁ x F ₁ (BC ₁)	40.66	30.49	0.75	P ₁ x F ₁ (BC ₁)	35.41	26.40	0.75
P ₂ x F ₁ (BC ₂)	36.93	28.18	0.76	P ₂ x F ₁ (BC ₂)	39.72	28.80	0.73

Table 5. Genetic parameters and components of variation for all studied characters in the cross 1 under normal (N) and drought stress (D) conditions

Characters	h/d	H _b	H _n	G _A	D	H	F	E _w	√H/D	F/√HXD	
NS	N	+2.65	0.69	0.28	25.06	19.80	9.07	+1.40	5.42	0.68	0.11
	D	+3.63	0.67	0.20	22.83	13.40	17.53	-0.80	5.50	1.14	-0.05
SW	N	+5.79	0.78	0.33	8.87	7.21	2.80	-0.11	1.19	0.62	-0.03
	D	+12.31	0.83	0.32	12.55	9.30	5.76	+0.60	1.27	0.79	-0.08
BY	N	+134.72	0.79	0.36	20.29	17.82	3.75	-1.60	2.56	0.46	-0.20
	D	+13.69	0.74	0.33	25.05	21.56	5.51	-1.55	4.34	0.51	0.14
GY	N	+8.94	0.79	0.29	56.63	40.35	29.26	-2.36	7.49	0.85	-0.07
	D	+13.67	0.74	0.24	46.60	28.68	33.12	+0.99	7.90	1.08	0.03
RWC	N	+2.70	0.74	0.32	21.36	18.03	5.42	-1.88	3.73	0.55	-0.19
	D	-2.40	0.76	0.37	23.76	22.37	1.39	-1.16	3.74	0.25	-0.21
CC	N	-5.31	0.73	0.21	23.43	13.25	19.00	-1.20	4.13	1.20	-0.08
	D	-6.18	0.74	0.18	19.92	9.20	20.27	2.24	3.36	1.48	0.16

Ns: No. of spikes/plant; sw: 100-seed weight in grams; BY: Biological yield/plant in grams; GY: Grain yield/plant in grams; RWC: Relative water content %; CC: Chlorophyll content

Table 6. Genetic parameters and components of variation for all studied characters in the cross 2 under normal (N) and drought stress (D) conditions

Characters	h/d	H _b	H _n	G _A	D	H	F	E _w	√H/D	F/√HXD	
NS	N	+0.67	0.73	0.30	27.98	22.50	9.33	-0.75	5.00	0.64	-0.05
	D	-1.50	0.70	0.29	25.06	20.70	7.27	-0.35	5.17	0.59	-0.03
SW	N	+11.06	0.81	0.32	10.66	8.09	4.51	+0.03	1.21	0.75	0.005
	D	+6.07	0.75	0.29	6.71	5.01	3.01	+0.50	1.08	0.78	0.13
BY	N	+16.48	0.66	0.31	17.92	16.21	2.37	-0.32	2.91	0.63	-0.02
	D	+13.32	0.77	0.26	22.80	14.91	13.45	-1.67	4.50	0.38	-0.27
GY	N	-8.51	0.71	0.24	47.64	30.87	30.76	-2.37	9.52	1.00	-0.08
	D	-11.79	0.75	0.30	53.60	41.71	20.66	+0.75	8.47	0.70	-0.03
RWC	N	-15.04	0.79	0.25	23.83	14.53	17.22	+2.08	3.16	1.09	0.11
	D	-8.63	0.77	0.19	19.62	20.62	8.74	+1.30	2.81	0.65	0.10
CC	N	-18.15	0.69	0.20	19.73	11.20	15.91	+0.88	4.25	1.19	0.07
	D	-3.03	0.80	0.26	20.01	12.34	14.16	+0.03	2.41	1.07	0.002

Ns: No. of spikes/plant; sw: 100-seed weight in grams; BY: Biological yield/plant in grams; GY: Grain yield/plant in grams; RWC: Relative water content %; CC: Chlorophyll content

The estimates of heterosis and inbreeding depression together provide information about type of gene action involved in the expression of various quantitative traits. The percentage of heterosis with regard to High Parent (HP) and Mid-Parent (MP) and Inbreeding

Depression (ID) (Figs. 1 and 2) exhibited that mid-parent and high parent heterosis were positive for NS, SW, BY, GY, RWC and CC in two crosses under both conditions except CC was negative (D) (C_1 & C_2) compared with high parent. Inbreeding depression was positive for all studied characters.




Fig. 1 (A and B). Percentage of heterosis and inbreeding depression under two environments in Cross 1 for characters investigated
 HP: High Parent; MP: Mid-Parent; ID: Inbreeding Depression




Fig. 2 (A and B). Percentage of heterosis and inbreeding depression under two environments in Cross 2 for characters investigated
 HP: High Parent; MP: Mid-Parent; ID: Inbreeding Depression

The joint scaling test [5] was employed to estimate the mean (m), additive effect (d), dominance effect (h), additive x additive (i), additive x dominance (j) and dominance x

dominance (1) values (Tables 7&8). The results of A, B, C and D scaling test for the two wheat crosses under both environments, revealed that significant of any of these tests indicates the presence of non-allelic gene interactions or epistasis on the scale of measurement used. Results of scaling test, showed that additive-dominance model is inadequate for explaining the inheritance of all studied characters, indicating the present of non-allelic gene interaction in two crosses under two environments. [36] studied the generation mean analysis in heat tolerance in wheat; they showed the adequacy of additive-dominance model for grain yield and its components.

The mean parameters (m) for all studied attributes of two crosses and environments (Tables 7&8) which reflect the contribution due to the over all mean plus the locus effects and interaction of the fixed loci were significant. The estimated of dominance gene action (h) was significant for the all studied characters (N&D) in two crosses, indicating the importance gene effects in inheritance of these characters. The significant [d] and [h] in the inheritance of RWC (D in C₂) revealed that both types of additive and dominance effects are involved in the genetics of RWC [1,2,13,15,37].

The genetic models fitted (Tables 7&8) for all studied characters (N&D) in two crosses except RWC (D in C₁), indicated dominance and additive x additive gene effects. indicated dominance and additive x additive gene effects. It is there fore suggested that selection should be carried out in late generation and the interaction should be fixed by selection under selfing conditions. The epistatic effect (dominance x dominance [1]) was significant for all studied characters (N&D) in two crosses, which confirm the important role of dominance x dominance gene interaction in the genetic system controlling, these result were reported by [13,15,24,38]. Both additive x additive [i] and dominance x dominance [1] effects were significant for all studied characters (N&D) in two crosses except RWC (D in C₁), supporting the presence of duplicate type of epistasis. This complementary interaction increases the variation between the generation and in the segregating population. The cross, which showed most promising in terms of narrow sense heritability and genetic gain, also showed highest means under both conditions, chance to find stress tolerant breeding material in segregating populations of this cross are promising. these finding are in line with [39], they studied genetic analysis of salt tolerance, and refer to High narrow sense heritability may be used as a useful indicator index for the selection of salt tolerant genotypes at the vegetative growth stage in wheat.

Table 7. Estimates of scaling test and types of gene action using generation means for all studied characters cross 1 under normal (N) and drought stress (D) conditions

Characters		Scaling test				Genetic parameters					
		A	B	C	D	m	[d]	[h]	[i]	[j]	[l]
NS	N	-1.60**	-1.40**	-5.00**	-1.00**	13.17**	2.00	5.30**	2.00**	-0.10	1.00**
	D	-0.50**	-0.70**	-4.00**	-1.40**	12.17**	1.60	5.80**	2.80**	0.10	-1.60**
SW	N	-0.91**	1.20**	-3.88**	-0.88**	5.11**	0.38	2.20**	1.77**	0.14	0.45**
	D	-0.57**	-0.80**	-2.23**	-0.43**	4.63**	-0.13	1.60**	0.86*	0.12	0.52**
BY	N	-17.97**	-14.91**	-85.85**	-26.49**	82.69**	0.47	63.32**	52.97**	-1.53	-20.09**
	D	-2.79**	-5.97**	-50.30**	-20.77**	52.95**	3.44	47.08**	41.54**	1.60	-32.78**
GY	N	-11.58**	-10.26**	-46.95**	-12.56**	32.52**	3.74	33.43**	25.11**	-0.66	-3.28**
	D	-11.81**	-13.25**	-49.78**	-12.36**	23.16**	2.31	31.58**	24.72**	0.72	-0.34**
RWC	N	-7.71**	-8.23**	-7.84**	4.05**	69.85**	-1.36	-3.67*	-8.10**	0.26	24.04**
	D	-14.89**	-15.54**	-29.40**	0.51**	55.21**	-3.21	7.69**	-1.03	0.32	31.46**
CC	N	-6.63**	-3.67**	-53.83**	-21.77**	39.37**	-8.10	43.02**	43.53**	-1.48	-33.23**
	D	-10.70**	-12.90**	-44.60**	-10.50**	33.93**	-3.50	21.63**	21.00**	1.10	2.60**

Ns: No. of spikes/plant; sw: 100-seed weight in grams; BY: Biological yield/plant in grams; GY: Grain yield/plant in grams; RWC: Relative water content %; CC: Chlorophyll content, * and ** significant at 5% and 1% levels of probability, respectively

Table 8 Estimates of scaling test and types of gene action using generation means for all studied characters in the cross 2 under normal (N) and drought stress (D) conditions

Characters		Scaling test				Genetic parameters					
		A	B	C	D	m	[d]	[h]	[i]	[j]	[l]
NS	N	-3.00**	-2.00**	-2.00**	1.50**	14.17**	1.50	1.00*	-3.00**	-0.50	8.00**
	D	-0.50**	-1.00**	3.50**	2.50**	13.67**	1.50	-2.25**	-5.00**	-0.25	6.50**
SW	N	0.13**	-0.43**	-3.88**	-1.57**	5.13**	0.34	3.76**	3.15**	0.28	-2.85**
	D	0.43**	-0.29**	-2.23**	-0.70**	4.52**	0.27	1.64**	1.40**	0.36	-1.54**
BY	N	-13.14**	-11.16**	-35.30**	-5.50**	96.58**	1.40	23.07**	11.00**	-0.99	13.30**
	D	-22.04**	-18.70**	-50.05**	-4.66**	59.19**	1.80	23.98**	9.32**	-1.67	31.41**
GY	N	-18.70**	-3.37**	-50.77**	-14.36**	30.39**	-4.32	36.77**	28.71**	-7.67	-6.65**
	D	-20.19**	-11.89**	-53.91**	-10.92**	22.14**	-2.41	28.42**	21.83**	-4.15	10.25**
RWC	N	-3.18**	6.39**	-83.50**	-43.36**	56.74**	-6.90	103.7**	86.71**	-4.78	-89.92**
	D	-19.35**	-16.88**	-64.55**	-14.16**	51.66**	-5.73**	49.44**	28.31**	-1.23	-7.91**
CC	N	-16.37**	-16.70**	-73.40**	-20.17**	43.87**	-2.83	51.37**	40.33**	0.16	-7.27**
	D	-2.87**	-4.23**	-20.17**	-6.53**	39.33**	-4.33	13.12**	13.07**	0.68	-5.97**

Ns: No. of spikes/plant; sw: 100-seed weight in grams; BY: Biological yield/plant in grams; GY: Grain yield/plant in grams; RWC: Relative water content %; CC: Chlorophyll content, * and ** significant at 5% and 1% levels of probability, respectively

4. CONCLUSION

Genetic analysis showed overdominance in the inheritance of all studied characters under normal and drought conditions in two crosses. The genetic advance in both of two crosses was high for grain yield under normal and drought conditions; meanwhile it was moderate in number of spikes, RWC and chlorophyll content. The complex genetic behavior especially both of additive x additive and dominance x dominance effects were significant for all studied characters (N&D) in two crosses except RWC under drought in cross 1, whereas several important characters are influenced by dominance and non-allelic gene interaction. It is recommended that selection for improvement of studied traits should be delayed to later generation of segregation population in wheat

ACKNOWLEDGEMENTS

The author would like to thank Dr. Ala A. S. Ahmed of Sohag University, Agronomy Dep. for his kind technical supports on the phenotypic measurements. The author also thank Dr. Mohamed Husein of Sohag University, Horticulture Dep. to provide the agricultural facilities in the research Farm.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Farshadfar E, Ghandha M, Zahravi M, Sutka J. Generation mean analysis of drought tolerance in wheat (*Triticum aestivum* L.). *Acta Agron. Hung.* 2001;49:59-66.
2. Farshadfar E, Mahjouri S, Aghaee M. Detection of epistasis and estimation of additive and dominant components of genetic variation for drought tolerance in Durum wheat. *J. Biol. Sci.* 2008b;8:598-603.
3. Iqbal M, Navabi A, Salmon DF, Young RC, Mardoch BM, Moore SS, Spaner D. Genetic analysis of flowering and maturity time in high latitude spring wheat. *Euphytica.* 2007;154:207-218.
4. Sprague GF. Orientation and objectives In *Statistical genetics and plant breeding* Nat. Acad. Sci. N.R.C. Pub. 1963;982:9-105.
5. Mather K, Jinks JL. *Biometrical Genetics.* 3rd Edn. Chapman and Hall Ltd. ;1982. ISBN-10: 0412228904.
6. Lamkey KR, Lee M. Quantitative genetics, molecular markers and plant improvement. In: *Focussed plant improvement- towards responsible and sustainable agriculture.* ed. imrie, b.c. and hacker, j.b. 1993;104-115.
7. Kearsey MJ, Jinks JL. A general method of detecting, additive, dominance and epistatic variation for metric traits.I. *Theor. Heredity.* 1968;23:403-409.
8. Burton GW. Quantitative inheritance in pearl millet (*Pennisetum glaucum* Agron. J) 1951;5.43:409-417.
9. Ritchie SW, Nguyen HT, Holiday AS. Water relations in winter wheat as drought resistance indicators. *Crop Sci.* 1990;28:526-531.
10. Smith HH. Fixing Transgressive Vigour in *Nicotiana Rustica.* In: *Heterosis.* Iowa State College Press, Ames, IA, USA; 1952.
11. Warner JN. A method of estimating heritability. *Agron J.* 1952;44:427-430.

12. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybean. *Agron J.* 1955;47:314-318.
13. Tammam AM. Generation mean analysis in bread wheat under different environmental conditions. *Minufiy J. Agri. Res.* 2005;30(3):937-956.
14. Farshadfar E, Aghaie M, Sharifi M, Yaghotipoor A. Assessment of salt tolerance inheritance in barley via generation mean analysis. *J. Biol. Sci.* 2008a;8:461-465.
15. Manette A, Schonfeld C, Richard J, Carre B, Morhinweg W. Water relations in winter wheat as drought resistance indicators. *Crop Sci.* 1988;28:526-531.
16. Amin IA. genetics behaviour of some agronomic traits in two durum wheat crosses under heat stress. *Alex.J. Agric. Res.* 2013;58(1):53-66.
17. Singh P, Narayanan SS. *Biometrical Techniques in Plant Breeding.* 1st Edn., Kalayani publishers, New Dehli, India; 1993.
18. Singh RK, Chaudhary BD. *Biometrical Methods in Quantitative Genetic Analysis.* 1st Edn., Kalyani Publisher, New Dehli, India; 1999. ISBN: 81-7663-307-0.
19. Ketata H, Smith EL, Edwards LH, McNew RW. Detection of epistatic, additive and dominance variation in winter wheat (*Triticum aestivum* L.). *Crop Sci.* 1976;16:1-4.
20. Moshref MK. *Genetical and statistical studies in wheat.* Ph.D. Thesis, Al-Azhar Univ., Egypt; 1996.
21. Zaazaa EI, Hager MA, El-Hashash EF. Genetic Analysis of Some Quantitative Traits in Wheat Using Six Parameters Genetic Model. *American-Eurasian J. Agric. & Environ. Sci.* 2012;12(4):456-462.
22. Gill KS, Dhillon SS, Bains KS. Combining ability and inheritance of yield components in crosses involving Indian and exotic wheat germplasm. *Indian J. Genet. Pl. Breeding.* 1972;32:421-430.
23. Sharma G, Singh RB. Inheritance of plant height and spike length in spring wheat. *Indian J. Genet. Pl. Breeding.* 1976;36:173-183.
24. Srivastava RB, Sharma SC, Yunus M. Additive and non-additive gene effects for yield and yield components in two crosses of wheat (*T. aestivum* L.). *India J. Genet.* 1992;52:297-301.
25. Kheiralla KA, El-Defrawy MM, Sherif THI. Genetic analysis of grain yield, biomass and harvest index in wheat under drought stress and normal moisture conditions. *Assiut J. Agric. Sci.* 1993;24:163-183.
26. Nyquist WE. Estimation of heritability and prediction of response in plant populations. *CRC Critical Reviews in Plant Sci.* 1991;10(3):235-322.
27. Tefra H, Peat HE. Genetics of grain yield and other agronomic characters in (*Eragrostis tef* Zuccrotter) II the triple test cross. *Euphytica.* 1997;96:193-202.
28. Umarahan P, Ariyanayagam RP, Haque SQ. Genetic analysis of yield and its components in vegetable cowpea (*Vigna unguiculata* L. walp). *Euphytica.* 1997;96:207-213.
29. Farshadfar E. Application of integrated selection index and rank sum for screening drought tolerant genotypes in bread wheat. *Intl J Agri Crop Sci.* 2012;4(6),325-332.
30. Alza JO, Martinez JMF. Genetic analysis of yield and related traits in sunflower (*Helianthus annus* L.) in dryland and irrigated environments. *Euphytica.* 1997;95:243-251.
31. Sharma SN. Genetics of spike length in durum wheat. *Euphytica.* 2003;130:155-161.
32. Sharma BD, Sood BC, Halotra VV. Studies on variability, heritability and genetic advance in chickpea research. *Indian J. Pulses Res.* 1991;3:1-6.
33. Sood S, Kalia NR, Bhateria S, Kumar S. Detection of genetic components of variation for some biometrical traits in *Linum usitatissimum* L. in sub-mountain Himalayan region. *Euphytica.* 2006;155:107-115.

34. Golparvar AR. Heritability and mode of gene action determination for grain filling rate and relative water content in hexaploid wheat.–Genetika. 2012;44(1):25-32.
35. Naroui Rad MR, Abdul Kadir M, Rafii MY, Jaafar ZEH , Naghavi5 MR, Ahmadi F. Genotype × environment interaction by AMMI and GGE biplot analysis in three consecutive generations of wheat (*Triticum aestivum*) under normal and drought stress conditions. AJCS. 2013;7(7):956-961.
36. Lal C, Maloo SR, Kumar V. Generation mean analysis for some heat tolerance and quantitative characters in bread wheat (*Triticum aestivum* L.). J. Wheat Res. 2013;5(2):22-26.
37. Farshadfar E, Mohammadi R, Aghaee M, Sutka J. Identification of QTLs involved in physiological and agronomic indicators of drought tolerance in Rye using multiple selection index. Acta Agron. Hung. 2003;51:419-428.
38. Kearsey M, Pooni HS. The Genetical Analysis of Quantitative Traits. 2nd Edn., Chapman and Hall, U.K. 2004; ISBN: 0-7487-4082-1.
39. Dashti H, Bihamta MR, Shirani H, Majidi MM. Genetic analysis of salt tolerance in vegetative stage in wheat (*Triticum aestivum* L) . POJ. 2012;5(1):19-23.

© 2014 Mohamed; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://www.sciencedomain.org/review-history.php?iid=547&id=24&aid=4947>