



Acacia Shrubs and Trees for Climate Change Mitigation and Economic Growth: An Imperative towards Achieving the Sustainable Development Goals in Yobe, Nigeria

I. N. Mohammed ^{a*}, A. Sule ^a, Y. K. Muhammad ^b
and K. D. Salami ^b

^a Department of Science Laboratory Technology, School of Sciences, Mai Idris Alooma Polytechnic, P.M.B. 1020 Geidam, Yobe State, Nigeria.

^b Department of Forestry and Wildlife Management, Federal University Dutse, Jigawa State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/SAJSSE/2023/v20i1691

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101253>

Review Article

Received: 10/04/2023
Accepted: 16/06/2023
Published: 09/07/2023

ABSTRACT

The need to mitigate the effect of climate change arising from over exploitation of natural resources and to improve the economic wellbeing of the common populace in Sub-Saharan Africa particularly the northern Sahelian state of Yobe. Acacia is one of the trees adapted to the Sahelian climate of Yobe that provides a lot of social and economic benefits, and it's exploited in wild state. This paper highlights the domestication of the tree using silvicultural practices and silvics of the tree by elucidating the origin, climatic and soil requirement, product they produce diseases and control

*Corresponding author: Email: ngurumuhammad@yahoo.co.uk;

methods. Findings reveal the following information based on cited literatures. The plant grows in drier Sahelian climates of Africa, parts of Arabia and parts of India and Pakistan. It produces gum Arabic, fodder for livestock and other medicinal products. Findings implies that Acacia Senegal is suitable for agro forestry based on the fact that Acacia is a nitrogen fixer and produce no thick canopy making it possible for crops to thrive well underneath their canopy.

Keywords: *Silviculture; acacia senegal; climate change; economic growth; Yobe State.*

1. INTRODUCTION

Sub-Saharan Africa has been inhabited by people for thousands of years. The area is characterized by dry and harsh climate with little rainfall which is around 800mm throughout the year [1]. In many parts of the region the major contributor to survival is the *Acacia* tree which is found across the continent [2]. Climate variability and human activities are the two major factors threatening forestry in Sudan Sahelian Africa. In this region half of the forest region are threatened [3]. In African context the major problems facing communities is deforestation of trees particularly *Acacia species* in northern States, Yobe inclusive. This has a negative impact on the life of farming communities in four major areas: - Lack of fuel, Shortage of income, Shortage of livestock fodder and Land degradation. Since 1980s, rural development agencies have been making efforts to encourage people to plant and manage local tree species that are beneficial particularly *Acacia species*. The major problems curtailing this effort include lack of information on the most appropriate species, variety or seed source for a particular locality, Lack of adequate knowledge of the silvicultural and silvics of the *Acacia species*.

The objective of this paper is to highlight the use of a tree-based rural development using taxonomy, ecology and variation of *Acacia species* which are appropriate for rural

development in the Sudano - sahelian region of Yobe State. The potentials of *Acacia* tree in addressing climate change mitigation and economic development are clearly demonstrated by adopting the silviculture and silvics of this multi-purpose Sudano-sahelian tree. *Acacia* can survive under extreme harsh conditions and give out the following benefits: -

- They act as natural repair -kits for depleted soils.
- They stabilized the microclimate by providing shade under which plants can grow and animals can feed.
- They provide pods for animal fodder, fuelwood, seeds, timber and gums which can all be used for domestic purposes and export for generating income for smallholder farmers and their families ‘

Table 1. Poverty indicators in Nigeria

Variable	Value
Population living on less than \$1.00/day	70%
Population living on less than \$2.00/ day	91%
Poverty Gap \$1.00/day	35%
Poverty Gap \$2.00/day	35%
Access to improved sanitation	59%
Access to improved water source	62%
Life expectancy (both sexes)	52 years

Source: Ibrahim [4]

Table 2. Trends in poverty level by zones (1980-2010)

Zone	1980	1985	1992	1996	2004	2010
South South	13.3	45.7	40.8	58.2	35.1	63.8
South East	12.9	30.4	41.0	53.5	26.7	67.0
South West	13.4	38.6	43.1	60.9	43.3	59.1
North Central	32.2	50.8	46.0	64.7	67.0	67.5
North East	35.6	54.9	54.0	70.0	72.2	76.3
North West	37.7	52.1	36.5	77.2	71.2	77.7

Source: Ibrahim [4]

2. POVERTY IN THE STUDY AREA

2.1 Poverty Indices and Economic Growth in Yobe State

Nigeria is blessed with huge mineral and natural resources and despite the abundance of the resources in the country, the poverty level is very alarming with nearly 100 million people living in abject poverty such that the country is referred to as the poverty capital of the World [5]. The poverty rate in the North east and North West is 77.7% and 76.3% respectively. The 10 worst states (Fig. 1) with high incidence of poverty in Nigeria are in the northern part of the country with Yobe appearing among the top ten with highest incidence of poverty [6]. Paradoxically, the country is rich, but the people are poor. Despite the abundance of mineral and natural resources, over exploitation and total mismanagement is the major problem [7].

Yobe is susceptible to environmental challenges, including desertification, erosion and flooding. Land degradation attributed to deforestation,

over-grazing by livestock, increasing demand for firewood and increased agricultural intensity has led to soil breakdown. Additionally, sand dune build up has resulted in a reduction of plant growth and productivity in the state [8]. One third of the World population uses fuelwood as a source of energy and the highest uses are from rural parts of Africa particularly northern Nigeria where it accounts for 80% of the total energy consumed [9].

In line with the Millennium Development Goals benchmark for minimum standard of livelihoods. One of the roles played by the Yobe state SDGs is to ensure that poverty is eradicated through wealth and job creation and sustainably harnessing the natural and mineral resources abundant in the state. This is in consonant with the views of Abbas and Ogiri, [10] that raising income is paramount tool in poverty reduction and overall wellbeing in any meaningful society. Yobe and Borno states have the best climate for *Acacia species* growth in Nigeria [4]. Based on this reason investing in *Acacia* domestication is a good idea.




Fig. 1. Map of Nigeria showing poverty count rate with no data for Borno as result of insecurity during the data collection time.

Source: National Bureau of Statistics, [11]




Fig. 2. Poverty headcount showing Yobe state among the top ten in the list

2.2 Silviculture and Silvics of *Acacia senegal*

The Northern state of Yobe is facing environmental degradations such as desert encroachment, flooding and drought. In addition, the global economic meltdown of 2009 and 2015 recession is another reason why we must diversify the economy [12]. Instability in the price of petroleum products, the negative impacts arising from the use of fossil fuel as a major cause of global warming and climate change necessitate the governments to focus on diversifying our economy by harnessing our natural products. In this case harnessing *Acacia*

and its related product is seen as a way of diversifying our economy.

Silviculture of *Acacia* is synonymous to domestication of *Acacia* trees. The tree is multi-purpose considering the wide number of uses it offers to people. Silviculture is the science and art of managing forest communities based on knowledge of tree silvics. More particularly, the theory and practice of controlling the establishment, composition and growth of forest vegetation [13]. Silviculture can be defined as the science of organising growth and development processes in trees and forest ecosystem to ensure environmental sustainability and optimise

management objectives on a prescribed area of forest land. It can also be defined as applied forest ecology. In other words, Silviculture is the art and science of cultivating forest crops for all their diverse range of uses. It is concerned with the growing of trees just as agriculture is concerned with the growing of food crops in fields. (Savill *et al.*, 1997) A generally accepted definition is that it is the study of the life history and general characteristics of trees with particular reference to environmental factors. Silviculture in forest development is charged with the responsibility of evolving appropriate techniques required for the formation of new and regeneration of old forests.

2.3 Origin and Introduction of *Acacia senegal*

Acacia senegal is a multipurpose African tree of the family *Leguminosae* and subfamily *Mimosoideae*. The trees function as "keystone" species in the arid and semi-arid regions of Africa and middle East [14]. It is highly valued for centuries for gum, fodder and timber production [15]. Today *A. Senegal* is grown primarily for gum, but plays a secondary role in agricultural systems, restoring soil fertility and providing fuel and fodder. *Acacia* are regarded as a forgotten resource. About 170 species are native to Africa [1,16]. The species is indigenous to Africa countries which appear in different variation and morphological characteristics. *Acacia senegal* is found in Western, Central, Eastern and Southern Africa, Oman, Pakistan and India. It has been introduced into Egypt, Australia, Puerto Rico and the Virgin Island [17].

2.4 Climatic and Soil Requirements of *Acacia senegal*

Acacia senegal is very drought resistant [18]. It grows on sites with annual rainfall between 100-950mm, mainly between 300-400mm, and 5-11-month dry periods or more), dry wind, and sandstorms. Generally, it cannot withstand frost [1]. *Acacia Senegal* prefers coarse-textured soils such as fossil dunes, but it will also grow on slightly loamy sands and skeletal soils such as Lithosols [19]. Although generally soils are drained, there are exceptions: in Kayers region, south-Kordofan, East Sudan, *Acacia Senegal* grows on heavy clay soils with approximately 800mm annual precipitation. The best sites have pH of 5 to 8. The tree ranges from 100-1700m

elevation in the Sudan to 1950m around Nakuru in Kenya. Distribution of gum trees in Chad is located between latitudes 11⁰ and 17⁰ north and rainfall of between 150mm and 900 mm per year [20]. *Acacia spp* is seen as an agricultural asset because it produces green leaves, which are in full growth during the dry season which are dropped unusually during the rainy season. For this reason they provide shade to cattle and farmworkers during hot sun at the same time do not obstruct the light needed by the growing crops during the rainy season [1]. This makes it a good choice for agroforestry practices.

3. RESULTS

3.1 Propagation Methods in *Acacia senegal*

Propagation in *Acacia senegal* is achieved both by direct sowing using seeds planted in nursery and by vegetative means using cuttings, grafting, layering and aerial layering. In *Acacia* propagation by using seeds, they should be harvested before pods have dried for easy collection and to avoid insect attack. Seed is easily extracted by hand. Freshly extracted seed should immediately be dusted with an insecticide.

Furthermore, one of the most notable examples of direct sowing for raising plantations is *Acacia nilotica* and *A. Senegal* in the Sudan. But nearly all tropical plantations are planted rather than direct sown, and at maturity, clear felled and replanted rather than naturally regenerated [21]. Vegetative propagation is also possible in *Acacia senegal* as fragments taken from crown of mature trees that are at least 13 years old. cuttings can be taken from branches 50 cm long.

3.2 Seed Production and Nursery in *Acacia senegal*

Acacia seeds are dormant and hard and needs to be given some sort of pre-treatment for them to germinate faster. This has been reported by Mohammed et al. [22], that generally members of the *Leguminosae* family have hard seed coat and contain less moisture. Adequate and dependable supply of water is needed during nursery sittings for adequate growth of the seedlings. Plants must be watered morning and evening daily by hand or irrigation system [23].



Plate 1. Supervision of planted seedlings in a nursery

Source: NAGAPPEN (2002)



Plate 2. Seed of *Acacia Senegal*

Source: Mohammed et al. [22]

A. *Senegal* is usually raised in the nursery in polyethylene pots, 2-4 seeds per pot, thinned to one seedling after 4-6 weeks. Direct seeding (5-8 seeds in 30 times 30 cm pits or larger) can also be used [24]. The planting stock can be kept in a wetted jute cloth to make the cuttings at the time of establishing nursery [18]. treatment by dilute Tetraoxo sulphate (vi) acid also greatly enhanced germination in *Acacia senegal* as reported by [25]. Experiment was conducted to assess different seedlings accessions of *Acacia senegal* in Bauchi Nigeria, eight different seedlings four from Borno and 4 from Yobe were collected. Seedling no eight from Yobe had the desirable seedlings characters namely fast growth habit highest seedling girth and could be transplanted within 57 days of sowing [26]. Shade are constructed to protect seedlings from direct sunlight for two to three weeks after pricking out [23]. In south Western highlands of Saudi Arabia *Acacia* species takes a significant portion of the floral composition and is significant for beekeeping production in the country. Despite the degraded condition of the vegetation of the region. It was pointed out that planting nursery grown seedlings is one of the best methods for successful restoration of the degraded land. Apart from environmental and climate change mitigation it also help in evaluating nursery level performances of selected native *Acacia* species in apicultural land restoration [27].

3.3 Wood and Other Uses of *Acacia senegal*

Gum: *Acacia Senegal* and its close relatives are the defined source of commercial gum Arabic food purposes. *A. Senegal* produces the only

Acacia gum evaluated toxicologically as a safe food additive [28]. Result of an experiment conducted by E. Muita et al shows that gum arabic from *A. senegal* var *kerensis* can be used as a stabilizer in low-fat yoghurt formulations and this increases consumer acceptability [29]. The phytochemical analysis of Gum Arabic have been carried out by Ibrahim and co-workers and was found to be rich and valuable in the food and other commercial industries [30]. The gum from other *Acacia* species (*A. seyal* etc.) is available commercially as gum tahla (approx. 10% of acacia gum marketed) for technological applications. Gum Arabic has been used for at least 4,000 years by local people for preparation in food, in human and veterinary medicine, in crafts and as a cosmetic [31]. Today, gum Arabic application is manifold. Formerly the international trade market largely absorbed all gum available. In Armania a trial was conducted of the coppicing ability of *Acacia nilotica* in order to cover the deficit for the high demand for biomass which is around 90% of the energy consumed. The specie coppiced well and is a medium to fast growing in comparison to other indigenous species. though recently international demand has declined together with gum prices [32]. *Acacia senegal* is the only tree that produces Grade 1 (Highest quality) gum Arabic. Sudan is the leading exporter of grade 1 gum Arabic. Nigeria has less than 5% of the World's gum Arabic market. Sudan and Chad makeup about 95% [17]. About 70 to 85 of the World total consumption of gum Arabic is produced in the Sudan [33]. World production of gum arabic potential is estimated at around 60,000 tons per annum, out of which 50% or more originate from Sudan [34]. "Gum Arabic is used in the food industry as a flavour fixative and emulsifier; stabilizer in frozen dairy products, for its viscosity and adhesive properties in bakery products, and as a foam stabilizer and clouding agent in beer" [35].

“In pharmaceuticals, it is used as a stabilizer for emulsions, binder and coatings for tablets, and as an ingredient in cough drops and syrups” [36]. “A soothing and softening agent, gum Arabic is extensively employed in folk medicines. Among many other uses it is used internally for coughs, diarrhoea, dysentery, hemorrhage, and externally to cover inflamed areas. Gum arabic is a dietary supplement for improving nutrition of type 2 diabetic patients, it exhibit an excellent property on improving their poor glycemic control” [37]. “Gum Arabic is used in cosmetics as an adhesive for facial masks and powders, and to give a smooth fell to lotions. Industrially, gum Arabic is applied as an adhesive, as a protective colloid and safeguarding agent for inks, sensitizer for lithographic plates, coating for special paper, sizing agent for cloth to give body to certain fabrics, and coating to prevent metal corrosion. Gum Arabic is used in the manufacture of matches and ceramic pottery” [38]. Many studies reported “the use of *acacia arabica* as an effective against variety of diseases including diabetes, skin disease and cancer. Fresh parts are considered as astringent, demulcent, aphrodisiac, anthelmintic, antimicrobial, antidiarrheal with good nutritional value in Indian traditional medicine system” [39,40]. Many studies reported “the use of *Acacia arabica* as an effective against variety of diseases including diabetes. Fresh parts are considered as astringent, demulcent, aphrodisiac, anthelmintic, antimicrobial, antidiarrheal with good nutritional value in Indian traditional medicine system” [41].

3.4 Tapping Techniques for Gum Arabic

The gum Arabic tree has erected rigid and strong stems and measures up to 13m in height. It is thorny and low branching. It requires moderate pruning of lower branches from the main stem up to 150 cm above ground level for easy access to the stem for tapping [42]. Tapping is an important management tool for enhancing gum productivity. The best time to tap a tree for high output is about 9:00 am and 5:00 pm [43]. Tappable girth size in gum Arabic lies between 25cm and 50cm at about 100cm above ground. The correct sign for tapping is after rains are over when about 50% of the leaves are shedded. This continues up to the next rain. Higher temperature encourages exudation of gum. exploitation can be natural through cracks on the stem or by the use of small axe or cutting knife [42].

There is a significant relationship between applied gum Arabic tapping techniques by using sunken and increasing gum yield and quality at ($p=0.05$) [44]. The gum is obtained as dry exudate from the stem and branches of the tree after the plant is injured. The nature of the injury can be organized and controlled (tapping). Injuries could be natural cracks/stem burst or injury as a result of pruning or cutting a stem or branches for fencing or animal fodder [24,31]. Research was conducted to find out whether there is a difference in yield between tapped and untapped *Acacia senegal* trees. Results show that tapping increases the gum Arabic yield by 77.42% [45]. Tapping date significantly ($P < 0.05$) produces a difference in *Acacia senegal* gum yield (g/picking) giving higher yield when tapped in the 15th of October and giving out lesser yield when tapped on the 1st of September as reported by Ilu and coworkers [46].

3.4.1 Wood

“*Acacia Senegal* is locally valued for fuel wood and charcoal although biomass yield per unit land area is not sufficient to plant *A. Senegal* purely for fuel wood. Wood is used in local construction for poles and fence posts, the light-coloured wood for tool handles and dark heartwood for weaver’s shuttles. Strong ropes are made from the bark fibres of the tree’s long surface roots” [47].

3.4.2 Food and fodder

“Dried and preserved seeds of *A. Senegal* are used by people as vegetables. The foliage and pods are browsed by sheep, goats, camels, impala, and giraffe. Leaves contain 10% - 13% digestible protein and 0.12% - 0.15% phosphorus, while the pods contain 15% digestible protein and 0.12% - 0.14% Phosphorus” [48].

3.4.3 Dune stabilization

Acacia Senegal is important for desertification control through sand dune stabilization and wind breaks.

3.4.4 Pests and diseases of *Acacia senegal*

One of the commonest pests of *A. senegal* is the buffalo tree hopper (*Stictocephala babilus*) which may destroy seed crops. Other pest includes the spiders (*Cyclops* sp.) which can smother young growing apices. Larval stage of *coleopterans* (*bruchids*), *Lepidoptera*, and

Hymenoptera damages the seed. Locusts (*Aecidium melanorhodon*) can defoliate vast areas overnight. *Acacia senegal* is also attacked by the fungi cladosporium herbarium, *Fusarium* sp, *Ravenelia acaciae-senegalae* and *R. acaciocola* [48].

3.4.5 *Acacia senegal* for agroforestry

It is a multipurpose nitrogen-fixing legume. They maintain and improve soil productivity and is the reason why they are adopted in agroforestry systems. Farmers cut trees and sell that are more than 20 years old which resulted in yield increase [49].

Acacia Senegal is grown in agro forestry systems especially in the Sudan and Uganda in "gum gardens" for gum as well as to restore soil fertility [14,50,51]. "Five-year-old trees are ready for tapping, and production peaks between 7 and 15 years. In Sudan, a traditional bush-fallow system is followed with a 20-year rotation during which time *Acacia senegal* is grown for 15 years. Agricultural crops are grown for five years (sesame, millet, sorghum, groundnuts), followed by five years with young, unproductive *A. Senegal* trees, which latter produce gum during the last 10 years of the rotation" [51]. Corresponding to this rotation, $\frac{1}{4}$ of the land is kept in agricultural crops, $\frac{1}{4}$ in unproductive young trees, and $\frac{1}{2}$ in productive trees. Controlled grazing is practiced after the trees have reached age four and under productive trees after the gum has been harvested. Wild trees are harvested during the dry season for gum exuded from cracks in the bark. Strict protection from fire and livestock grazing, and efficient control of weed competition during at least the first two years is important to seedling survival. Minimum spacing for block planting is 4 times 4m. At 10 times 10m spacing, agriculture inter cropping is possible, for example interplanting with millet, beans, or groundnuts.

4. CONCLUSION

In conclusion, Yobe is an agrarian state known with marginal agricultural soil as a result of population expansion, land degradation and desert encroachment. It may here be seen that the northern state of Yobe is an agrarian state, as a result of population expansion the marginal agricultural soil is over cultivated and trees were cleared massively for fuel wood all resulting in land degradation and desert encroachment. Poverty index is very high in the state with over

70%, one of the highest in the country. Domestication of *Acacia* species (silviculture and silvics) is pointed by this paper as a way forward for improving the economy (income generation of the masses) and improving the harsh Sahelian climate of the state. *Acacia senegal* is a legume and a nitrogen fixer. *Acacia* produce seeds in a pod. *Acacia* is a plant of dry tropical climate, *Acacia Senegal* produce gum Arabic and fuel wood, *Acacia* is mostly grown in underdeveloped African nations and some developing Asian countries. *Acacia* is planted through seed with little chance of natural regeneration because of grazing animals. The tree can also be propagated by vegetative means.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hayward B, Wales Z, ES. The *Acacia* tree: A sustainable resource for Africa. DFID Forestry Research Programme; 2004.
2. Hayward B. The *Acacia* tree: A sustainable resource for Africa. DFID Forestry Research Programme; 2004.
3. Diatta O, Sarr MS, Hansen JK, Diallo AM, Nielsen LR, Ræbild A, et al. Survival and growth of *Acacia senegal* (L.) Wild. (*Senegalia senegal* (L.) Britton) provenances depend on the rainfall at the site of origin. *Ann Forest Sci.* 2021;78(4):1-16. DOI: 10.1007/s13595-021-01098-5
4. Ibrahim AA. MDGs and poverty in Yobe State. *Int J Sustain Dev.* 2012;5(11):1-14.
5. Jaiyeola AO, Bayat A. Assessment of trends in income poverty in Nigeria from 2010-2013: An analysis based on the Nigeria General Household survey. *J Pover.* 2019;00(00):1-18. DOI: 10.1080/10875549.2019.1668900
6. Jaiyeola AO, Choga I. Assessment of poverty incidence in Northern Nigeria. *J Pover.* 2020;00(00):1-18. DOI: 10.1080/10875549.2020.1783424
7. Edoumiekumo SG, Karimo TM, Tombofa SS. Income poverty in Nigeria: incidence, gap, severity and correlates. *Am J Hum Soc Sci.* 2014;2(1):1-9. DOI: 10.11634/232907811402499
8. Hunger AA. A Cost of the Diet analysis in Yobe State, Nigeria; 2019.

9. Ali B, Saadun N, Alias MA, Norizah K. Assessment of the fuelwood value chain in Yobe, Nigeria. IOP Conf S Earth Environ Sci. 2019;268(1):0-8.
DOI: 10.1088/1755-1315/268/1/012063
10. Abbas AI, Ogiri IA. Political economy of financing education at sub national level in Nigeria: trends, achievements, gaps and challenges. Int J Dev Sustain. 2017;6(12):2145-62.
11. Statistics NB. of. 2020;2019 Poverty and Inequality in Nigeria : Executive summary.
12. Yk M, Kj I, Au, Ah J, G. Kd, S, & aa, L. Analysis of profitability of neem seed marketing in Yobe. International Journal of Wood Science, Technology and Forestry. 2022;7(2):158-67.
13. Forestry OM. of N.R. and. 2015. Forest management guide to silviculture in the Great Lakes-St. Lawrence and boreal forests of Ontario. Queen's printer for Ontario.
14. Agea JG, Obua J, Namiremhe S, Buyim M, Waiswa D. Ecology and Conservation of *Acacia senegal* in the Rangelands of Luwero and Nakasongola Districts. Uganda. J Agric Sci. 2005;11(1):40-6.
15. Jibo AU. Influence of Drought on *Acacia senegal* (L.) Willd: gum Yield within a Soil Moisture Gradient in North Eastern Nigeria. J Appl Sci Environ Manag. 2021;25(7):1201-6.
DOI: 10.4314/jasem.v25i7.15
16. Githae EW, Gachene CKK, Njoka JT, Omondi SF. Nitrogen fixation by natural populations of *acacia senegal* in the drylands of Kenya using 15-N natural abundance. Arid Land Res Manag. 2013;27(4):327-36.
DOI: 10.1080/15324982.2013.784377
17. Harrison G, Ryan Roberge ARD, I. Best Practices, Best Markets Training the Nigerian Gum Arabic Producer and small Trader; 2002.
18. Danthu P, Leblanc JM, Badji S, Colonna JP. Vegetative propagation studies of gum arabic trees .2 The vegetative propagation of adult *Acacia senegal*. Agroforest Syst. 1992;19(1):15-25.
DOI: 10.1007/BF00130091
19. Qadir MSZAC, SA. *Acacia senegal* (L.). Vegetatio. 1973;27(1-3):131-62.
20. Daakkreo G, Adamou I, Valery NN, Djangrang M. Comparison of carbon stocks in *Acacia senegal* and *Acacia seyal* stands in Chari – Baguirmi region (Chad). Int J Adv Res Biol Sci. 2019;6(2):195-203.
DOI: 10.22192/ijarbs
21. Julian E. Plantation forestry in the tropics. Oxford: Clarendon press; 1996.
22. Mohammed E, Mustafa A, Al-kamali HH, Habeeballa RS. Taxonomic characters of some Sudanese *Acacia*. Eur J Adv Res Biol Life Sci. 2017;5(1):9-18.
23. Shettima ML, Mohammed GA, Badawi HL, Kofan A, Edet BE, Marte ML, et al. Comparative Techniques of Raising Seedlings of *Acacia seyal* in the Arid Zone of Borno State, North – Eastern Nigeria. International J Environ Prot Policy. 2017;5(6–1):40-9.
DOI: 10.11648/j.ijepp.s.2017050601.16
24. Alkali UU. silvicultural practices of gum arabic tree (*Acacia senegal*): A review Alkali, U.U. Journal of Environmental Issues and Agriculture in Developing Countries. 2010;2(3):140-5.
25. Bosah KO, BO. Germination response of *Acacia senegal*.pdf. Agric J. 2007;2(6):681-4.
26. Ibrahim N, Gani AM, Abdul SD et al. Performance of Different Seedlings Accessions of *Acacia senegal* (L.) willed in Bauchi, Nigeria. Int J Sci Eng Res. 2014;5(6):5-7.
27. Al-ghamdi AA, Tadesse Y, Adgaba N. Evaluation of major *Acacia* species in the nursery towards apicultural landscape restoration around Southwestern Saudi Arabia. Saudi J Biol Sci. 2020;27(12):3385-9.
DOI: 10.1016/j.sjbs.2020.09.002, PMID 33304146.
28. Anderson DM. NFT gums: Ancient and modern commercial products; 1991. Available:<http://vl.winrock.org/forestry/factpub/factsh/aSenegal.html>
29. Muita E, Mahungu SM, Chikamai BN, Johnson K. Evaluation of gum arabic from *Acacia senegal* var *kerensis* and *Acacia senegal* var *senegal* as a Stabilizer in Low-fat Yoghurt. Int J Food Stud. 2020;9(414):110-24.
30. Gashua IB, Ukekpe US, Yusuf I. Biophysical investigation of plant exudate of *Acacia senegal* (L) Wild. from Sudano-savannah ecological zone of Nigeria. Int J Adv Res. 2013;1(4):228-32.
31. Mokwunye MUB, AO. Restoring Nigeria's lead in gum arabic Production: Prospects and Challenges. 2010;2(4);.
32. Gessesse AT, Gezahegn TT, Wolle HS. Study on coppice management of *Acacia nilotica* Tree for better woody biomass

- production. Agric For Fish. 2015;4(3):138-41.
DOI: 10.11648/j.aff.20150403.19
33. Mustafa AF, Elamin KH, AAS, WM. Water use efficiency studies of *Acacia senegal* (L.) Willed provenances in Sudan; 1965.
 34. Mujawamariya G. Economics of the gum arabic value chain in Senegal. Wageningen University; 2012.
 35. Studer MJ, JN, M. Commodities at a glance Special issue on gum arabic. United Nations Conference on Trade and Development, 8, 1283; 2018.
 36. Hamouda Y. Factors affecting the quality of *Acacia senegal* gums. University of Chester; 2017.
 37. Babiker R, Elmusharaf K, Keogh MB, Banaga ASI, Saeed AM. Metabolic effects of gum arabic (*Acacia senegal*) in patients with type 2 diabetes mellitus (T2DM): Randomized, placebo controlled double blind trial. Funct Foods Health Dis. 2017;7(3):219-31.
DOI: 10.31989/ffhd.v7i3.325
 38. NFTA. *Acacia senegal* – gum Tree with Promise for Agroforestry. A quick guide to useful nitrogen fixing trees from around the world; 1991.
 39. Abduljawad EA. Review of some evidenced medicinal activities of *Acacia nilotica*. Arch Pharm Pract. 2020;11(4):20-5.
 40. Amadou I, Soulé M, Salé A. An overview on the importance of *Acacia nilotica* (L.) wild. Ex Del.: a review. Asian J Res Agric For. 2020;5(3):12-8.
DOI: 10.9734/AJRAF/2020/v5i330085
 41. Rajvaidhya S, Nagori BP, Singh GK, Dubey BK, Desai P, Alok S, et al. A review on *Acacia arabica* – an Indian medicinal plant. Int J Pharm Sci Res. 2012;3(07):1995-2005.
 42. Aghughu O, Wuranti V. A review of tapping techniques of gum arabic (*Acacia senegal* (L.) wild) in the sahel region of north-eastern nigeriae; 2005.
 43. Yau IA, Babura MS, Bashir BH, Ahmed S. Effect of weeding status and tapping date on the yield of gum arabic (*Acacia senegal*) in Jigawa State, Nigeria. Bayero J Pure Appl Sci. 2016;9(2):263-6.
 44. Hamad AAEH, Merghani ESA, Abdelmagid TD, Amien TM, Eltahir MES. Situation analysis of gum arabic production and practices in North Kordofan, Sudan. Asian J Res Biosci. 2021;3(1):38-43.
 45. Wekesa C, Makenzi P, Chikamai BN, Lelon JK, Luvanda AM, Muga M. Gum arabic yield in different varieties of *Acacia senegal* (L.) Willed in Kenya. Afr J Plant Sci. 2009;3(11):263-76.
 46. Ilu KJ, Salami KD, Mohammed KY et al. Influence of tapping dates on the yield of *Acacia senegal* (L.) wild at two different. Fudma Journal of Sciences. 2020;4(1):246-9.
 47. Mohammed MH, Rahman HMA, Khatir AA. Provenance variation in seed morphological characteristics, germination, and seedling growth of *Acacia senegal* high-yielding gum trees. Forestist. 2022;72(3):233-40.
DOI: 10.5152/forestist.2022.21017
 48. Orwa C, Mutua KR, Jamnadass R, SA. *Acacia senegal* (L.) Willed. Agrofor Database 4.0. 2009;0.
 49. Bargali SS, Bargali K. *Acacia nilotica*-based traditional agroforestry system: Effect on paddy crop and management. Curr Sci. 2009;96(4):1-8.
 50. Mohamed AG. Improvement of traditional *Acacia senegal* Agroforestry: Ecophysiological characteristics as indicators for tree-crop interaction on sandy soil in western Sudan (Issue March). University of Helsinki; 2005.
 51. Temgoua LF, Momo Solefack MC, PO, GTB. Floristic structure and carbon sequestration potential of *Acacia senegal* (L.) willed. (fabaceae) improved fallows in far north Region of Cameroon. Eur J Agric For Res. 2018;6(2):29-41.

© 2023 Mohammed et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/101253>