



Effectiveness of Cognitive Behavioral Therapy for Post-partum Depression in Low and Middle-income Countries: A Systematic Review

Abimbola Eunice Arisoyin ^{a*}, Mujeeb Adedeji Salawu ^b,
Ibrahim Lanre Folorunsho ^a, Kehinde Tolulope Olaleye ^c,
Adebola Yewande Afolayan ^b, Damilola Abiola Adeyemo ^d,
Rheiner Nnenna Mbaezue ^e, Chidalu Nkeyonyere Ibeneme ^f,
Jashanpreet Singh Ballagan ^g, Shilpa Rai ^h,
Olasumbo Elizabeth Fagbenle ^g,
Ifeyinwa Monica Oraekwute ⁱ,
Stefany Carolina Lopez Pantoja ^j, Omolola Okunromade ^k
and Mark Majid Haddad ^l

^a College of Medicine, University of Lagos, Nigeria.

^b College of Health Science, University of Ilorin, Nigeria.

^c Obafemi Awolowo University, Ile Ife, Nigeria.

^d Ladoke Akintola University of Technology, Nigeria.

^e Department of Health, Cape Town, South Africa.

^f College of Medicine, Abia State University, Nigeria.

^g Windsor University School of Medicine, Saint Kitts and Nevis.

^h Isra University, Pakistan.

ⁱ O.O. Bogomolets National Medical University, Kiev, Ukraine.

^j Pontificia Universidad Catolica del Ecuador, Ecuador.

^k Georgia Southern University, USA.

^l Caribbean Medical School, Curaçao.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMMR/2023/v35i185119

*Corresponding author: E-mail: abimbolaarisoyin@gmail.com;

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102313>

Review Article

Received: 05/05/2023

Accepted: 07/07/2023

Published: 14/07/2023

ABSTRACT

Background: Post-partum depression (PPD) is a prevalent psychological condition affecting 10-20% of women following childbirth, with higher rates in low and middle-income countries (LMICs) due to limited resources, poor access to mental health care, and sociocultural stressors. Cognitive Behavioral Therapy (CBT) is an evidence-based intervention for treating PPD in high-income countries, but its effectiveness in LMICs remains underexplored.

Aim: This systematic review aims to synthesize evidence on CBT's effectiveness for post-partum women in LMICs, examine cultural and contextual factors, and provide recommendations for culturally-sensitive interventions.

Methods: Following PRISMA Statement 2020 guidelines, a systematic search was conducted in electronic databases such as PubMed, PsycINFO, and Web of Science to identify studies evaluating CBT's effectiveness for post-partum women in LMICs. A narrative synthesis was used to analyze and synthesize findings, considering the heterogeneity in study designs, populations, and outcome measures.

Results: The review included five studies, all of which reported a more significant decrease in Edinburgh Postnatal Depression Scale (EPDS) scores for the CBT intervention group compared to the control group. The studies varied in terms of populations, settings, and CBT delivery modes, with telephonic CBT, in-hospital CBT, and in-person CBT being utilized.

Conclusion: The findings from this systematic review suggest that CBT may be effective in reducing depressive symptoms among post-partum women in LMICs. Further research is needed to strengthen the evidence base, refine intervention strategies, and develop more effective, accessible, and culturally-appropriate interventions for PPD in LMICs. Addressing these recommendations will contribute to better mental health outcomes for post-partum women and their families in developing countries.

Keywords: Postpartum; depression; CBT; psychiatry; LMIC; developing country.

1. INTRODUCTION

Post-partum depression (PPD) is the most common psychological condition following childbirth, affecting approximately 10-20% of women during pregnancy or within the first 12 months post-partum [1]. PPD is characterized by a range of emotional, behavioral, and cognitive symptoms, including persistent sadness, anxiety, irritability, and difficulty bonding with the newborn [2]. If left untreated, PPD may have detrimental effects on the social and cognitive health of spouses, infants, and children, leading to long-term consequences for family well-being and child development [3]. The prevalence of PPD is considerably higher in low and middle-income countries (LMICs) compared to high-income

countries due to factors such as limited resources, poor access to mental health care, and various sociocultural stressors [4]. As a result, there is an urgent need to identify and implement effective interventions to address PPD in these contexts, where the burden of the condition is disproportionately high.

Cognitive Behavioral Therapy (CBT) is a well-established, evidence-based psychological intervention that has demonstrated effectiveness in treating post-partum depression (PPD) in high-income countries [5]. It is a structured, short-term therapeutic approach that aims to help individuals identify and modify maladaptive thought patterns and behaviors that contribute to emotional distress and psychological problems

[6]. By addressing these cognitive and behavioral factors, CBT can lead to significant improvements in emotional well-being and overall mental health. In the context of PPD, CBT typically involves working with the individual to recognize and challenge negative thoughts and beliefs associated with their experience of motherhood, relationships, and self-worth [7,8]. Through a combination of psychoeducation, cognitive restructuring, and behavioral activation, women are encouraged to develop healthier and more balanced thought patterns and coping strategies, which can ultimately reduce depressive symptoms and enhance their ability to bond with their newborns [9]. Despite the documented success of CBT in high-income countries, there is limited research exploring its effectiveness and adaptability for PPD in LMICs [10]. This knowledge gap is particularly concerning given the higher prevalence of PPD in LMICs, where women often face additional challenges such as limited access to mental health care, inadequate social support, and unique sociocultural stressors.

Given the significant impact of PPD on maternal and child health, particularly in LMICs, the primary aim of this systematic review is to synthesize the existing evidence on the effectiveness of CBT for post-partum women in developing countries. The specific objectives of this review are to identify and analyze studies evaluating the effectiveness of CBT for PPD in LMICs, to examine the cultural and contextual factors that may influence the effectiveness of CBT in LMICs, and to provide recommendations for the development of culturally-sensitive and contextually-appropriate CBT interventions for PPD in LMICs. By addressing these objectives, this systematic review will contribute to a better understanding of the potential benefits and challenges of implementing CBT for PPD in LMICs, ultimately informing the development of more effective and accessible mental health care interventions for this vulnerable population.

2. METHODS

Adhering to PRISMA Statement 2020 guidelines [11], a systematic search was conducted to identify studies evaluating the effectiveness of CBT for post-partum women in LMICs. The search strategy included a combination of relevant keywords and MeSH terms such as cognitive-behavioral therapy, post-partum women, developing countries, and low and middle-income countries. Electronic databases,

including PubMed, PsycINFO, and Web of Science, were searched from inception to April 13, 2023. The PubMed search string is as follows: ("Cognitive Behavioral Therapy"[Mesh]) AND ("Postpartum Period"[Mesh]) AND ("Developing Countries"[Mesh]) AND ("Income"[Mesh]). Additional MeSH terms and keywords were adjusted. Reference lists of eligible articles and relevant reviews were also screened to identify any additional studies.

Two independent reviewers screened the titles and abstracts of identified articles for eligibility. The full texts of potentially eligible articles were then assessed to determine their inclusion in the review. Any discrepancies were resolved through discussion and consensus between the reviewers. The following information was extracted from each included study using a standardized data extraction form: author, year, country, study type, the primary mode of CBT delivery, sample size (interventional group and standard group), age (interventional group and standard group) as mean (SD), and Edinburgh Postnatal Depression Scale (EPDS) scores at follow-up compared to baseline (intervention group and control group).

Due to the heterogeneity in study designs, populations, and outcome measures, a narrative synthesis approach was employed to analyze and synthesize the findings. This method allowed for a more comprehensive understanding of the effectiveness of CBT in LMICs for post-partum women while taking into account the diverse contexts and characteristics of the included studies. The narrative synthesis was structured around the following themes: characteristics of the included studies, the primary mode of CBT delivery, and effects of CBT on EPDS scores at follow-up compared to baseline.

3. RESULTS

The study selection process for this systematic review followed the PRISMA guidelines, intending to identify relevant literature on the effectiveness of CBT for post-partum women in LMICs. A total of 96 records were initially identified through database searches, with 37 from PubMed, 21 from PsycINFO, and 33 from Web of Science. Additionally, five studies were identified through other sources, such as reference lists of eligible articles and relevant reviews. After removing duplicates, 71 records remained for screening.

During the screening phase, titles and abstracts of the 71 records were carefully examined, leading to the exclusion of 51 records that did not meet the inclusion criteria. Consequently, 20 full-text articles were assessed for eligibility. After a thorough evaluation, 15 of these articles were excluded for reasons such as not being specific to LMICs, not evaluating CBT, or not targeting post-partum women. Ultimately, a total of 5 studies were deemed eligible and were included in this systematic review, providing valuable insights into the effectiveness of CBT for post-partum women in developing countries (Fig. 1).

Cognitive Behavioral Therapy (CBT) in various countries and through different modes of delivery. In a randomized controlled trial conducted by Chungu et al. (2017) in Zambia, telephonic CBT was administered to 32 participants, and the results showed a decrease of 6.69 points (SD=3.44) in EPDS scores compared to the control group's decrease of 3.97 points (SD=3.53). Similarly, Hou et al. [13] conducted a controlled trial in China, where in-hospital CBT was provided, resulting in a decrease of 3.4 points (SD=1.9) in EPDS scores for the intervention group, compared to the control group's decrease of 1.5 points (SD=1.6).

Table 1 summarizes the findings from five distinct trials that examined the effects of




Fig. 1. PRISMA flowchart depicting the study selection process

Table 1. Characteristics of the included studies

Author, Year	Country	Study type	Primary mode of CBT delivery	N (IG)	N (SG)	Age (IG) as mean (SD)	Age (SG) as mean (SD)	EPDS scores at follow-up compared to baseline (IG)	EPDS scores at follow-up compared to baseline (SG)	Study Conclusions
Chungu et al. [12]	Zambia	Randomized Controlled Trial	Telephonic	32	32	15-35		-6.69 (3.44)	-3.97 (3.53)	When combined with pharmacotherapy, CBT improved mental health in young mothers with PPD, highlighting the need for greater CBT awareness among health professionals and mothers
Hou et al. [13]	China	Controlled trial	In-hospital	104	109	28(3)	28 (4)	-3.4 (1.9)	-1.5 (1.6)	CBT and SFT significantly reduced depression scores and improved sleep quality in mothers with mild to moderate postpartum depression
Fathi-Ashtiani et al. [14]	Iran	Randomized Controlled Trial	In-hospital	64	71	25(3.58)	26 (3.82)	-4.28 (6.87)	-1.04 (5.81)	Cognitive-behavioral therapy improved PPD and self-esteem in at-risk Iranian mothers
Ngai et al. [15]	China	Randomized Controlled Trial	Telephonic	197	200	31.1(3.8)	30.4 (4.4)	-4.14 (2)	0.96 (2.1)	Telephone-based CBT significantly reduced depressive symptoms in minor depressed mothers at 6 weeks and 6 months postpartum
Hussain et al. [16]	Pakistan	Randomized Controlled Trial	In-person	123	124	28.20 (5.47)	27.26 (5.5)	-5.6 (2.7)	-1.99 (3.08)	Learning through Play Plus, combined with CBT, reduced maternal depression and improved mothers' child development knowledge

Abbreviations: CBT: Cognitive behavioral therapy; EPDS: Edinburgh Postnatal Depression Scale; IG: Interventional group; PPD: Postpartum depression; SFT: Systemic family therapy; SG: Standard group; SD: Standard deviation.

Fathi-Ashtiani et al. [14] performed a randomized controlled trial with in-hospital CBT in Iran. They observed a decrease of 4.28 points (SD=6.87) in EPDS scores for the intervention group as opposed to the control group's drop of 1.04 points (SD=5.81). Ngai et al. [15] conducted another randomized controlled trial in China, where telephonic CBT led to a decrease of 4.14 points (SD=2) in EPDS scores for the intervention group and an increase of 0.96 points (SD=2.1) for the control group. Lastly, Hussain et al. [16] implemented a randomized controlled trial in Pakistan with in-person CBT, resulting in a decrease of 5.6 points (SD=2.7) in EPDS scores for the intervention group and a reduction of 1.99 points (SD=3.08) for the control group.

All of the studies reported a more significant decrease in EPDS scores for the intervention group compared to the control group, highlighting the potential effectiveness of CBT in the respective populations.

4. DISCUSSION

The findings of the five included studies in this systematic review suggest that Cognitive Behavioral Therapy (CBT) may be an effective intervention for reducing depressive symptoms among post-partum women in low and middle-income countries (LMICs). However, it is essential to discuss the studies' characteristics and outcomes in more detail to better understand the factors contributing to the intervention's effectiveness.

In the study by Chungu et al. [12] in Zambia, telephonic CBT was used, and the intervention group showed a more substantial decrease in EPDS scores compared to the control group. This mode of delivery might be particularly beneficial in LMICs, where access to mental health care services can be limited. Telephonic CBT can overcome geographical barriers and reduce travel-related expenses for participants, potentially increasing the intervention's accessibility and affordability.

Hou et al. [13] and Fathi-Ashtiani et al. [14] both used in-hospital CBT in China and Iran, respectively. These studies also reported significant reductions in EPDS scores for the intervention group compared to the control group. Delivering CBT in a hospital setting might enhance the integration of mental health care into existing maternal health services, facilitating early identification and intervention for women at risk of post-partum depression.

The study by Ngai et al. [15], conducted in China, employed telephonic CBT and demonstrated a decrease in EPDS scores for the intervention group. However, the control group's scores slightly increased, which might indicate that the standard care provided in the study setting was insufficient in addressing post-partum depression symptoms. This highlights the importance of evaluating and improving the quality of mental health care services available to post-partum women in LMICs.

Hussain et al. [16] implemented a randomized controlled trial in Pakistan with in-person CBT, which significantly reduced EPDS scores for the intervention group compared to the control group. In-person CBT allows for more personalized and tailored interventions, taking into account each individual's unique needs and circumstances.

It is important to acknowledge the limitations of this review. The small number of included studies limits the generalizability of the findings. The heterogeneity in study designs, populations, and outcome measures precluded the use of meta-analysis, necessitating a narrative synthesis approach. The small sample sizes in some of the included studies might limit the power to detect significant differences between intervention and control groups. Furthermore, the studies included in this review varied in terms of intervention duration, therapist qualifications, and follow-up periods, making it challenging to compare the findings directly.

Based on the findings and limitations of this review, the following recommendations are proposed for future research and practice:

- Conduct more rigorous and well-powered studies to further establish the efficacy of CBT for post-partum depression in LMICs [17].
- Assess the long-term effects of CBT on post-partum depression, as well as potential impacts on maternal functioning, infant development, and family relationships [18].
- Investigate the cost-effectiveness of different modes of CBT delivery, such as telephonic, in-person, or group-based interventions, to inform resource allocation decisions in LMICs [19].
- Explore the cultural adaptation of CBT to ensure its acceptability, relevance, and effectiveness in various cultural and socio-economic contexts [20].

- Examine the potential role of non-specialist or community health workers in delivering CBT interventions, as a strategy to improve access to mental health care in resource-limited settings [21].

5. CONCLUSION

This systematic review provides key evidence for the effectiveness of CBT in reducing depressive symptoms among post-partum women in LMICs. Despite the limitations, the findings have important implications for policymakers, clinicians, and researchers working to improve maternal mental health in developing countries. Further research is required to strengthen the evidence base, refine intervention strategies, and ultimately improve the mental health outcomes for post-partum women and their families in LMICs. By addressing the recommendations outlined above, the global health community can work towards developing and implementing more effective, accessible, and culturally-appropriate interventions for post-partum depression in LMICs.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kim THM, Connolly JA, Tamim H. The effect of social support around pregnancy on post-partum depression among Canadian teen mothers and adult mothers in the maternity experiences survey. *BMC Pregnancy Childbirth*. 2014;14(1):1–9.
2. Rai S, Pathak A, Sharma I. Post-partum psychiatric disorders: Early diagnosis and management. *Indian J Psychiatry*. 2015;57(Suppl 2):S216.
3. Johansson M, Benderix Y, Svensson I. Mothers' and fathers' lived experiences of post-partum depression and parental stress after childbirth: a qualitative study. *Int J Qual Stud Health Well-Being*. 2020;15(1):1722564.
4. Parsons CE, Young KS, Rochat TJ, Kringelbach ML, Stein A. Postnatal depression and its effects on child development: a review of evidence from low-and middle-income countries. *Br Med Bull*. 2012;101(1):57.
5. Jannati N, Mazhari S, Ahmadian L, Mirzaee M. Effectiveness of an app-based cognitive behavioral therapy program for post-partum depression in primary care: A randomized controlled trial. *Int J Med Inform*. 2020;141:104145.
6. McCraty R, Tomasino D. Emotional stress, positive emotions, and psychophysiological coherence. *Stress Heal Dis*. 2006;342–65.
7. Whelan A, Haywood P, Galloway S. Low self-esteem: Group cognitive behaviour therapy. *Br J Learn Disabil*. 2007;35(2): 125–30.
8. Westbrook D, Kennerley H, Kirk J. An introduction to cognitive behaviour therapy: Skills and applications. Sage; 2011.
9. Cassin SE, Sockalingam S, Wnuk S, Strimas R, Royal S, Hawa R, et al. Cognitive behavioral therapy for bariatric surgery patients: preliminary evidence for feasibility, acceptability, and effectiveness. *Cogn Behav Pract*. 2013;20(4):529–43.
10. Poudyal A, Van Heerden A, Hagaman A, Maharjan SM, Byanjankar P, Subba P, et al. Wearable digital sensors to identify risks of post-partum depression and personalize psychological treatment for adolescent mothers: protocol for a mixed methods exploratory study in rural . *JMIR Res Protoc*. 2019;8(9): e14734.
11. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021 Mar;372:n71.
12. Chungu MH. The effects of cognitive behavioral therapy on the improvement of the health of youth mothers with post-partum depression at the University Teaching Hospital of Lusaka, Zambia. The University of Zambia; 2017.
13. Hou Y, Hu P, Zhang Y, Lu Q, Wang D, Yin L, et al. Cognitive behavioral therapy in combination with systemic family therapy improves mild to moderate post-partum depression. *Brazilian J Psychiatry*. 2014; 36:47–52.
14. Fathi-Ashtiani A, Ahmadi A, Ghobari-Bonab B, Azizi MP, Saheb-Alzamani SM. Randomized Trial of Psychological Interventions to Preventing Post-partum Depression among Iranian First-time Mothers. *Int J Prev Med*. 2015;6:109.

15. Ngai F-W, Wong PW-C, Leung K-Y, Chau P-H, Chung K-F. The effect of telephone-based cognitive-behavioral therapy on postnatal depression: a randomized controlled trial. *Psychother Psychosom.* 2015;84(5):294–303.
16. Husain N, Zulqernain F, Carter L-A, Chaudhry IB, Fatima B, Kiran T, et al. Treatment of maternal depression in urban slums of Karachi, Pakistan: a randomized controlled trial (RCT) of an integrated maternal psychological and early child development intervention. *Asian J Psychiatr.* 2017;29:63–70.
17. Islam A, Mozumder TA, Rahman T, Shatil T, Siddique A. Forced Displacement, Mental Health, and Child Development: Evidence from the Rohingya Refugees. Work Pap. 2021;
18. Cooper PJ, Murray L, Wilson A, Romaniuk H. Controlled trial of the short-and long-term effect of psychological treatment of post-partum depression: I. Impact on maternal mood. *Br J Psychiatry.* 2003; 182(5):412–9.
19. Ward C, Sanders MR, Gardner F, Mikton C, Dawes A. Preventing child maltreatment in low-and middle-income countries: Parent support programs have the potential to buffer the effects of poverty. *Child Abus Negl.* 2015;
20. Heim E, Kohrt BA. Cultural adaptation of scalable psychological interventions. *Clin Psychol Eur.* 2019;1(4):1–22.
21. Barnett ML, Gonzalez A, Miranda J, Chavira DA, Lau AS. Mobilizing community health workers to address mental health disparities for underserved populations: a systematic review. *Adm Policy Ment Heal Ment Heal Serv Res.* 2018; 45:195–211.

© 2023 Arisoyin et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/102313>