



## **Tooth Mobility Pattern in Periodontitis Patients with Diabetes Mellitus**

**Aarthi Muthukumar<sup>a</sup>, M. Jeevitha<sup>b\*</sup>, Selvaraj Jayaraman<sup>c</sup>  
and M. Naveen Kumar<sup>d</sup>**

<sup>a</sup> Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, India.

<sup>b</sup> Department of Periodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, India.

<sup>c</sup> Department of Biochemistry, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, India

<sup>d</sup> Department of Orthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i64B35326

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/80279>

**Original Research Article**

**Received 06 November 2021**  
**Accepted 29 December 2021**  
**Published 30 December 2021**

## **ABSTRACT**

**Background:** Tooth mobility may be a common problem among patients seeking dental treatment. Tooth mobility is defined as an extent of horizontal and vertical tooth displacement created by external forces, trauma and periodontal diseases.

**Aim:** The aim of the present study is to evaluate the pattern of tooth mobility in patients with diabetes mellitus.

Study design: Retrospective study

**Materials and methods:** The case sheet records ( DIAS data ) of tooth mobility in diabetic patients were extracted. A total of 1568 case sheets were analysed for the study. Age, gender and pattern of tooth mobility with respect to the region were collected and statistically analysed using SPSS statistical software. Descriptive statistics (percentage, mean, SD) and inferential test (Chi square test) were performed to determine the association between age, gender and region of tooth mobility in diabetes mellitus patients.

**Results:** Results showed that tooth mobility was more common among males when compared to females. According to age, patients between 41 to 50 years with diabetes mellitus were more affected by tooth mobility. Tooth mobility was more commonly seen in the mandibular anterior region.

**Conclusion:** The present study showed a possible relationship between diabetes mellitus and tooth mobility. Results showed that 80% of patients with diabetes mellitus had tooth mobility. Therefore, early diagnosis, preventive therapeutic measures and oral hygiene reinforcement is needed to prevent progression of periodontal disease leading to tooth mobility.

*Keywords: Diabetes mellitus; novel method; periodontitis; tooth mobility; periodontal disease.*

## 1. INTRODUCTION

Physiological tooth mobility is defined as a slight displacement of the clinical crown of the tooth that is allowed by the resilience of an intact and healthy periodontium, under the application of moderate force [1]. Tooth mobility can cause occlusal instability, improper mastication and impaired quality of life [2]. Assessment of tooth mobility is an important part of periodontal assessment, because tooth mobility is a major sign in diagnosis of periodontal disease [2,3].

Miller's tooth mobility index graded the tooth mobility from I to III [4]. The mechanisms of periodontitis mainly involved to cause tooth mobility are inflammatory disruption of periodontal tissues [5], widening of periodontal ligament, loss of attachment, horizontal or vertical loss of alveolar bone and trauma from occlusion [6]. Tooth mobility can occur as a result of secondary trauma as the destruction of tissue occurs in the presence of normal occlusal forces in teeth with weakened supporting tissues [7]. Factors that result in tooth mobility are abnormality in crown root ratio [7,8], shape and length of root, structural abnormality of tooth and periodontal disease [9,10]. The initial signs of tooth mobility as shared by patients are tenderness on mastication followed by pain or sudden displacement of tooth [11]. Displacement of anterior teeth will affect aesthetic appearance [12,13].

Chronic periodontitis is an advanced form of periodontal disease causing destruction of both soft tissue and hard tissue components of tooth supporting structures resulting in tooth mobility [14]. The adoption of proper and adequate steps in the management of tooth mobility will definitely help in increasing the longevity of teeth and preventing tooth loss [15]. The aim of the present study is to evaluate the pattern of tooth mobility in diabetic patients with periodontal disease.

Our team has extensive knowledge and research experience that has translate into high quality publications [16-35].

## 2. MATERIALS AND METHODS

The data comprising demographic details and tooth mobility in diabetes mellitus patients were collected from the duration of June 2019 to April 2020. A total of 1568 case sheets were analysed. The samples were collected by a simple random sampling method. Cross verification of data was done. FDI tooth numbering system was followed in the study. Incomplete datas was excluded. The analysis was done using SPSS version 19. Dependent variables are the number of mobile teeth and independent variables are age and gender. The data was statistically analysed using Chi-square test. The level of significance was set at 0.05.

## 3. RESULTS AND DISCUSSION

100 diabetic patients diagnosed with periodontal disease were studied. The study sample consisted of 48.5% females and 51.5% males. The proportion of males was greater than the females [Fig. 1]. Majority of the participants were in the age group between 51-60 years (31.3%) [Fig. 2]. 49% of the mobile teeth were present in both anterior and posterior regions, 30% were present in the anterior region and 20% of the mobile teeth were present in the posterior region. Thus the results showed that tooth mobility was more commonly seen in the anterior region [Fig. 3]. Mobile teeth were present in the mandibular arch (36.3%) than the maxillary arch (13%) [Fig. 4]. Majority of the patients had tooth mobility in the anterior region (30.3%) than other regions. Most of the male patients had mobile teeth in both regions (28%) compared to female patients (21%), however it was statistically not significant ( $p>0.05$ ) [Fig. 5]. The patients in the age group of 51-60 years present with a higher number of mobile teeth in both anterior and posterior


regions (31%) ( $p>0.05$ ) [Fig. 6]. Tooth mobility was more commonly seen in the mandibular arch than the maxillary arch and more common among males (51.5%) than females [Fig. 7]. Mobile teeth were observed in patients in the age group of 51-60 years more commonly seen in the mandibular region (36.6%) than the maxillary region. However, this is statistically not significant [Fig. 8].

Tooth mobility is a common problem among patients seeking dental health attention with the presenting complaints of mobile teeth, painful teeth and shaking teeth [36]. The high chance of tooth mobility in this study was related to chronic periodontitis [37,38]. Tooth mobility may be associated with various factors of periodontal disease [39] among patients in relation to age, gender, irregular dental check up, chronic medical condition and poor oral hygiene [40]. In this present study, males (51.5%) with mobile teeth and females (48.5%) with mobile teeth coincides with the previous study performed by Henry et al in 2017 [41] he habit of smoking was thought to be a reason behind the increased susceptibility of males with tooth mobility. The results of this study showed that the tooth mobility was highest in the fifth and sixth decades with a peak incidence of 51-60 years (31.3%). This study coincided with Azado et al in 2017 reported that tooth mobility to be one of the main symptoms of periodontal disease after the age of 40 years. The possible explanation for the difference is due to the effect of increasing age,


there could be a destruction of periodontium, loss of alveolar bone, loss of attachment and malnutrition [42].

Tooth mobility in the anterior region (30%) more than the posterior region (20%). This finding is similar to a study proposed by Muhlemann HR et al in 1960, he stated single rooted teeth have a higher tooth mobility measurement than the multirooted teeth [43] results of the present study showed tooth mobility in the mandibular arch (36%), maxillary arch (13.3%) and both maxillary and mandibular arch (50.5%). Tooth mobility was more commonly seen in the mandibular arch. This result contradicts the finding of Giannakoura et al in 2019 [14], which reported that the maxillary arch showed more tooth mobility than the mandibular arch. The possible explanation of the difference is occlusal trauma and primary tooth exfoliation [44].


In the present study gender and age was not significantly associated with tooth mobility. The present study is important to raise clinicians awareness in the appropriate way of managing tooth mobility. It is very important to properly diagnose the cause of a particular pattern of tooth mobility. The present study had geographic limitations, small sample size and also other confounding factors were not studied. The future scope of the study is to implement preventative therapeutic measures, emphasize oral hygiene in the mandibular anterior region and awareness programs for patients with periodontal diseases.




**Fig. 1.** Bar graph showing the gender distribution of patients. X-axis represents gender and Y-axis represents the percentage of patients. 51.5% of the patients are males (red) and 48.4% are females (blue)




**Fig. 2. Bar graph showing the distribution of the age group of the study participants. X axis represents the patient age group and Y axis represents the percentage of patients. Majority of the patients are in the age group of 51-60 years (31.3%)**




**Fig. 3. Bar graph showing the distribution of location of tooth mobility. X axis represents the location of the tooth mobility (anterior, posterior, both) and Y axis represents the percentage of patients with periodontal diseases. Mobile teeth were more commonly seen in the anterior region (30.3%)**




**Fig. 4.** Bar graph showing the distribution of the region of the tooth mobility. X axis represents the region of tooth mobility (maxillary, mandibular, both) and Y axis represents percentage of patients with periodontal diseases. Mobile teeth were more common in the mandibular arch (36.3%) than the maxillary arch (13.3%)




**Fig. 5.** Bar graph representing the association of gender and location of mobile teeth. X axis represents the location of the mobile teeth and Y axis represents the number of patients who had tooth mobility. Majority of the male patients (blue) had tooth mobility in the anterior and posterior region than female patients (green)



**Fig. 6.** Bar graph representing the association of age group and location of mobile tooth. X-axis represents the location of the mobile tooth and Y-axis represents the number of patients who had tooth mobility, The patients in the age group of 51-60 years presents with a higher number of mobile teeth in both anterior and posterior regions



**Fig. 7.** Bar graph representing the association of gender and region of tooth mobility. X axis represents the region of the mobile tooth (Jaws involved) and Y axis shows the number of patients who had tooth mobility. Tooth mobility is more commonly seen in mandibular arch than the maxillary arch and more common among males (blue) than females (green)



**Fig. 8.** Bar graph representing the association of age group and region of tooth mobility. X axis represents the region of the mobile tooth and Y axis represents the number of patients. Patients in the age group of 51-60 years present with higher number of tooth mobility and more commonly in the mandibular region than maxillary region

#### 4. CONCLUSION

Therefore, from this study, the pattern of tooth mobility in diabetic patients with periodontal disease was analysed. Patients' age, gender, site and region of mobile tooth are not associated with tooth mobility. Tooth mobility due to periodontal disease can occur equally in both male and females. The present study showed that 80% of the diabetic patients with periodontal disease had tooth mobility. Therefore, early diagnosis, preventive therapeutic measures are needed to prevent progression of periodontal disease which leads to tooth mobility and tooth loss.

#### CONSENT

It is not applicable.

#### ETHICAL APPROVAL

As per international standard or university standard ethical approval has been collected and preserved by the authors.

#### ACKNOWLEDGEMENT

This research was supported by Saveetha Dental College and Hospitals. We thank the department

of Periodontics, Saveetha Dental College for providing insight and expertise that greatly assisted the research. The present study was supported by the following Saveetha Institute of Medical and Technical sciences (SIMATS), Saveetha Dental College and Hospitals, Saveetha University and Kumaran TV center and furnitures, Attur, Salem.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Mühlemann HR. Tooth mobility: The Measuring method. initial and secondary tooth mobility. Vol. 25, Journal of Periodontology. 1954;22-9. Available:<http://dx.doi.org/10.1902/jop.1954.25.1.22>
2. Dombret J, Marcos E. Tooth mobility and containment. Rev Belge Med Dent. 1989;44(2):98-109.
3. Ravi S, Malaiappan S, Varghese S, Jayakumar ND, Prakasam G. Additive effect of plasma rich in growth factors with guided tissue regeneration in treatment of intrabony defects in patients with chronic

- periodontitis: A split-mouth randomized controlled clinical trial. Vol. 88, Journal of Periodontology. 2017;839–45.  
Available:<http://dx.doi.org/10.1902/jop.2017.160824>
4. Kokane VB, Patil SN. Management of grade III mobile anterior tooth in function using endostabilizer - A case report. *J Clin Diagn Res.* 2014;8(12):ZD06–7.
  5. Thamaraiselvan M, Elavarasu S, Thangakumaran S, Gadagi JS, Arthie T. Comparative clinical evaluation of coronally advanced flap with or without platelet rich fibrin membrane in the treatment of isolated gingival recession. *J Indian Soc Periodontol.* 2015;19(1):66–71.
  6. Serio FG. Patient and tooth specific risk indicators for posterior tooth fracture are identified. *J Evid Based Dent Pract.* 2005;5(3):164–5.
  7. Bernal G, Carvajal JC, Muñoz-Viveros CA. A review of the clinical management of mobile teeth. *J Contemp Dent Pract.* 2002;3(4):10–22.
  8. Kavarthapu A, Thamaraiselvan M. Assessing the variation in course and position of inferior alveolar nerve among south Indian population: A cone beam computed tomographic study. *Indian J Dent Res.* 2018;29(4):405–9.
  9. Lang NP, Lindhe J. *Clinical Periodontology and Implant Dentistry*, 2 Volume Set. John Wiley & Sons. 2015;1480.
  10. Ramesh A, Varghese S, Doraiswamy J, Malaiappan S. Herbs as an antioxidant arsenal for periodontal diseases. Vol. 5, *Journal of Intercultural Ethnopharmacology.* 2016;92.  
Available:<http://dx.doi.org/10.5455/jice.20160122065556>
  11. Davies S, Gray R, Linden G, James J. Occlusal considerations in periodontics. Vol. 191, *British Dental Journal.* 2001;597–604.  
Available:<http://dx.doi.org/10.1038/sj.bdj.4801245a>
  12. Varghese SS, Thomas H, Jayakumar ND, Sankari M, Lakshmanan R. Estimation of salivary tumor necrosis factor-alpha in chronic and aggressive periodontitis patients. *Contemp Clin Dent.* 2015;6(Suppl 1):S152–6.
  13. Sippel V, Pierlot GM, Renault B, Groenen PMA, Strasser DS. Activation of IL5R and CRTH2 on human eosinophils elicit a similar molecular response and reveal a synergistic effect. Vol. 5, *European Journal of Molecular and Clinical Medicine.* 2018;1–11.  
Available:<http://dx.doi.org/10.5334/ejmcm.1>
  14. Giannakoura A, Pepelassi E, Kotsovilis S, Nikolopoulos G, Vrotsos I. Tooth mobility parameters in chronic periodontitis patients prior to periodontal therapy: A cross-sectional study. Vol. 5, *Dental, Oral and Craniofacial Research;* 2019.  
Available:  
<http://dx.doi.org/10.15761/docr.1000284>
  15. Gajendran P, Parthasarathy H, Tadepalli A. Comparative evaluation of cathepsin K levels in gingival crevicular fluid among smoking and nonsmoking patients with chronic periodontitis. Vol. 29, *Indian Journal of Dental Research.* 2018;588.  
Available:[http://dx.doi.org/10.4103/ijdr.ijdr\\_95\\_17](http://dx.doi.org/10.4103/ijdr.ijdr_95_17)
  16. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol.* 2018;89(10):1241–8.
  17. Paramasivam A, Priyadharsini JV, Raghunandhakumar S, Elumalai P. A novel COVID-19 and its effects on cardiovascular disease. *Hypertens Res.* 2020;43(7):729–30.
  18. SG, TG, KV, Faleh AA, Sukumaran A, PNS. Development of 3D scaffolds using nanochitosan/silk-fibroin/hyaluronic acid biomaterials for tissue engineering applications. *Int J Biol Macromol.* 2018;120(Pt A):876–85.
  19. Del Fabbro M, Karanxha L, Panda S, Bucchi C, Nadathur Doraiswamy J, Sankari M, et al. Autologous platelet concentrates for treating periodontal infrabony defects. *Cochrane Database Syst Rev.* 2018;11:CD011423.
  20. Paramasivam A, Vijayashree Priyadharsini J. Mitomi Rs: New emerging microRNAs in mitochondrial dysfunction and cardiovascular disease. *Hypertens Res.* 2020;43(8):851–3.
  21. Jayaseelan VP, Arumugam P. Dissecting the theranostic potential of exosomes in autoimmune disorders. *Cell Mol Immunol.* 2019;16(12):935–6.
  22. Vellappally S, Al Kheraif AA, Divakar DD, Basavarajappa S, Anil S, Fouad H. Tooth implant prosthesis using ultra low power and low cost crystalline carbon bio-tooth sensor with hybridized data acquisition



- algorithm. *Comput Commun.* 2019;148:176–84.
23. Vellappally S, Al Kheraif AA, Anil S, Assery MK, Kumar KA, Divakar DD. Analyzing relationship between patient and doctor in public dental health using particle memetic multivariable logistic regression analysis approach (MLRA2). *J Med Syst.* 2018;42(10):183.
  24. Varghese SS, Ramesh A, Veeraiyan DN. Blended module-based teaching in biostatistics and research methodology: A retrospective study with postgraduate dental students. *J Dent Educ.* 2019;83(4):445–50.
  25. Venkatesan J, Singh SK, Anil S, Kim S-K, Shim MS. Preparation, characterization and biological applications of biosynthesized silver nanoparticles with chitosan-fucoidan coating. *Molecules.* 2018;23(6). Available:<http://dx.doi.org/10.3390/molecules23061429>
  26. Alsubait SA, Al Ajlan R, Mitwalli H, Aburaisi N, Mahmood A, Muthurangan M, et al. Cytotoxicity of different concentrations of three root canal sealers on human mesenchymal stem cells. *Biomolecules.* 2018;8(3). Available:<http://dx.doi.org/10.3390/biom8030068>
  27. Venkatesan J, Rekha PD, Anil S, Bhatnagar I, Sudha PN, Dechsakulwatana C, et al. Hydroxyapatite from cuttlefish bone: Isolation, characterizations, and applications. *Biotechnol Bioprocess Eng.* 2018;23(4):383–93.
  28. Vellappally S, Al Kheraif AA, Anil S, Wahba AA. IoT medical tooth mounted sensor for monitoring teeth and food level using bacterial optimization along with adaptive deep learning neural network. *Measurement.* 2019;135:672–7.
  29. PradeepKumar AR, Shemesh H, Nivedhitha MS, Hashir MMJ, Arockiam S, Uma Maheswari TN, et al. Diagnosis of vertical root fractures by cone-beam computed tomography in root-filled teeth with confirmation by direct visualization: A systematic review and meta-analysis. *J Endod.* 2021;47(8):1198–214.
  30. R H, Ramani P, Tilakaratne WM, Sukumaran G, Ramasubramanian A, Krishnan RP. Critical appraisal of different triggering pathways for the pathobiology of pemphigus vulgaris-A review. *Oral Dis;* 2021. Available:<http://dx.doi.org/10.1111/odi.13937>
  31. Ezhilarasan D, Lakshmi T, Subha M, Deepak Nallasamy V, Raghunandhakumar S. The ambiguous role of sirtuins in head and neck squamous cell carcinoma. *Oral Dis;* 2021. Available:<http://dx.doi.org/10.1111/odi.13798>
  32. Sarode SC, Gondivkar S, Sarode GS, Gadbail A, Yuwanati M. Hybrid oral potentially malignant disorder: A neglected fact in oral submucous fibrosis. *Oral Oncol.* 2021 Jun 16;105390.
  33. Kavarthapu A, Gurumoorthy K. Linking chronic periodontitis and oral cancer: A review. *Oral Oncol.* 2021;105375.
  34. Vellappally S, Abdullah Al-Kheraif A, Anil S, Basavarajappa S, Hassanein AS. Maintaining patient oral health by using a xeno-genetic spiking neural network. *J Ambient Intell Humaniz Comput;* 2018. Available:<https://doi.org/10.1007/s12652-018-1166-8>
  35. Aldhuwayhi S, Mallineni SK, Sakhamuri S, Thakare AA, Mallineni S, Sajja R, et al. Covid-19 knowledge and perceptions among dental specialists: A cross-sectional online questionnaire survey. *Risk Manag Healthc Policy.* 2021;14:2851–61.
  36. Zhou X. Is laser treatment effective for patients with periodontal disease?. <http://isrctn.com/>. Available:<http://dx.doi.org/10.1186/isrctn18400416>
  37. High prevalence of gingivitis and periodontitis and tooth loss in saudi adult male population and the possible association with the inheritance of abo blood groups. Vol. 8, *International Journal of Biology, Pharmacy and Allied Sciences;* 2019. Available: <http://dx.doi.org/10.31032/ijbpas/2019/8.7.4748>
  38. Periodontitis and chronic obstructive pulmonary disease. Vol. 57, *Dental Abstracts.* 2012;327. Available:<http://dx.doi.org/10.1016/j.denabs.2012.04.097>
  39. Priyanka S, Kaarthikeyan G, Nadathur JD, Mohanraj A, Kavarthapu A. Detection of cytomegalovirus, Epstein-Barr virus, and Torque Teno virus in subgingival and atheromatous plaques of cardiac patients with chronic periodontitis. *J Indian Soc Periodontol.* 2017;21(6):456–60.

40. Ramesh A, Vellayappan R, Ravi S, Gurumoorthy K. Esthetic lip repositioning: A cosmetic approach for correction of gummy smile - A case series. J Indian Soc Periodontol. 2019;23(3):290–4. Available:[http://dx.doi.org/10.4103/ijohr.ijohr\\_42\\_17](http://dx.doi.org/10.4103/ijohr.ijohr_42_17)
41. Benoist HM, Seck-Diallo A, Diouf A, Yabbre S, Sembene M, Diallo PD. Profile of chronic and aggressive periodontitis among Senegalese. J Periodontal Implant Sci. 2011;41(6):279–84.
42. Azodo C, Ogbemor O. Tooth mobility in a nigerian specialist periodontology clinic. Indian Journal of Oral Health and Research. 2017;3:62.
43. Mühlemann HR. Periodontometry, a method for measuring tooth mobility. Vol. 4, Oral Surgery, Oral Medicine, Oral Pathology. 1951;1220–33. Available:[http://dx.doi.org/10.1016/0030-4220\(51\)90080-1](http://dx.doi.org/10.1016/0030-4220(51)90080-1)
44. Rosales C, Uribe-Querol E. Neutrophil role in periodontal disease. Role of Neutrophils in Disease Pathogenesis; 2017. Available:<http://dx.doi.org/10.5772/67789>

---

© 2021 Muthukumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/80279>