



# Evaluation of Different Ornamental Ficus under Shadenet Conditions of Prayagraj

**Bhagyasri Nandam<sup>a+++\*</sup>, Devi Singh<sup>a#</sup> and C. John Wesley<sup>b</sup>**

<sup>a</sup> Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh, India.

<sup>b</sup> Centre for Geospatial Technologies VIAET, SHUATS, Prayagraj, Uttar Pradesh, India.

## **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## **Article Information**

DOI: 10.9734/IJECC/2023/v13i102614

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/104704>

**Original Research Article**

**Received: 04/06/2023**

**Accepted: 08/08/2023**

**Published: 12/08/2023**

## **ABSTRACT**

An experiment entitled Evaluation of different ornamental ficus under shadenet conditions of prayagraj was conducted in shadenet house, Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj during July-October 2022 on ficus. The experiment was conducted in Completely Randomized Design (CRD) with eight varieties replicated thrice. The varieties used were *Ficus microcarpa* golden, *Ficus microcarpa* moclame, *Ficus microcarpa* crassifolia, *Ficus triangularis* variegata, *Ficus benjamina* exotica, *Ficus benjamina* reginald, *Ficus benjamina* starlight, *Ficus benjamina* safari. The results revealed that better performance was seen in *Ficus microcarpa* golden in all parameters like plant height at 90 days (56.33 cm), no. of leaves at 90 days (94.43 cm), leaf length at 90 days (10.13 cm), leaf breadth at 90 days (4.53 cm), no. of primary branches at 90 days (11.23 cm), plant spread at 90 days (36.93 cm), leaf petiole length (1.6 cm).

<sup>++</sup>M.Sc. Scholar;

<sup>#</sup>Associate Professor;

<sup>\*</sup>Corresponding author: E-mail: bhagyasrinandam1702@gmail.com;

**Keywords:** *Ficus*; *Ficus microcarpa golden*; CRD; ornamental.

## 1. INTRODUCTION

*Ficus* a genus of about 850 species woody trees, shrubs, vines, epiphytes and hemiepiphytes in the Family Moraceae Collectively known as fig trees or figs, they are native throughout the tropics with a few species extending into the semi-warm temperate zone.

The maximum diversity of the genus exhibits in Asiatic mainland (170 spp.), New Guinea (132 spp.) and Borneo (129 spp.). Many species of *Ficus* L. are very common in different biogeographic regions. Although, the great majority of the species grow in lowlands but some of them reach up to about 2,000 m altitudes [1-3]. *Ficus* is also considered one of the most diversified genera with regard to its habits (deciduous and evergreen trees, shrubs, herbs, climbers and creepers) and life forms (free standing tree, epiphytes, semi-epiphytes in the crevices, rheophytes and lithophytes) [4-7].

Ornamental plants or garden plants are plants that are primarily grown for their beauty but also for qualities such as scent or how they shape physical space. Many flowering plants and garden varieties tend to be specially bred cultivars that improve on the original species in qualities such as color, shape, scent, and long-lasting blooms. There are many examples of fine ornamental plants that can provide height, privacy, and beauty for any garden. These ornamental perennial plants have seeds that allow them to reproduce [8]. One of the beauties of ornamental grasses is that they are very versatile and low maintenance. Almost any types of plant have ornamental variety/varieties: trees, shrubs, climbers, grasses, succulents, aquatic plants, herbaceous perennials and annual plants. Non-botanical classifications include houseplants, bedding plants, hedges, plants for cut flowers and foliage plants. The cultivation of ornamental plants comes under floriculture and tree nurseries, which is a major branch of horticulture [9,10].

*Ficus benjamina* is a tree reaching 30 m (98 feet) tall in natural conditions, with gracefully drooping branchlets and glossy leaves 6–13 cm, oval with an acuminate tip. The bark is light gray and smooth [11-13]. The bark of young branches is brownish. The widely spread, highly branching tree top often covers a diameter of 10 meters. It is a relatively small-leaved fig. The changeable leaves are simple, entire and stalked. The petiole

is 1 to 2.5 cm (3/8 to 1 inch) long. The young foliage is light green and slightly wavy, the older leaves are green and smooth; the leaf blade is ovate to ovate-lanceolate with wedge-shaped to broadly rounded base and ends with a short dropper tip. The pale glossy to dull leaf blade is 5 to 12 cm long and 2 to 6 cm wide. Near the leaf margins are yellow crystal cells ("cystolites"). The two membranous, deciduous stipules are not fused, lanceolate and 6 to 12 mm (1/4 to 1/2 inch) (rarely to 15 mm or 9/16 inch) long.

*F. benjamina* is monoecious. The inflorescences are spherical to egg-shaped, shiny green, and have a diameter of 1.5 cm (1/2 inch).

*Ficus microcarpa* is a tropical tree with smooth light-gray bark and entire oblanceolate leaves about 2–2.5 inches (5.1–6.4 centimetres) long which in Mediterranean climates grows to about forty feet (twelve meters) tall and with an equal spread of banyan habit (tropical and humid subtropical) numbers of prop roots.

## 2. MATERIALS AND METHODS

The investigation was carried out at the shadenet, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj during the year July-October, 2022. The experiment was conducted in completely randomized design with 8 varieties in three replications viz. V<sub>1</sub> *Ficus microcarpa golden*, V<sub>2</sub> *Ficus microcarpa moclame*, V<sub>3</sub> *Ficus microcarpa crassifolia*, V<sub>4</sub> *Ficus triangularis variegata*, V<sub>5</sub> *Ficus benjamina exotica*, V<sub>6</sub> *Ficus benjamina reginald*, V<sub>7</sub> *Ficus benjamina starlight*, V<sub>8</sub> *Ficus benjamina safari* was planted in eight inches pots holes on both sides was done at 30, 60, 90 Days after emergence. The data was recorded for the following parameters viz Plant height (cm), Number of leaves, Leaf length (cm), Leaf breadth (cm), No. of primary branches per plant, Plant spread (cm), Leaf petiole length (cm).

## 3. RESULTS AND DISCUSSION

### 3.1 Vegetative Characters

#### 3.1.1 Plant height

Plant height varies significantly between different Varieties. Longer plant height at 90days after planting was found in variety V<sub>1</sub> *Ficus microcarpa golden* (56.33 cm) followed by 54.33 cm in V<sub>5</sub>

*Ficus benjamina* exotica whereas shorter plant height 33.7cm was recorded in V<sub>8</sub> *Ficus benjamina* safari in Table 1.

In plants higher auxin transport just below the apical meristem zone might have increased the plant height by cell elongation. The plants grown under low light level were found to be more apical dominant than those grown under high or full light intensities.

The results are in conformity with findings of Thangam and Thamburaj, (2008) who stated that partial shading improves the plant height than full sunlight, which compared the plant growth under shadenet condition.

### 3.1.2 Number of leaves

Number of leaves varies significantly between different Varieties. Highest number of leaves at 90days after planting was found in variety V<sub>1</sub> *Ficus microcarpa* golden (94.43 cm) followed by 90.23 cm in V<sub>5</sub> *Ficus benjamina* exotica whereas lowest number of leaves 64.86 cm was recorded in V<sub>8</sub> *Ficus benjamina* safari in Table 1. As light capturing ability and chlorophyll content is high in shade condition, at this condition N leads to increase numbering of leaves.

### 3.1.3 Leaf length

Leaf length varies significantly between different Varieties. Longer leaf length at 90days after planting was found in variety V<sub>1</sub> *Ficus microcarpa* golden (10.13 cm) followed by 9.56 cm in V<sub>5</sub> *Ficus benjamina* exotica whereas shorter leaf length 4.93 cm was recorded in V<sub>8</sub> *Ficus benjamina* safari in Table 1.

The number of leaves is an important parameter to consider because it donate health status of plant [14]. The various physiological functions like photosynthesis,transpiration and capability to tolerate air pollution etc depend on number of leaves in a plant.

### 3.1.4 Leaf breadth

Leaf breadth varies significantly between different Varieties. Longer leaf breadth at 90days after planting was found in variety V<sub>1</sub> *Ficus microcarpa* golden (4.53 cm) followed by 4.16 cm in V<sub>5</sub> *Ficus benjamina* exotica whereas shorter leaf breadth 2.76 cm was recorded in V<sub>8</sub> *Ficus benjamina* safari in Table 1 [15-20].

The boundary layers help to improve leaf growth and photosynthetic activity under low red: far red ratio which elongates leaf breadth.

The results were in agreement with Das (2010) who stated the leaf breadth was influenced by low light irradiance.

### 3.1.5 Number of primary branches

Number of primary branches varies significantly between different Varieties. Highest number of primary branches at 90days after planting was found in variety V<sub>1</sub> *Ficus microcarpa* golden (11.23 cm) followed by 10.26 cm in V<sub>5</sub> *Ficus benjamina* exotica whereas shorter number of primary branches 5.33 cm was recorded in V<sub>8</sub> *Ficus benjamina* safari in Table 1.

Shade netting that increases light scattering and has been shown to increase branching and vegetative growth.

Similar results were obtained by Brand (1997) studied the effect of shade during nursery production on the growth, no. of branches and foliage colour. Results showed that shade improved no. of branches effectively and the best ornamental quality was also obtained [21,22].

### 3.1.6 Plant spread

Plant spread varies significantly between different Varieties. Maximum plant spread at 90days after planting was found in variety V<sub>1</sub> *Ficus microcarpa* golden (36.93 cm) followed by 33.0 cm in V<sub>5</sub> *Ficus benjamina* exotica whereas minimum plant spread 19.5cm was recorded in V<sub>8</sub> *Ficus benjamina* safari in Table 1.

Plant spread is an important character considering the ficus for interior plantscaping. The minimum the spread, more compact will be the plants and this makes them suitable for indoors. The plants with more branches/laterals were found to have more spread and plants with vertical growth rather than lateral were having lesser spread. Such differences in plant spread were also observed by Russ and Pertuit (2001) in different foliage plant species [23-35].

### 3.1.7 Leaf petiole length

Leaf petiole length varies significantly between different Varieties. Longer leaf petiole length at 90days after planting was found in variety *Ficus microcarpa* golden (1.6 cm) followed by 1.3 cm in V<sub>5</sub> *Ficus benjamina* exotica whereas shorter leaf petiole length 0.36 cm was recorded in V<sub>8</sub> *Ficus benjamina* safari in Table 1.

**Table 1. Plant height, no.of leaves, leaf length, leaf breadth, no. of primary branches, plant spread, leaf petiole length, of different ficus ornamental plants for 90 days**

Varieties	Plant height	No.of leaves	Leaf length	Leaf breadth	No. of primary branches	Plant Spread	Leaf petiole length
<i>Ficus microcarpa</i> golden	56.33	94.43	10.13	4.53	11.23	36.93	1.6
<i>Ficus microcarpa</i> moclame	54.06	87.4	8.93	3.8	9.66	31.06	1.23
<i>Ficus microcarpa</i> crassifolia	50.96	78.53	7.7	3.5	7.56	26.4	1.11
<i>Ficus triangularis</i> variegata	46.56	71.86	6.6	3.3	6.6	25.63	1.0
<i>Ficus benjamina</i> exotica	54.33	90.23	9.56	4.16	10.26	33	1.3
<i>Ficus benjamina</i> reginald	53.06	84.5	8.2	3.53	8.56	28.6	1.18
<i>Ficus benjamina</i> starlight	35.1	67.4	5.6	3.1	6.13	25	0.81
<i>Ficus benjamina</i> safari	33.7	64.86	4.93	2.76	5.33	19.5	0.36
<b>Mean</b>	48.01	79.90	7.70	3.5875	8.17	28.26	1.06
<b>F-Test</b>	S	S	S	S	S	S	S
<b>SE(M)</b>	0.60	1.04	0.13	0.07	0.20	0.90	0.07
<b>CD at 5%</b>	1.82	3.12	0.40	0.21	0.61	2.72	0.21
<b>CV</b>	2.19	2.25	3.05	3.46	4.35	5.57	11.57

**Table 2. Leaf texture, leaf margin, leaf type, leaf arrangement, leaf venation, leaf shape, pigmentation of different ficus ornamental plants for 90 days**

Varieties	Leaf texture	Leaf margin	Leaf type	Leaf arrangement	Venation	shape	Pigmentation
<i>Ficus microcarpa</i> golden	Leathery	Entire	Simple/ Unifoliate	Alternate	Pinnate	Oval-Elliptic	Younger leaves tend to be more yellowish or golden. Mature leaves tend to be yellow/ golden green.
<i>Ficus microcarpa</i> moclame	Leathery	Entire	Simple	Alternate	Pinnate	Ovoid lanceolate	Green
<i>Ficus microcarpa</i> crassifolia	Leathery, thick, waxy.	Entire	Simple	Alternate	Pinnate	Oblong-Elliptic-Ovate	Young/immature: Bronze to yellow-green. Mature: Bright medium green to dark green.
<i>Ficus triangularis</i> variegata	Smooth, glossy, waxy.	Entire	Simple	Alternate	Pinnate	Obovate (Triangle shaped)	The triangular leaf has irregular yellowish-white edges, with shades of green sections in the centre.
<i>Ficus benjamina</i> exotica	Both sides waxy, smooth and leathery.	Entire	Simple	Alternate	Pinnate	Oblong-Elliptic	Young/immature: Light green. Mature: Bright medium green to dark green.
<i>Ficus benjamina</i> reginald	Leathery	Entire	Simple	Alternate	Pinnate	Elliptic	Yellow-green leaves that have irregular dark green markings around the midrib.
<i>Ficus benjamina</i> starlight	Smooth, glossy/ shiny	Entire	Simple/ Unifoliate	Alternate	Pinnate	Ovate-Elliptic	Leaves are glossy green with irregularly white margins.
<i>Ficus benjamina</i> safari	Leathery	Entire	Simple	Entire	Pinnate	Vine	The leaf has irregular yellowish-white edges, with shades of green sections in the center.

### 3.2 Leaf Characters

Leaf texture, leaf margin, leaf arrangement, leaf shape, leaf type, leaf venation, pigmentation are taken as qualitative characters as it helped to identify the morphological characters of leaf. The data estimated on qualitative leaf characters were recorded and significant differences were observed at 90 days after planting shown in Table 2.

### 4. CONCLUSION

Based on the results of the present investigation entitled "Evaluation of different ornamental ficus under shadenet conditions of prayagraj" it was concluded that the variety V<sub>1</sub> *Ficus microcarpa* golden reported significantly better performance in all of the parameters like plant height(cm), No. Of leaves, leaf length (cm), leaf breadth (cm), no. Of primary branches, plant spread (cm), leaf petiole length (cm) .Hence the variety *Ficus microcarpa* golden could be recommended for the prayagraj agro-climatic condntions.

### ACKNOWLEDGEMENT

The author is thankful to the Advisor and Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and sciences, Prayagraj (U.P.) India for providing necessary facilities to undertaken this research. This manuscript has not received any grant from any sources.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Barrett JE, Nell TA. *Ficus benjamina* response to growth retardants. Proc. Fla. State Hort. Soc. 1983;96:264-265.
2. Berg CC. Flora Malesiana Precursor for the Treatment of Moraceae 4: *Ficus* subgenus *Synoecia*. Blumea. 2003;48 (3).
3. Chang, Lester. Auntie Sarah's Banyan. The Garden Island. Lihue; 2003.
4. Denisowski Paul. Chinese–English Dictionary; 2007. Retrieved November 1, 2008.
5. Fails BS, Lewis AJ, Barden JA. Anatomy and morphology of sun- and shade-grown *Ficus benjamina*. J. Amer. Hort. Sci. 1982;107(5):754-757.
6. Johnson CR, Ingram DL, Barrett JE. Effects of irrigation frequency on growth, transpiration, and acclimatization of *Ficus benjamina* L. Hort Science. 1981;16(1):80-81.
7. Kislev Mordechai E. Hartmann Anat, Bar-Yosef Ofer. Early Domesticated Fig in the Jordan Valley. Science. 2006;312(5778): 1372–1374. DOI:10.1126/science.1125910 PMID 16741119. S2CID 42150441.
8. Lewington, Anna, Parker, Edward. Ancient trees: Trees that live for 1000 years. London: Collins & Brown; 1999. ISBN 978-18-5585-704-9.
9. Conover CA, Poole RT. *Ficus benjamina* leaf drop. Florists' Rev. 1973;151(3925): 29, 67, 68.
10. Corner E.J.H. Check-list of *Ficus* in Asia and Australasia with keys to identification. The Gardens' Bulletin Singapore. (digitised, online, via biodiversitylibrary.org). 1965;21(1):1–186. Retrieved 5 Feb 2014. pages 22–23.
11. Chen, Ying-Ru, Wen-Chung Chuang, Wen-Jer Wu. Chalcids wasps on *Ficus microcarpa* L. in Taiwan (Hymenoptera: Chalcidoidea). Journal of Taiwan Museum. 1999;52:39–79.
12. Condit Ira J. *Ficus: The exotic species*. University of California, Division of Agricultural Sciences. OCLC. 1969; 1086743649.
13. Conklin E. A guide to interior plantings. Everett Conklin and Co., Montvale, N. J. 1970;25.
14. Lev-Yadun Simcha, Ne'eman Gidi, Abbo Shahal, Flaishman Moshe A. Comment on Early Domesticated Fig in the Jordan Valley. Science. 2006;314(5806):1683a. Bibcode: 2006 Sci. 314.1683L. DOI:10.1126/science.1132636 PMID 17170278.
15. Littlecott, Lorna. Hawai'i First. American Forests. 1969;75(2):61.
16. Peterson, J.C. 1986. The impact of leaf anatomy on *Ficus* survival indoors. Interior Landscape Industry 3(4):30-35.
17. Peterson NC, Blessington TM. Postharvest effects of dark storage and light source on keeping quality of *Ficus benjamina* L. Hort Science. 1981;16(10):681-682. 34. and. Antitranspirant and dark storage effects on the postharvest quality of *Ficus benjamina* L. Florists Review. 1982; 170(4402):12, 13, 41.

18. Riefner, Richard E. Jr. *Ficus microcarpa* (Moraceae) naturalized in Southern California, U. S. A.: Linking plant, pollinator, and suitable microhabitats to document the invasion process *Phytologia*. 2016;98(1):42-75. ISSN 0031-9430.
19. Rønsted N, Weiblen GD, Clement WL, Zerega NJC, Savolainen V. Reconstructing the phylogeny of figs (*Ficus*, *Moraceae*) to reveal the history of the fig pollination mutualism (PDF). *Symbiosis*. 2008;45. ISSN 0334-5114.
20. Rønsted Nina, Weiblen George D, Cook, James M, Salamin, Nicholas, Machado, Carlos A, Savoianen Vincent. 60 million years of co-divergence in the fig-wasp symbiosis. *Proceedings of the Royal Society B: Biological Sciences*. 2005; 272(1581):25932599. DOI: 10.1098/rspb.2005.3249 PMC 1559977. PMID 16321781.
21. Shanahan M, SG, So, Samson, Corlett Richard. Compton, Fig-eating by vertebrate frugivores: A global review. *Biological Reviews*. 2001;76(4):529–572. DOI: 10.1017/S1464793101005760 PMID 11762492. S2CID 27827864.
22. Somenzari, Marina, Linda Lacerda da Silva, Rosanna GQ Benesi. Atração de aves por *Ficus elastica* Roxb. e *Ficus microcarpa* L. em ambiente urbano (abstract) (PDF). XIV Congresso Brasileiro de Ornitologia; 2006. Archived from the original (PDF) on 2009-02-06.
23. Sommer, Anthony. State's Biggest Banyan Bears Heavy Fire Damage. *Star Bulletin*. Honolulu; 2000.
24. Sommer Anthony. Ancient Banyan Tree Returns Protector's Favor. *Star Bulletin*. Honolulu; 1999.
25. Stenitz B, Ben-Jaacov J, Ackerman A, Hagiladi A. Dark storage of three cultivars of bare-root *Ficus benjamina* foliage plants. *Scientia Horticulturae*. 1987;32: 315-322.

© 2023 Nandam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/104704>