



Impacts of the Use of *Chrysophyllum albidum* G. Don on Its Vulnerability and Its Conservation in Benin

Blaise Alowanou-Kélé ^{a,b*}, Antoine Abel Missihoun ^{a,b},
Relique Ignace Agbo ^{a,b}, Eben-Ezer B. K. Ewédjé ^c,
Hounnankpon Yédomonhan ^d and Clément Agbangla ^{a,b}

^a Faculty of Sciences and Techniques, Laboratory of Molecular Genetics and Genome Analysis, University of Abomey-Calavi, 01 PO Box 526, Cotonou, Benin.

^b Laboratory of Genetics and Biotechnologies, University of Abomey-Calavi, 01 PO Box 526, Cotonou, Benin.

^c Laboratory of Botany, Applied Plant Ecology and Forest Genetics, National Higher School of Biosciences and Applied Biotechnologies of Dassa (ENSBBA-Dassa), National University of Sciences, Technologies, Engineering and Mathematics of Abomey, UNSTIM-Abomey, Bénin.

^d Laboratory of Botany and Plant Ecology, Faculty of Science and Technology, University of Abomey-Calavi, 01 PO Box 4521 Cotonou, Benin.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARRB/2023/v38i1130613

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111614>

Original Research Article

Received: 23/10/2023

Accepted: 29/12/2023

Published: 30/12/2023

ABSTRACT

Aims / Objectives: *Chrysophyllum albidum* is a fruit tree of great socio-economic importance in West Africa. In several localities in Benin, despite the species over-exploitation of the species, few data relating to its viability and the conservation of its genetic diversity exist. The objective of this

*Corresponding author: E-mail: alowanoukelblaise@gmail.com;

study is to determine the values and forms of uses, of various organs of the species by the populations, in order to identify their impacts on the vulnerability of the species and its sustainable use.

Methodology: A semi-structured survey has collected from 877 an (eight hundred and seventy-seven) person's taken at random, ethnobotanical data on the species. The frequencies of use of the different organs, the index vulnerability of the species, as well as the dendrometric parameters were evaluated. In addition, Cramer's V test was used to analyze the influence of sex and socio-cultural groups on the different uses made of the species.

Results: It appears that the species is more and more essential, because it is better known and has priority used in human and animal food, the cure of certain bodily and spiritual diseases but also serves as energy wood and lumber. The frequency of use organs varies from 2.28% (flower) to 99.54% (fruits), passing through 25.88% (root), 35.23% (bark), 49.82% (leaves) and 82.78% (wood). Furthermore, the high number of categories of use (NCU = 6), the frequency of use (FUO = 49.25%), the method of collection (picking and cutting), the types of organs used (wood, fruits, bark, leaves, flower and root), the presence or density of trees of the species in the study environment (less abundant), and the stage of development (adult), made it possible to have the value 2.5 as a vulnerability index. Thus, the degree of stress on the different organs of *C. albidum* makes the species very vulnerable (VI = 2.50) in Benin.

Conclusion: These different data reveal the need to design strategies for the sustainable management of the plant genetic resources of this species useful in Benin.

Keywords: *Chrysophyllum albidum*; anthropogenic activity; sustainable use; traditional uses; vulnerability; Benin.

1. INTRODUCTION

Apart from agriculture and breeding Non-Timber Forest Products (NTFP) constitute a major source of medicine, food and income for the rural populations [1]. NTFPs are resources plant genetics that effectively contribute to the food self-sufficiency of our local populations and communities [2].

Chrysophyllum albidum is one of the forest tree species that provide NTFP of great importance to the inhabitants of rural and urban areas of West Africa with great export potential [3]. Fruit tree growing as a wild plant, *C. albidum* belongs to the family Sapotaceae and to the order Ericales. Its fruit, widely used in food, is sold in local markets. Their fruit pulp contains 8.8% protein, 15.1% fat, 3.4% ash, 4% crude fiber and 68.7% carbohydrate [4]. The species is used in the traditional pharmacopoeia where its leaves, its roots and its bark intervene in the treatment of many diseases or symptoms such as malaria, sexual weakness, asthma, jaundice. Its wood is used both as firewood and lumber for local populations [5].

Ranked among the ten priority woody food species in Benin [6], *C. albidum* is therefore a species on which action should be taken aimed at better understanding the socio-economic importance, uses and pressures that threaten its

preservation and conservation, especially since it does not yet benefit until day, of a real research program for its sustainable development. Among others factors causing the disappearance of *C. albidum* in Benin, we note aging, urbanization and the sacred nature of the species [7], the systematic collection of fruits of the species for marketing [8]. However, in Benin, the favorable habitats of *C. albidum*, are not sufficiently threatened by the climate changes [9]. So the probability of survival of the species in the face of climate change is high. Furthermore, previous ethnobotanical studies carried out on the species have shown that it provides multiple goods and services to local populations, because it presents several categories of use [5, 8] which vary according to time and socio-cultural groups holding knowledge diversified endogenous species transmitted from generation to generation [10]. Certainly, this dynamic in the forms of use makes the species more essential to the rural populations but also increases their vulnerability [7]. Consequently, the anthropogenic actions remain one of the greatest factors causing the disappearance of the species in Benin. However, such losses have a negative and long-term impact on the endogenous knowledge of several plant species [11]. Clearly, the taking into account the endogenous perceptions of local populations are essential to the design of adequate strategies that can allow the sustainable exploitation of resources plant

genetics [12]. Use values determine the interest of the rural populations towards plant species [13], while the frequency of use of the organs and the number of categories of use of the species, and provide information on the vulnerability index [14].

The demographic and economic growth experienced by the localities of Benin, increases the anthropogenic actions exerted on the populations of the species, due to the ever-increasing demand for his organs. Hence the relevance of this study which has general objective of assessing the level of vulnerability of the species in a context where its different organs are used daily. Specifically, this study aims to: (i): identify the different forms of use of the organs of *Chrysophyllum albidum* trees and (ii) determine the frequencies of use of plant organs and the value of its index of vulnerability.

2. MATERIALS AND METHODS

2.1 Study Zone

This study took place in Benin in different localities located in the six phytogeographical districts where *C. albidum* individuals are present (Fig 1). It's about localities in the Coastal, Pobè, Ouémé Valley and Plateau districts of the Guineo- Congolese located between 6°25'N and 7°30'N; and the Zou and Bassila districts of the zone of Sudano-Guinean transition located between 7°30'N and 9°45'N. These areas vary depending on the nature of the ground, the climate and the type of activity carried out by the inhabitants. Groups plants that host the species are those of concessions, the edge of lowlands, forests relics of fallows then fields and plantations [15].




Fig. 1. Prospected localities

2.2 Sampling and Ethnobotanical Surveys

A preliminary survey made it possible to interview 34 people chosen in a way randomly by phytodistrict, i.e. 204 people in all six phytodistricts constituting the study area. The data collected made it possible to identify the localities where the species is present and to calculate the size of the sample in each of the phytodistricts, from the formula of Dagnelie [16] presented below:

$$n = \frac{U_{1-\alpha/2}^2 \sum P_i (1-P_i)}{d^2} ; \text{ with:}$$

n, the size of the sample to be considered by phytodistrict in this study;

$U_{1-\alpha/2}$, the value of the normal random variable for a risk $\alpha = 0.05$ ($U_{1-\alpha/2} = 1.96$ for $\alpha = 0.05$);

P_i , the proportion of people who know and use *Chrysophyllum albidum*;

d, the allowable margin of error. It is 0.05.

2.3 Collection of Data

877 people, made up of 256 women and 621 men, were interviewed with success, based on a semi-structured questionnaire [17] and a clear picture of the species. These respondents were chosen at random from all the localities, and interviewed individually (Fig. 2). The established questionnaire concerned the categories of use of the species, the parts of the plant that are exploited, the method of harvesting the organs of the plant, the opinion of the respondent on the abundance of the species in the localities of the study area, the probable reasons for the scarcity of *C. albidum* trees and the strategies to adopt for the sustainable development of the species. Moreover, in the localities where the actual presence of feet of the species, it was measured the dendrometric data such as the total height of the plants of the species and the diameter at chest height (Fig. 3).

2.4 Data Analysis

The information collected made it possible to identify the different categories of uses and uses made of the various organs of *C. albidum*, and to calculate the following parameters explained:

- Frequency of use (FUO)

The frequency of use made it possible to know the level of use of each of the organs of the plant by the target populations. It is expressed by the formula used by Agbo et al [19] and is as follows:

$$FUO = \frac{N \times 100}{N_t} ; \text{ with:}$$

N, the number of people who cited this body;

N_t , the total number of respondents.

- Average Value of Use (AVU)

According to Gomez-Beloz [18], the average value of use makes it possible to evaluate the local knowledge of the populations on a species. It is the total number of reported uses by an interviewee, and is calculated in this study according to socio-cultural groups and sex, through the following formula:

$$AVU = \frac{N_{Ut}}{N} ; \text{ with:}$$

N_{Ut} , the total number of uses reported by all respondents in a category data;

N, the total number of respondents considered.

- Vulnerability index (VI)

The vulnerability index was calculated to assess the impact of the forms of use of the species on its viability. The formula used is as follows:

$$VI = \frac{\sum N_x}{N} ; \text{ with:}$$

N_x , the value given to x according to its degree of vulnerability;

N, the number of parameters serving as major indicators of pressure and threat on the species.

By parameter, the vulnerability scale considered varies from 1 to 3 [14]. For interpretation of our vulnerability scale, it was taken as reference values, the interpretation thresholds of Traoré et al [11]. For Traoré et al [11], a species is said to be weakly vulnerable when its vulnerability index is strictly less than 2; a species is said moderately vulnerable when its vulnerability index $VI \in [2; 2.5]$; a species is said very vulnerable when its vulnerability index is greater than or equal to 2.5.



Fig. 2. Ethnobotanic survey interview



Fig. 3. Diameter measurement

Table 1. Sample size by phytodistrict

Phytodistricts	Number of people knowing <i>C. albidum</i> among the 34 presurveyed per phyto-district	Proportion (Pi)	Size of the sample considered in the phyto-district (n)
Bassila	25	0,735	300
Pobè	31	0,911	125
Plateau	32	0,941	86
Coastal	32	0,941	86
Valley of Ouémé	32	0,941	86
Zou	29	0,852	194
Total	-	-	877

Table 2. Adapted parameters measuring the vulnerability of *Chrysophyllum albidum*; [11], [19] and [20]

Parameters (Pi)	Types of vulnerability		
	Weak (Scale = 1)	Average (Scale = 2)	Full (Scale = 3)
P ₁ : Frequency of use (FUO)	FUO < 20%	20 % ≤ FUO ≤ 60%	FUO ≥ 60%
P ₂ : Number of categories of use (NCU)	NCU < 2	2 ≤ NCU ≤ 4	NCU ≥ 5
P ₃ : Plant organ used	Latex ; leaf	Branch ; fruit	Wood ; seed ; flower ; bark ; root
P ₄ : Presence in the environ-ment	Abundant	Less abundant	Rare
P ₅ : Collection mode	Collection	-	Picking ; cutting
P ₆ : Stage of development	Old/ senescent	Adult	Young

Statistical analyzes were performed with R version 4.2.0 software. To do this, field data underwent Cramer's V test and Correspondence Analysis Multiples (CAM), to identify the possible influence of gender and socio-cultural groups investigated on the forms of use of the species.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 *Chrysophyllum albidum*, multi-purpose species

In Benin, local people use *Chrysophyllum albidum* for several purposes. Thus six (6) categories of use of the species have been identified. The ripe fruits of the species constitute for men, food (human food) very well appreciated; its wood is intensively used as primary fuel (wood energy) and timber; its bark, leaves and roots are used as fodder and very phytomedicine very effective. The flowers and leaves contribute to the fight against evil spirits (uses medico-magic). 99.54% of informants exploit the species as food while 75.25% and

46.18% use it respectively as energy wood and phytomedicine. By against, 35.68% of respondents say they rather exploit the species for its wood used as wood work. In addition, 18.47% and 2.73% of the interviewees declare using the species respectively as fodder and medico-magical species (Fig. 4).

Several uses are made of plant organs (Table 3). The fruits and wood are the most exploited organs with respectively 99.54% and 82.78% of frequencies. Follow after, the leaves, the bark and the root which have respectively for frequencies 49.82%, 35.23% and 25.88%. On the other hand, the flower is little used because; it has a more or less low frequency of use (2.28%) more or less weak. Better, the fruits of this species are more sought after by the Gouns and the Fons; its leaves are mainly exploited by Kotafons and Nagots; the wood of the species is more used by the Gouns and the Nagots; the roots and barks are used more by the Nagots and the Kotafons (Fig 5). The average use values of *Chrysophyllum albidum* organs (Fig 6) reveal that men (AVU = 6.75) have almost the same knowledge as women (AVU = 6.42), in the

use of the species. Within socio-cultural groups, average use values vary, from 3 to 8.42 and show that the Kotafons (AVU = 8.42), the Aïzos (AVU = 7.90), Nagots and Adjias (AVU = 7) respectively hold more than knowledge in the use of the species, that the Fons (AVU = 6.34), the Gouns (AVU = 5.55), the Mahi (AVU = 5.39) and the Ouatchis (AVU = 3).

The results of Cramer's V test (Table 4) reveal that the link between gender and categories of use of *C. albidum* is low (P-value = 0.024; value very close to 0), so that the link between socio-

cultural groups and these categories of use is strong (P-value = 0.83 → value close to 1). Multiple Correspondence Analysis (MCA) indicates that the link between sex and certain forms of use of the plant genetic resources of the species (human food, fodder, phytomedicine, timber and fuel wood) is low; on the other hand, medico-magical uses are significantly influenced by gender (Fig. 7). Furthermore, the socio-cultural groups identified influence the forms of use of the different organs of *C. albidum*.

Table 3. Categories of use and frequencies of use of *Chrysophyllum albidum* organs

Organs	Categories of use	Frequencies of use (%)	Uses
Fruit	Human food	99,54	Directly consume at ripeness
	Wood		
	Firewood		Wood bundles / Wood coal
	Lumber	82,78	Woodwork / Woodart and craft
	Animal feeding		Provender
Leaf	Phytomedicine	49,82	Treat diseases
	Medical and magical uses		Fight against evil spirits
	Bark		
	Phytomedicine	35,23	Treat diseases
Root	Phytomedicine	25,88	Treat diseases
Flower	Medical and magical uses	2,28	Fight against evil spirits

Table 4. Links between the categories of use of *Chrysophyllum albidum* and the socio-demographic parameters (gender and socio-cultural group)

Cramer's V test	
Socio-demographic parameters	P-value
Gender	0,024
Socio-cultural groups	0,83




Fig. 4. Frequencies of *Chrysophyllum albidum* use categories in Benin




Fig. 5. Use of *Chrysophyllum albidum* organs by socio-cultural groups investigated




Fig. 6. Variation in the Average Value of Use (AVU) according to gender and sociocultural groups

3.1.2 Vulnerability of *Chrysophyllum albidum* in Benin

Due to the high number of usage categories (NCU = 6), the frequency of use (FUO = 49.25%), mode of collection (picking and cutting), types of organ exploited (wood, fruit, bark, leaf, flower and root), the presence in the study environment of feet of the tree (less abundant), and of the stage of development (adult), *Chrysophyllum albidum* has a vulnerability index

of 2.50. Therefore the species is very vulnerable ($VI \geq 2.50$) in Benin (Table 5).

3.1.3 Conservation status of *Chrysophyllum albidum* in Benin

Nowadays, *Chrysophyllum albidum* is mainly present in certain houses. However, many people cut off the feet of the species that were in their homes, because of the lack of space that these trees often create, when the environment

life is favorable to them. Similarly, some people have testified to having cut the old trees of *C. albidum*, because for them, the old age of these plants decreases the quality and quantity of fruit produced. The plants of *C. albidum*, identified in Pobè phytodistrict, are more large (TrCir =

202.40 cm) and taller (Hpl = 23.86 m) for the most part (Fig. 8). Furthermore, people familiar with the plants of *C. albidum*, as a plant strongly medicinal, overexploit some of its organs such as roots, bark and leaves, to treat many bodily and/or spiritual illnesses (Fig 9).




Fig. 7. Correspondence Analysis Graph Multiples (CAM) on the categories of use of the species in sex function




Fig. 8. Diagram showing the height of the plants and the circumferences of the trunks of *Chrysophyllum albidum* by phyto-district of effective presence of the species



Fig. 9. Rised debarking of the trunk of *Chrysophyllum albidum*

Table 5. Scale of vulnerability by considered parameters

Considered parameters	Results obtained and comments	Affected vulnerability scales
Frequency of use	49,25%	N1 = 2
Number of categories	NCU = 6	N2 = 3
Exploited plant organs	Wood, fruit, bark, leaf, flower and root	N3 = 3
Presence in the medium	Less abundant	N4 = 2
Mode of collection	Picking and cutting	N5 = 3
Stage of development	Mainly adult	N6 = 2

3.2 Discussion

3.2.1 Links between the use of *Chrysophyllum albidum*, vulnerability and its availability

To live, the local populations are involved in agriculture whose production is complemented by non-timber forest products [21]. *Chrysophyllum albidum* with its multiple uses (NCU = 6), is a plant widely used by rural populations. However, the populations judge a plant important, when they can take its wood, its bark, roots, fruits and leaves for useful purposes

[11]. It is therefore because the different organs of *C. albidum* have not escaped this mesh. From where anthropic action contributes a lot to the destruction of the vegetation cover through the world. This pressure on the plant genetic resources of trees is even more increased with the increase in population size. As a result, plants become more vulnerable day by day. This is the case of *C. albidum*, whose index of vulnerability VI calculated at 2.50 ($VI \geq 2.50$) clearly reveals that the species is very vulnerable in Benin, according to the interpretation thresholds defined by [11]. The parameters that used to calculate this vulnerability are the high

number of use categories (NCU = 6), the frequency of use (FUO = 49.25%), the mode of collection (picking and cutting), the types of organ exploited (wood, fruit, bark, leaf, flower and root), the presence in the environment study of the feet of the tree (less abundant), and the stage of development (adult). *Detarium microcarpum* with an index of 2.67 was also considered a very vulnerable species in the Zou phytodistrict of Benin, by [19]. The average usage values vary between 8.42 (Kotafons) and 7.90 (Aïzos), indicate that according to socio-cultural groups, there is an uneven distribution of knowledge about the uses of the species. So the socio-cultural groups hold a diversity of endogenous knowledge on the use of the species [10].

Unlike the work of [8] who state that men have significantly more knowledge in the use of the species (ID = 5.44 bits) than women (ID = 2.10 bits), the present study reveals after eight (8) years that the knowledge are almost the same in kind. This difference in results can be justified by several reasons, including the size of the sample considered, which is fixed by phytodistrict, the increase in the interest that the fairer sex increasingly grants to knowledge related to the use of plants [22, 23], the presence of more in addition marked plants of the species in the houses to the detriment of its other habitats [8], etc. Apart from the medico-magical uses of the species which remain the well-guarded hunting ground of men, perhaps for reasons of culture, men and women have almost the same knowledge in the use of *C. albidum*. In clear, women have made great progress in the knowledge and use of the species.

The very high frequency of fruit use (FUO = 99.54%) of trees of the species shows that apart from the systematic collection of the fruits of the species for their commercialization [8], the populations also engage in harvesting and the cutting of the branches bearing them. However, according to the testimonies collected on the ground, the cutting the branches of these trees, profoundly reduces its yield. The wood is used for a frequency of 83.78% due to the aging of certain trees of the species and urbanization [7]. The felling of *C. albidum* trees is also linked to the fact that the wood would be a good wood for the manufacture of vegetable toothbrushes and for the manufacture economical coal. For some people, the older the tree, the less the fruits are numerous and of good quality. The irrational exploitation of the bark and roots of plants of the species for medicinal purposes, stunts their

growth and fruiting, when well even the disproportionate use of the leaves does not impact the development of the trees [24]. Ultimately, the ever-increasing reduction in density in trees of *C. albidum* in natural habitats is mainly due to poor management of resources of this very useful plant.

3.2.2 Prospects for the conservation of *Chrysophyllum albidum*

Rural populations are mostly poor, so they are greatly affected by the vulnerability of the income plants they exploit. In order to come to their aid, the State must take measures promoting the influence of agricultural activities, such as financial subsidies, technical supervision, access to water for all, and the fight against transhumance through the sensitization of breeders and farmers for the strict respect of animal circulation corridors. Thus, the local populations deliver more to fields and livestock, so that their unhealthy uses of resources phyto-genetics of forest trees are limited. Likewise, political leaders must make available land reserved for plantations of woody species whose that of *Chrysophyllum albidum* to prevent their felling for lack of space. In this framework, we can also experiment with these plantations in forests and other sacred places for solve the problem of poaching and premature killing. The scientific community must not only develop strategies for making juice from the ripe fruits of the species to avoid the massive loss of fruits due to lack of storage over a long period; but also work for the construction of seed banks of the good fruits of the species for promote the sowing of seeds likely to produce good quality fruit later.

4. CONCLUSION

This research has made it possible to identify the links between perceptions, uses and vulnerability or even the availability of *Chrysophyllum albidum* in Benin. Results inventoried, it appears that the species under study is used for its fruits which are used as food for humans; its wood exploited as energy wood and lumber; its leaves, bark and its roots prized for their medicinal and spiritual virtues. Unfortunately, the rural populations have daily access to the various organs of trees of the species through use of very unsavory practices, thus making this species very vulnerable. This vulnerability contributes dangerously to the loss of the species. Hence the need to design conservation strategies for this species, very useful in Benin.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ezebilo EE, Leif Mattsson. (). Socio-economic benefits of protected areas as perceived by local people around Cross River National Park, Nigeria. *Forest Policy and Economics*. 2010 ;12(3): 189-193.
2. Akouèhou GS, Houndonougbo A, Tenté B. The dynamics of production systems in the riparian agricultural terroirs of the Fita-Agbado intercommunal forest in Dassa-Zoumè and Savalou communes, Benin Central Hills Department. *Int. J. Biol. Chem. Sciences*. 2013;7(5):1877-1891.
3. Nwoboshi LC. The Nutrient Factor in Sustainable Forestry. Ibadan University Press, Ibadan, Nigeria. 2000;303.
4. Edem DO, Eka O, Ifon ET. Chemical evaluation of the nutritive value of the fruits of African star apple (*Chrysophyllum albidum*). *Food Chem*. 1984;14:303–311.
5. Houessou GL, Loubégnon OT, Gbesso GHF, Anagonou SL, Sinsin B. Ethnobotanical study of the African star apple (*Chrysophyllum albidum* G. Don) in the Southern Benin (West Africa). *Journal of Ethnobiology and Ethnomedicine*. 2012;8:40.
6. Dah-Dovonon JZ. Report from Benin. In Network Food Linear Species. Account of the first meeting of the Network: 11–13 December 2000. Edited by Eyog- Matig O, Gaoué OG, Dossou B. IPGRI: Ouagadougou, Burkina Faso. 2002; 2–19.
7. Gbesso GHF. Structure, population dynamics and socioeconomic and cultural importance of *Chrysophyllum albidum* on the Allada Plateau (Southern Benin). DEA memoir, Abomey-Calavi University, Abomey-Calavi. 2010 ;87.
8. Loubégnon OT, Anagonou L, Houessou GL, Gbesso GHF, Sinsin B. (2015). Socioeconomic importance of the organs (fruits, bark and roots) of the white star apple (*Chrysophyllum albidum* G. Don, 1831) on the Plateau of Allada. *Ann. Univ. de Lomé, sér. Lett., Tome*. 2015;XXXI-1: 294:157-165.
9. Gbesso FHG, Tenté BHA, Gouwakinnou NG, Sinsin BA. (2013a). Influence of climatic changes on the geographical distribution of *Chrysophyllum albidum* G. Don (Sapotaceae) in Benin. *Int. J. Biol. Chem. Sci*. 2007-20187(5).
10. Nguenang GM, Fedoung EF, Nkongmeneck BA. Importance of secondary forests for useful plant collection among the Badjoué of Eastern Cameroon. *Tropicultura*. 2010;28(4): 238-245.
11. Traoré L, Ouedraogo I, Ouedraogo A, Thiombiano A. Perceptions, uses and vulnerability of woody plant species resources in Southwest Burkina Faso. *Int. J. Biol. Chem. Sci*. 2011;5(1):258-278. DOI : 10.4314/ijbcs.v5i1.68103
12. Bonou A. Estimating the economic value of Nonlinear Forest Products (NFP) of vegetable origin in Sampéto village (Banikoara municipality). DEA memoir, Faculty of Agricultural Sciences, University of Abomey-Calavi, Bénin. 2008;66.
13. Lynam T, Cunliffe R, Mapaure I. Assessing the importance of woodland landscape locations for both local communities and conservation in Gorongosa and Muanza Districts, Sofala Province, Mozambique. *Ecology and Society*. 2004;9(4):1.
14. Betti JL. Vulnerability of plants used as antimalarials in Mintom district south of the Dja Biosphere Reserve (Cameroon). *Systematics and Geography of Plants*. 2001;71:661- 678.
15. Gbesso FGH, Loubégnon TO, Tenté BAH, Mensah G and Sinsin BS. Phytoecological and structural characterization of plant groups sheltering *Chrysophyllum albidum* (G. Don) on the Allada Plateau in Southern Benin. *Afrique Science*. 2013b;09(3)147 – 158.
16. Dagnelie P. Theoretical and applied statistics. Brussels: De Boeck Services, Brussels, Belgium ; 1998.
17. Alexiades MN. Selected guidelines for Ethnobotanical Research. A field manuel: The New York Botanical Garden; 1996.
18. Gomez-Beloz A. Plant use knowledge of the Winikina Warao: The case for questionnaires in ethnobotany. *Economy Botany*. 2002;56(3):231– 241.
19. Agbo IR, Missihoun AA, Vihotogbé R, Assogbadjo EA, Ahanhanzo C, Agbangla C. Impacts of traditional uses on the vulnerability of *Detarium microcarpum* Guill. & Perr. (Caesalpinaceae) in the Zou geographical photodistrict in Benin. *International Journal of Biological and Chemical Sciences*. 2017;11(2): 730-743.

20. Yaovi CR, Hien M, Kaboré SA, Séhoubou YJ, Somda I. Use and vulnerability of plant species and adaptation strategies of riparian populations in the Kou Classified Forest (Burkina Faso). *Int. J. Biol. Chem. Sci.* 2021;15(3):1140- 1157.
21. Loubelo E. Impact of non-timber forest products (NTFP) on the household economy and food security: case of the Republic of Congo. Doctoral thesis in economics. University Rennes 2. France. 2012;260.
22. Leach M. Gender and the Environment: traps and opportunities. *Development in Practice.* 1992;2:12-22.
23. FAO. Gender and Forestry Website ; 2006. Available:<http://www.fao.org/gender/en/fore-e.htm> [Accessed 07.03.2006].
24. Ayéna AC, Agassounon Djikpo Tchiboza M, Assogbadjo AE, Adoukonou-Sagbadja H, Mensah GA, Agbangla C, Ahanhanzo C. Uses and vulnerability of *Pterocarpus santalinoides* Her. ex from (Papilionoidae), a plant used in the treatment of gastroenteritis in southern Benin. *European Scientific Journal.* 2016;1 :1857–788.

© 2023 Alowanou-Kélé et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111614>