



Evaluation of Crop Establishment Methods and Nano Urea Versus Conventional Urea on Growth Indices in Rice Crop (*Oryza sativa* L.)

Kamaram Azam ^{a++*}, A.K. Singh ^{a#}, Vikash Singh ^{a++},
Manju Singh ^{a++}, Rajesh Kumar ^{a†} and Vishuddha Nand ^{a‡}

^a Department of Agronomy, Acharya Narendra Deva University of Agriculture and Technology, Ayodhya-224229, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i71010>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118243>

Original Research Article

Received: 10/04/2024

Accepted: 14/06/2024

Published: 17/06/2024

ABSTRACT

The present study aimed to evaluate Crop Establishment Methods and Nano Urea Versus Conventional Urea on Growth Indices in Rice Crop (*Oryza sativa* L.). Method of crop establishment influences the performance of rice through its effect on growth and development. Although, system of rice intensification has been reported to be the best establishment method. In traditional fertilizers, there is a huge loss of fertilizers by various manners like drift, leaching, runoff, microbial

⁺⁺ Research Scholar;

[#] Professor;

[†] Associate Professor;

[‡] Assistant Professor;

*Corresponding author: E-mail: anilkumarsinghmausam@nduat.org;

Cite as: Azam, Kamaram, A.K. Singh, Vikash Singh, Manju Singh, Rajesh Kumar, and Vishuddha Nand. 2024. "Evaluation of Crop Establishment Methods and Nano Urea Versus Conventional Urea on Growth Indices in Rice Crop (*Oryza Sativa* L.)". *Journal of Advances in Biology & Biotechnology* 27 (7):493-501. <https://doi.org/10.9734/jabb/2024/v27i71010>.

degradation, hydrolysis and photolysis. Bio accessibility of nutrients to plant is much lower relative to the rate of conventional fertilizers applied. The experiment was conducted during two consecutive seasons of *Kharif* 2022 and 2023 at the Agronomy Research Farm, Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya (U.P.), India. The nitrogen level had significantly affected on crop growth rate, relative growth rate and net assimilation rate at all the stages of crop growth during both the years of investigations. The maximum Net assimilation rate at 30-60 and 60-90 DAS/DAT and crop growth rate, relative growth rate at 30-60, 60-90 and 90 to harvest DAS/DAT was recorded under 60% RDN through conventional Urea + 40% RDN through Nano Urea which was at par with 40% RDN through conventional Urea + 60% RDN through Nano Urea treatment, while significant over rest of the treatment during both the years of investigations.

Keywords: Nano urea; conventional urea; crop establishment methods; bio accessibility; rice Crop.

1. INTRODUCTION

“Rice (*Oryza sativa* L.) is one of the most important cereal crops of the Poaceae family, it is said to have come from South-East Asia. In both regions of the temperate region and the tropics. A variety of agricultural ecologies, including irrigated uplands, rain-fed lowlands, and rice habitats that are vulnerable to flooding are used to grow rice. In terms of area and output, India is the second-largest producer of rice in the world after China. Rice is cultivated World-wide over an area of about 167.24 million ha with an annual production of about 756.74 million tonnes and productivity 4.60 tone ha⁻¹ in 2020-2021” [1]. “Rice is the most important staple food in Asia, providing on an average 32 % of total calorie uptake because of growing population, the demand for rice is expected to increase in the coming decades” [2].

“Method of crop establishment influences the performance of rice through its effect on growth and development” [3-5]. “Although, transplanting has been reported to be the best establishment method but due to high water and labour requirement, some alternatives like dry and wet direct seeding are being explored to ensure optimum yield at a lower cost. The results revealed that both transplanted and direct seeded method needed nearly equal investment on cultivation, but transplanted rice required more initial expenditure as compared to direct seeded rice” [6]. Nanotechnology is one of the cornerstone technologies in the 21st century. In traditional fertilizers, there is a huge loss of fertilizers by various manners like drift, leaching, runoff, microbial degradation, hydrolysis and photolysis. Bio accessibility of nutrients to plant is much lower relative to the rate of conventional fertilizers applied. It also gives rise to air and water adulteration. Nutrient use efficiency for standard fertilizers rarely surpasses 30-40% i.e.,

NUE of conventional nitrogen (30-35%), phosphorus (18-20%), and potassium (35-40%). Shang et al. [7]. Nano fertilizer nutrient use efficiency (NUE) is high as it is delivered to the site according to the crop demand. Nano fertilizer nutrient release character is according to the crop absorption rate, unlike conventional fertilizer in which nutrient release rate is substantial than crop absorption rate, causing low fertilizer use efficiency and environmental jeopardy [7].

2. MATERIALS AND METHODS

The experiment was conducted during two consecutive seasons of *Kharif* 2022 and 2023 at the Agronomy Research Farm, Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya (U.P.) which is situated on Ayodhya-Raebareli Road at the distance of 42 km from Ayodhya district head quarter. The experimental site falls under sub-tropical conditions with remarkable humidity and lies between 24.40 North latitude and 82.12° East longitudes with an altitude 113 meters above mean sea level. The experimental field was well leveled having good irrigation and drainage facilities. The source of irrigation was tube well. The experiment was layout in split plot design (SPD) with three replications taking three crop establishment methods direct seeded rice (DSR), System of Rice Intensification (SRI), transplanting rice in main plot and seven nitrogen level N₁: Control, N₂: 100% RDN through conventional Urea, N₃: 100% RDN through Nano Urea, N₄: 80% RDN through conventional Urea + 20% RDN through Nano Urea, N₅: 60% RDN through conventional Urea + 40% through Nano Urea, N₆: 40% RDN through conventional Urea + 60% through Nano Urea and N₇: 20% RDN through conventional Urea + 80% through Nano Urea in sub plot. Soil was sampled before sowing and after harvest of the crop to know the fertility

status of the experiment field. The growth analysis was done as per standard procedures

2.1 Crop Growth Rate

It represents the dry weight gained by a unit area of crop in unit time. The crop growth rate (CGR) was estimated by using the formula suggested by Watson [8] and expressed as $g\ m^{-2}\ day^{-1}$.

$$CGR = \frac{1}{A} + \frac{W_2 - W_1}{T_2 - T_1}$$

Where,

A is area, W_1 and W_2 Whole plant dry weight at T_1 and T_2 time, respectively.

2.2 Relative Growth Rate

The parameter indicates rate of growth per unit dry mater. It is similar to compound interest wherein the increment in any interval adds to the capital for subsequent growth. This rate of increment is known as relative growth rate [9]:

$$RGR = \frac{\text{Log}_e W_2 - \text{Log}_e W_1}{t_2 - t_1}$$

Where,

W_1 = Weight of dry matter (g) at time t_1
 W_2 = Weight of dry matter (g) at time t_2
 $t_2 - t_1$ = The interval in days

loge = Natural logarithms (Logarithms to the base of 2.3026)

Relative growth rate is expressed in $g/plant/day$

2.3 Net Assimilation Raate

It is an indirect expression of net photosynthesis. It represents the dry matter produced per square meter of leaf area per day). NAR is calculated by using the formula as suggested by Gregory [10] and expressed as mass unit-1 leaf area present per unit time ($g\ cm^{-2}\ day^{-1}$).

$$NAR = \frac{W_2 - W_1}{t_2 - t_1} \times \frac{\text{Log}_e LA_2 - \text{Log}_e LA_1}{LA_2 - LA_1}$$

Where,

W_1 and W_2 is dry weight of whole plant at time t_1 and t_2 respectively. LA_1 & LA_2 is the leaf area at times T_1 and T_2 respectively.

3. RESULTS AND DISCUSSION

3.1 Crop Growth Rate

Crop Growth rate was significantly affected due to crop establishment methods and nitrogen level in Table 1 and depicted in Fig. 1a and 1b. Crop growth rate of rice increased with crop age up to 60-90 DAS/DAT and there after declined in the present investigation.




Fig. 1a. Effect of different crop establishment methods and nitrogen levels on crop growth rate ($g\ m^{-2}\ day^{-1}$) of rice during 2022




Fig. 1b. Effect of different crop establishment methods and nitrogen levels on crop growth rate (g m⁻² day⁻¹) of rice during 2023

Table 1. Crop growth rate (g m⁻² day⁻¹) of rice as influenced by crop establishment methods and nitrogen levels

Treatments	Crop growth rate (g m ⁻² day ⁻¹)					
	30-60		60-90		90 - At harvest	
	2022	2023	2022	2023	2022	2023
Methods of crop establishment						
M ₁ : Direct Seeded Rice	10.67	10.81	9.73	9.87	5.98	5.64
M ₂ : System of Rice Intensification (SRI)	18.58	18.82	10.40	10.55	6.35	6.17
M ₃ : Transplanted Rice	16.86	17.17	10.23	10.43	6.01	5.97
SEm±	0.233	0.23	0.14	0.14	0.07	0.05
CD at 5%	0.94	0.94	0.56	0.58	0.29	0.24
Nitrogen Levels						
N ₁ : Control	10.09	10.04	10.05	8.72	4.89	4.85
N ₂ : 100% RDN through conventional Urea	14.68	14.78	10.12	10.43	5.43	5.41
N ₃ : 100% RDN through Nano Urea	14.99	15.00	10.19	10.66	5.58	5.55
N ₄ : 80% RDN through conventional Urea + 20% RDN through Nano Urea	15.35	16.81	10.15	10.52	5.87	5.79
N ₅ : 60% RDN through conventional Urea + 40% RDN through Nano Urea	18.65	18.60	11.12	11.63	7.25	7.05
N ₆ : 40% RDN through conventional Urea + 60% RDN through Nano Urea	17.58	17.71	10.68	11.18	6.53	6.49
N ₇ : 20% RDN through conventional Urea + 80% RDN through Nano Urea	16.26	16.23	10.22	10.84	6.18	5.96
SEm±	0.45	0.46	0.22	0.23	0.33	0.33
CD at 5%	1.29	1.31	0.63	0.65	0.94	0.94

Data further revealed that maximum crop growth rate (18.58 and 18.82, at 30-60 DAS/DAT during 2022 and 2023 respectively) recorded under SRI method and (10.40 and 10.55, 6.35 and 6.17 at 60-90 and 90-120 DAS/DAT during 2022 and

2023 respectively) recorded under SRI method which was at par with transplanted rice and direct seeded rice at 60-90 and 90-120 DAS/DAT while significantly higher than rest of the treatments during both years.

The nitrogen level had significantly affected on crop growth rate at all the stages of crop growth during both the years of investigations. Higher crop growth rate (18.65 and 18.60, 11.12 and 11.63, 7.25 and 7.05, 17.58 and 17.71, 11.68 and 11.18, 6.53 and 6.49) was noted with 60% RDN through conventional Urea + 40% RDN through Nano Urea treatment, which was at par with 40% RDN through conventional Urea + 60% RDN through Nano Urea and significantly superior over rest of the treatment at various interval of crop growth during both the years of investigations. Although, lowest crop growth rate was observed under control treatment at all the stages during both the years of investigation. The reason for this is that nano-urea increases the formation of chlorophyll, and the rate of photosynthesis leads to overall growth in the plant, which may lead to the formation of a greater number of leaves). "might due to the increased in leaf area, leaf number and vegetative growth of plants which increasing the photosynthetic activity; consequently, the higher dry matter produced and increased crop growth rate (CGR). The increase in CGR ultimately increases the total dry matter" [11,12].

3.2 Relative Growth Rate

The data on rice relative growth rate recorded at 30-60, 60-90 and 90-120 DAS/DAT for both the cropping seasons has been presented in Table 2 and depicted in Fig. 2a and 2b clearly indicated that establishment methods and nitrogen level

had significant effect on relative growth rate at all stages of crop growth during both the years of experiment.

Data further revealed that maximum relative growth rate (39.85 and 39.81 at 30-60 DAS/DAT respectively) recorded under SRI method which was at par with transplanted rice during both years and (14.43 and 14.44, 8.41 and 8.23 at 60-90, 90-120 DAS/DAT during 2022 and 2023 respectively) recorded under DSR method which was significantly higher than rest of the treatments during both years.

The nitrogen level had significantly affected in respect of relative growth rate at various intervals of crop growth during both the years, except between 90-120 DAS/DAT. The maximum relative growth rate was recorded under control between 60-90, 90-120 DAS/DAT, which was at par with N₁, N₂ and N₃ between 60-90 DAS/DAT during both the years of investigation. While, between 30-60 DAS/DAT the maximum relative growth rate (39.70 and 39.33 g g⁻¹ day⁻¹) was recorded with SRI method which significantly superior over rest of treatments. This might be due to higher plant height, leaf area index with maximum number of tillers contributed to the growth parameters of rice resulted in higher relative growth rate. As the result of maximum number of leaves resulted in high Leaf area index which will harvest maximum solar radiation within the canopy resulting in production of high dry matter in crop. The results are in close conformity [13,14].




Fig. 2a. Effect of different crop establishment methods and nitrogen levels on relative growth rate (g g⁻¹ day⁻¹) of rice during 2022




Fig. 2b. Effect of different crop establishment methods and nitrogen levels on relative growth rate (g g⁻¹ day⁻¹) of rice during 2023

Table 2. Relative growth rate (g g⁻¹ day⁻¹ x 10⁻³) of rice as influenced by crop establishment methods and nitrogen levels.

Treatments	Relative growth rate (g g ⁻¹ day ⁻¹ x 10 ⁻³)					
	30-60		60-90		90 - At harvest	
	2022	2023	2022	2023	2022	2023
Methods of establishment						
M ₁ : Direct Seeded Rice	28.75	28.82	14.43	14.44	8.41	8.23
M ₂ : System of Rice Intensification (SRI)	39.85	39.81	11.15	11.17	6.92	6.96
M ₃ : Transplanted Rice	38.13	38.18	11.66	11.70	6.79	6.64
SEm±	0.52	0.51	0.17	0.17	0.10	0.10
CD at 5%	2.08	2.07	0.68	0.69	0.41	0.41
Nitrogen Levels						
N ₁ : Control	27.83	27.52	14.44	14.59	8.85	8.64
N ₂ : 100% RDN through conventional Urea	34.66	34.62	13.78	13.81	6.33	6.15
N ₃ : 100% RDN through Nano Urea	35.28	35.09	13.87	13.93	7.99	7.89
N ₄ : 80% RDN through conventional Urea + 20% RDN through Nano Urea	35.77	37.38	12.29	10.34	7.14	7.30
N ₅ : 60% RDN through conventional Urea + 40% RDN through Nano Urea	39.70	39.33	11.24	11.43	7.03	6.93
N ₆ : 40% RDN through conventional Urea + 60% RDN through Nano Urea	38.75	38.62	10.86	11.16	6.84	6.88
N ₇ : 20% RDN through conventional Urea + 80% RDN through Nano Urea	37.06	36.66	11.80	12.44	7.43	7.14
SEm±	0.79	0.79	0.27	0.27	0.16	0.16
CD at 5%	2.27	2.27	0.78	0.82	0.46	0.45

3.3 Net Assimilation Rate

The data on rice net assimilation rate recorded at 30-60 and 60-90 DAS/DAT for both the cropping seasons has been presented in Table 3 and

depicted in Fig. 3a and 3b clearly indicated that establishment methods and nitrogen level had significant effect on net assimilation rate at all stages of crop growth during both the years of experiment.




Fig. 3a. Effect of different crop establishment methods and nitrogen levels on net assimilation rate (g m⁻² day⁻¹) of rice during 2022




Fig. 3b. Effect of different crop establishment methods and nitrogen levels on net assimilation rate (g m⁻² day⁻¹) of rice during 2023

Data further revealed that maximum net assimilation rate (3.45 and 3.50, 2.21 and 2.14 at 30-60 and 60-90 DAS/DAT during 2022 and 2023 respectively) recorded under SRI method which was at par with transplanted rice (3.31 and 3.35, 2.11 and 2.07 at 30-60 and 60-90 DAS/DAT during 2022 and 2023 respectively) and was found significantly superior over direct seeded rice at 30-60 and 60-90 DAS/DAT during both the years [15].

The nitrogen level had significantly affected on net assimilation rate at all the stages of crop

growth during both the years of investigations. The maximum Net assimilation rate at 30-60 and 60-90 DAS/DAT was recorded under 60% RDN through conventional Urea + 40% RDN through Nano Urea which was at par with 40% RDN through conventional Urea + 60% RDN through Nano Urea treatment, while significant over rest of the treatment during both the years of investigations. Increased in net assimilation rate enhances photosynthetic capacity of leaves with improved nutrition of the plants thereby increasing dry matter accumulation at final harvest. The results are in close conformity [16,17].

Table 3. Net assimilation rate (g m⁻² day⁻¹) of rice at different growth stages as affected by crop establishment methods and nitrogen levels

Treatments	Net Assimilation Rate (g m ⁻² day ⁻¹)			
	30-60 DAS/DAT		60-90 DAS/DAT	
	2022	2023	2022	2023
Methods of establishment				
M ₁ : Direct Seeded Rice	2.92	2.98	1.69	1.62
M ₂ : System of Rice Intensification (SRI)	3.45	3.50	2.21	2.14
M ₃ : Transplanted Rice	3.31	3.35	2.11	2.07
SEm±	0.046	0.047	0.029	0.028
CD at 5%	0.18	0.19	0.12	0.11
Nitrogen Levels				
N ₁ : Control	2.69	2.75	1.45	1.38
N ₂ : 100% RDN through conventional Urea	3.14	3.19	1.90	1.83
N ₃ : 100% RDN through Nano Urea	3.25	3.31	2.01	1.94
N ₄ : 80% RDN through conventional Urea + 20% RDN through Nano Urea	3.19	3.24	1.95	1.88
N ₅ : 60% RDN through conventional Urea + 40% RDN through Nano Urea	3.53	3.58	2.28	2.21
N ₆ : 40% RDN through conventional Urea + 60% RDN through Nano Urea	3.38	3.43	2.17	2.12
N ₇ : 20% RDN through conventional Urea + 80% RDN through Nano Urea	3.29	3.34	2.05	1.98
SEm±	0.071	0.072	0.044	0.042
CD at 5%	0.20	0.21	0.13	0.12

4. CONCLUSIONS

It is concluded that, 60% RDN through conventional Urea + 40% RDN through Nano Urea which was at par with 40% RDN through conventional Urea + 60% RDN through Nano Urea treatment for different crop establishment methods and nitrogen levels was found better for all growth indices crop growth rate (CGR), relative growth rate (RGR), and net assimilation rate (NAR) under transplanted rice.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. FAO F. Food and Agriculture Organization of the United Nations; 2021
Available: <https://www.fao.org/faostat/en/#data>
2. Pingali PL, Hossain M, Gerpacio RV. Asian rice bowls: The returning crisis. *Int. Rice Res. Inst*; 1997.
3. Abisankar MS, Augustine R, Manuel RI, Balaganesh B, Kumar D. Study of Physiological Growth Indices and Yield of Chickpea (*Cicer arietinum* L.) to Soil and Foliar Application Through Integrated Nutrient Management Practices. *Asian Journal of Soil Science and Plant Nutrition*, 2024;10(2):190–197.
Available: <https://doi.org/10.9734/ajsspn/2024/v10i2275>
4. Baboo K, Nand V. Effect of Crop Establishment Methods and Fertility Management on Growth Parameters of Rice (*Oryza sativa* L.). *International Journal of Plant & Soil Science*. 2023;35(16): 24–32.
Available: <https://doi.org/10.9734/ijpss/2023/v35i163125>
5. Verma KK, Song XP, Degu HD, Guo DJ, Joshi A, Huang HR, Xu L, Singh M, Huang DL, Rajput VD, Li YR. Recent advances in nitrogen and nano-nitrogen fertilizers for sustainable crop production: A mini-review. *Chemical and Biological Technologies in Agriculture*. 2023;10(1):111.
6. Singh KM, Pal SK, Verma UN, Thakur R, Singh MK. Effect of time and methods of

- planting on performance of rice cultivars under medium land of Bihar plateau. Indian Journal of Agronomy. 1997;(3):443-445.
7. Shang Y, Hasan MK, Ahammed GJ, Li M, Yin H, Zhou J. Applications of nanotechnology in plant growth and crop protection: A review. *Molecules*. 2019;24(14):2558.
 8. Watson DJ. Leaf growth in relation to crop yield. In: *The Growth of Leaves* (Ed.F.L. Milthorpe). Butterworth, U.K. 1956:178–191.
 9. Fisher RA. *Statistical method for research works*. Oliver and Boyd and Co. Inc. Endinburgh, United Kingdom; 1921.
 10. Gregory FG. Physiological conditions in cucumber houses. Third Annual Report, Experimental and Research Station, Cheshunt. 1917.
 11. Gulser F. Effect of ammonium sulphate and urea on NO₃ and NO₂ accumulation nutrient contents and yield criteria in spinach. *Scientia Horticulturae*. 2005;106: 330-340.
 12. Kumar P, Singh G, Singh PD, Singh T, Singh A, Lakra K. Effect of different crop establishment methods and nitrogen levels on the performance of kharif season rice (*Oryza sativa* L.) in the Indo-Gangatic plains of eastern Uttar Pradesh. *Crop Research*. 2021;56(1and2):1-7.
 13. Baloch MS, Awan IU, Hassan G, Zubair M. Studies on plant population and stand establishment techniques for increasing productivity of rice in Dera Ismail Khan, Pakistan. *Rice Science*. 2007;14(2):118-124.
 14. Tripathi AM, Pohanková E, Fischer M, Orság M, Trnka M, Klem K, Marek MV. The evaluation of radiation use efficiency and leaf area index development for the estimation of biomass accumulation in short rotation poplar and annual field crops. *Forests*. 2018;9(4):168.
 15. Blackman VH. The compound interest law and plant growth. *Annals of botany*. 1919;33(131):353-360.
 16. Ahmad N, Ahmad R, Bokhari S, Ghani A. Physiological determinants of growth and yield in wheat as affected by different levels of nitrogen and phosphorous. *Pakistan J. Agri. Sci*. 1990; 27:390-404.
 17. Meena MAMTA, Patel MV, Das TANIA, Verma HP. Effect of organic sources and nitrogen levels on growth and yield of kharif rice (*Oryza sativa* L.) under SRI technique. *Agriculture for Sustainable Development*. 2014;2(1):39-42.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/118243>