



# **Gonado Somatic Index in G5 Transgenic Mutiara Catfish (*Clarias gariepinus*) at Different Temperatures**

**Ibnu Dwi Buwono <sup>a\*</sup>, Iskandar <sup>a</sup>, Ine Maulina <sup>a</sup>  
and Ulfah Novia Suhendar <sup>a</sup>**

<sup>a</sup> *Department of Aquaculture, Faculty of Fisheries and Marine Sciences, Universitas Padjadjaran, Indonesia.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119639>

**Original Research Article**

**Received: 10/05/2024**

**Accepted: 09/07/2024**

**Published: 16/07/2024**

## **ABSTRACT**

Temperature is one of the factors that can affect the maturity of the gonads in fish. This study aims to find out what is the best temperature for maturation of broodstock to prepare G5 transgenic mutiara catfish broodstock that are ready to hatch based on gonado somatic index (GSI) performance, relative fecundity, fertilization rate (FR), hatching rate (HR). The research was carried out in the Hatchery Building 4, Fisheries Biotechnology Laboratory, Faculty of Fisheries and Marine Sciences. The method used in this research is an experimental method with a complete randomized design (CRD), which consists of four treatments and three replicates. Treatment A: 26

\*Corresponding author: E-mail: [ibnu.dwi.buwono@unpad.ac.id](mailto:ibnu.dwi.buwono@unpad.ac.id), [0812ibnu@gmail.com](mailto:0812ibnu@gmail.com);

$\pm 0.5$  °C, B :  $28 \pm 0.5$  °C, C :  $30 \pm 0.5$  °C for transgenic catfish, C\*:  $30 \pm 0.5$  °C (for nontransgenic catfish) with maintenance for 8 weeks. The reproductive performance of the female broodstock of G5 transgenic mutiara catfish with the highest relative fecundity was found in treatment B of kg/brood. The best temperature for gonad maturation was obtained in treatment B with a temperature of  $28 \pm 0.5$  °C with a gonadosomatic index value of  $52.42 \pm 5.93\%$  for females and  $5.22 \pm 0.17\%$  for males. The spawning performance in the parent pair of treatment B produced the best relative fecundity, fertilization rate (FR), and hatching rate (HR), with values of 89,179 eggs/kg,  $84.98 \pm 5.01\%$ , and  $90.87 \pm 3.12\%$ , respectively.

**Keywords:** Gonad maturation; transgenic catfish; temperature manipulation; G5.

## 1. INTRODUCTION

Catfish culture activities are one of the cultivation in the fisheries sector that continues to experience rapid development. This is because catfish is one of the fish that is in great demand by the community. Natural spawning is the spawning that is most carried out by culture. However, this natural spawning will produce a smaller quantity of fish fry than artificial and semi-artificial spawning, besides that natural spawning also tends to experience many failures. Spawning by utilizing hormones and natural compounds has been widely carried out and has shown significant results [1]. The hormone ovaprim, which is often used for catfish spawning, has also been shown to significantly increase ovulation [2]. Problems that occur in natural spawning occur because natural spawning depends a lot on the readiness of mature broodstock and generally only occurs in certain seasons.

Catfish spawning using hormones can be done to make the spawning process easier. The use of hormones can make it easier for farmers to be able to produce fingerlings outside of the spawning season, where the fish to be spawned have gone through the previous maturation process until they are ready to spawn [3]. The maturation process in the broodstock that is not optimized will make it difficult for the fish to mature the gonads at a certain time. One of the efforts to optimize the broodstock can be done by environmental manipulation.

One of dominant factors that influence fish gonadal maturity is temperature. In addition, temperature can also affect the reproductive activity of fish because it can increase or decrease the weight of the gonads. Previous research to related environmental factors that have a significant role in influencing the development of fish gonads are temperature, food, light, and season [4]. In the maturation

process, the ideal temperature to maintain gonad maturation in catfish groups is 27-30°C [5]. Maturation must be accompanied by spawning induction, because maturation can be considered successful if the broodstock that is matured with temperature can be induced to spawn. The indicator can determine how much temperature affects the fecundity that will produce the number of eggs and larvae to be produced.

The increase in egg production is shown by fecundity. The quality of fecundity depends on a good broodstock strain. Transgenic mutiara catfish (*Clarias gariepinus*) strains produced from the use of transgenic technology (addition of exogenous GH, *Clarias gariepinus* Growth Hormone, CgGH) have higher reproductive performance than non-transgenic catfish. The propagation of transgenic strain catfish through the breeding process has produced the transgenic mutiara catfish G5 [6,7]. Research showed by the transgenic G1 and G2 mutiara catfish could produce more catfish fingerlings than non-transgenic catfish [8]. Currently, transgenic mutiara catfish have reached the fifth generation (G5). Therefore, it is necessary to carry out research on transgenic and non-transgenic fish with different temperature variations.

## 2. MATERIALS AND METHODS

The research was carried out in the Hatchery Building 4, Fisheries Biotechnology Laboratory, Faculty of Fisheries and Marine Sciences, Padjadjaran University. The time for conducting the research is from May to August 2023 which starts from preparation, implementation, and observation.

The tools used for the research are, round fiber tubs, heaters, thermometers, aeration installations, water pumps, brushes, basins, scales, bars, sterofoam, injections, kakaban, black plastic, containers, cameras, plastic

containers, label paper. The ingredients used are test animals, Prima Feed (PF)128 broodfish feed, ovaprim hormone, aquabides, chorulon hormone (hCG, human chorionic gonadotropin).

The method used in the study was an experimental method with a Complete Random Design (CRD) consisting of four treatments and three replicates. The temperature treatments used were as follows: A (26 ±0.5 °C), B (28 ±0.5 °C), C (30 ±0.5 °C) for G5 transgenic mutiara catfish and C\* (30 ±0.5 °C) for non-transgenic catfish as a control to compare with fish that do not have growth hormon. The parameters used during the research were as follows :

#### a. Gonado Somatic Index (GSI)

The success of the gonad maturation process can be presented from the Gonado Somatic Index (GSI) value which reflects the growth of female or male gonads. The gonadal maturation rate calculated using the GSI value reflects whether the parent can be castrated or not. The calculation of the GSI value is calculated using the following formula according to [9,10]:

$$GSI = \frac{Gw}{Bw} \times 100\%$$

#### Information:

GSI = Gonado Somatic Index (%)  
 Gw = Gonadal weight (g) (weight before spawning – weight before after spawning)  
 Bw = Body weight (g)

#### b. Relative Fecundity (RF)

Fish fecundity can be calculated in several ways, namely: the number method, the volumetric method and the gravimetric method. This study uses the gravimetric method [11] in the following way:

$$RF = \frac{Wg}{We} \times 100\%$$

#### Information:

RF = Relative fecundity  
 Wg = weight gonad (g)  
 We = Mean weight of egg sampel (g)

#### c. Fertilization Rate (FR)

The rate of fertilization is calculated as the percentage of the number of eggs fertilized

(undergoing the embryogenesis process) compared to the number of eggs from spawning calculated on a sieve as a container for treatment samples. Fertilized catfish eggs are yellow-green, while unfertilized eggs are milky white or pale. Fertilized eggs are counted after the fertilization process. The rate of fertilization is calculated using the formula, as follows [12]:

$$FR = \frac{\sum \text{fertilized eggs}}{\sum \text{total egg}} \times 100\%$$

#### d. Hatching Rate (HR)

The rate of hatching is calculated as a percentage of the number of eggs hatched compared to the number of fertilized eggs. The calculation of hatching rate is carried out 18 - 24 hours after the fertilization process (at the water temperature of the hatching medium ranges from 29°C-30°C). The rate of hatching is calculated using the formula [12], as follows:

$$HR = \frac{\sum \text{fertilized eggs}}{\sum \text{total egg}} \times 100\%$$

#### e. Data Analysis

The data from the research results were analyzed quantitatively using one way ANOVA with advanced tests Duncan's Multiple Range Test (Sigma plot 14.5)

## 3. RESULTS AND DISCUSSION

### 3.1 Female Gonado Somatic Index

Observation of the Gonado Somatic Index (GSI) for eight weeks in female G5 mutiara and non-transgenic catfish in the form of percentages (%) is presented in (Fig. 1).

The results of gonad maturation with temperature treatment showed that the highest female gonado somatic index value was found in treatment B (temperature 28 ± 0.5 °C) with a gonado somatic index value of 52.42 ± 5.93% followed by treatment A (temperature 26 ± 0.5 °C) with a gonado somatic index value of 43.54 ± 8.55%, treatment C (temperature 30 ± 0.5 °C) with a GSI value of 42.39 ± 10.60%, while the lowest GSI value of the female is found in the treatment C\* (temperature 26 ± 0.5 °C) by producing a gonado somatic index (GSI) value of 19.02 ± 2.28%.




Fig. 1. Graph of GSI values of transgenic mutiara catfish G5 (A,B,C) and non-transgenic (C\*). The mean followed by different letter notations showed significance ( $p < 0.05$ )




Fig. 2. GSI value chart of male transgenic mutiara catfish G5 (A,B,C) and non-transgenic (C\*). The mean followed by different letter notations showed significance ( $p < 0.05$ )

The growth of the gonads of transgenic mutiara catfish can be optimized by using temperature. In Lin's research, the temperature of 28°C has the best influence on gonadal growth [13]. The high GSI value in treatment B females is one of the reasons for the use of the optimum maintenance temperature of 28°C. In addition, the rapid growth of fish can affect gonadal growth. Transgenic mutiara catfish in treatment B had the highest GSI value also influenced by the presence of exogenous GH inserts. Transgenesis technology has been carried out [14] to produce G0 transgenic mutiara catfish through transgenesis technology by increasing growth by two and a half times (125%) compared to non-transgenic catfish. Exogenous GH inserts have a positive effect on the reproduction of G1 transgenic mutiara catfish, namely it can increase gonadal fertility, increase egg size and increase the production of estradiol hormone [15].

Temperature fluctuations are thought to have an effect on the CgGH excretion level. The temperature of 28°C produced the highest GSI value because at 28°C it was related to a high level of expression which when compared to treatment A (26°C) and treatment C (30°C). And vice versa, the low GSI value obtained at temperature 26 is suspected to be due to a decrease in the expression level of CgGH. Rapid growth can affect the value of GSI. The effect of CgGH on fish gonad growth related to the induction of exogenous GH stimulates the secretion of IGF-1 growth factor which works in cooperation with GH and steroids (estrogen, E2 or testosterone, T) to multiply gamete cells (sperm or ovum). The multiplication of these gamete cells leads to an increase in gonadal growth as a consequence of which the GSI value increases at the optimum temperature of 28°C. Previous research on the G1 transgenic mutiara catfish, showed that the growth of the gonads of transgenic fish was higher when the broodstock was kept in the temperature range of 27-28°C [16].

### 3.2 Male Gonado Somatic Index

The observation results during 8 weeks of maintenance can be seen in (Fig. 2)

The highest male gonado somatic index value was found in treatment B (temperature 28±0.5°C) with a gonado somatic index value of 5.22 ± 0.17%, followed by treatment C (temperature 30±0.5°C) with a gonado somatic index value of

5.13 ± 0.36, and treatment A (temperature 26±0.5°C) of 4.92±0.37%. Meanwhile, the C\* treatment (temperature 30±0.5) obtained the lowest GSI value with a male parent GSI value of 1.95 ± 0.96%. The difference in treatment A, B, C with treatment C\* is suspected because in transgenic mutiara catfish there is an insert of the GH gene [16]. Exogenous GH has a positive effect on the reproduction of G1 transgenic mutiara catfish, namely it can increase gonadal fertility, increase egg size and increase the production of estradiol hormone [15]. The results of the research that have been carried out show that the gonado somatic index value in the male broodstock of transgenic mutiara catfish is better compared to the gonado somatic index value in non-transgenic catfish. The gonado somatic value of the male brood index in non-transgenic catfish obtained is still relatively good because it has a value of 1.95 ± 0.96%. The gonado somatic value of the male broodstock index of mutiara catfish ranged from 0.22%-1.47% [17]. Meanwhile, the GSI value of male catfish in the waters of the Nile is reported to be around 1% [18]. The GSI value of male broodstock in the G4 transgenic mutiara catfish research got the best score of 3.55 ± 0.24% [19]. In this study, the G5 transgenic mutiara catfish had a gonado somatic index value of 5.22 ± 0.17%. This value is much higher compared to non-transgenic catfish. A significant difference between transgenic mutiara catfish and non-transgenic mutiara catfish is suspected because transgenic mutiara catfish have GH gene inserts [17]. Exogenous GH has a positive effect on the reproduction of G1 transgenic mutiara catfish, namely it can increase gonad fertility [15]. GH in male fish spurs the secretion of testosterone produced by leydig cells which functions to carry out the process of spermatogenesis until it becomes spermatozoa or ripe sperm [20].

### 3.3 Relative Fecundity

The relative fecundity obtained from fish that successfully spawned during the 8 weeks of the study can be seen in Table 1.

Based on Table 1, the relative fecundity of the broodstock catfish of the G5 transgenic mutiara benita was highest in treatment B (♂ B1 X ♀ B3) of 89,179 egg/kg of broodstock, followed by treatment A (♂ A3 X ♀ A2) of 87,530 and treatment C (♂ C2 X ♀ C2) of 83,857. According to Putri research, transgenic mutiara catfish in the first offspring produced a relative fecundity of 83,333-84,000 egg/kg [15].

**Table 1. Relative fecundity**

Treatment	Relative Fecundity (egg/kg broodstock)
(A) ♂ A3 X ♀ A2	87,530
(B) ♂ B1 X ♀ B3	89,179
(C) ♂ C2 X ♀ C2	83,857

In the study, G4 transgenic mutiara catfish succeeded in spawning with a relative fecundity of 85,103 eggs/kg [19]. The results of the research on the spawning of G5 transgenic mutiara catfish, obtained the highest value of 89,179 egg /kg. This value is classified as a good number of fecundity [21] which states a good fecundity value for catfish, which is around 50,000-100,000 egg/kg of fish body weight.


The success of G5 transgenic mutiara catfish in spawning in a relatively faster time with a good fecundity value is because in the transgenic mutiara catfish there is a GH insert of *C. gariepinus* [14]. One of the factors that can affect the fecundity of fish is the age of the fish. The age of fish affects the development of the gonads, in male fish maturing the gonads generally experience gonadal maturation at an earlier time, namely 8 months [3]. Meanwhile, the gonads of female fish mature at the age of 10 months and the spawning cycle is once every 1.5 months [22,23]. Meanwhile, in female fish, the physical characteristics are the enlargement of

the abdomen and when striping is carried out, there are eggs that come out.


### 3.4 Fertilization Rate

The results of the fertilization rate in catfish broodstock that successfully spawned after eight weeks of rearing can be seen in Fig. 3.

Based on Fig. 3, the highest fertility value in G5 transgenic mutiara catfish was in treatment B, which was  $84.98 \pm 5.01$ , followed by treatment C of  $82.15 \pm 3.23$  and the lowest value was in treatment A of  $72.51 \pm 2.97$ . The value of the rate of fertilization is still relatively good, referring to the Indonesian National Standardization (SNI) [21] no. 01-6484.3.2000, catfish have a fertilization rate in the range of 70% to 90% [24]. The rate of fertilization that reaches a value above 70% is categorized as high [25]. Data analysis resulted in Fig. 4, the rate of fertilization between cross C and cross A and B does not have a significant difference, This is suspected because the gonad maturity level of



**Fig. 3. Graph of fertilization rate of G5 transgenic mutiara catfish eggs (A,B,C). The average followed by different letter notation shows significance ( $p < 0.05$ )**



**Fig. 4. Graph of hatching rate of G5 transgenic mutiara catfish eggs (A,B,C). The average followed by different letter notation shows significance ( $p < 0.05$ )**

the cross pair of G5 transgenic catfish broodstock is relatively uniform. Allegedly, this is the cause of the rate of hatching in each egg produced in the G5 transgenic catfish parent pair is relatively the same. Furthermore, in non-transgenic mutiara catfish that use temperature treatment, it does not cause maturation in the broodstock so that it causes spawning failure.

### 3.5 Hatching Rate

The following are the results of the rate of egg hatching that can be seen on (Fig. 4). Based on Fig. 4, the data on the rate of hatching of G5 transgenic mutiara catfish eggs shows that the highest value of hatching rate is in treatment B of  $90.87 \pm 3.12$  followed by treatment C of  $87.92 \pm 5.23$  and the lowest value is in treatment A of  $80.97 \pm 4.93$ . Hatching rate value Transgenic mutiara catfish is relatively high compared to sangkuriang catfish used in artificial spawning in Poland with a value of 66.14-68.50% [26] and in Nigeria which is  $84.15 \pm 2.05\%$  [27] and around 60-66% [28].

The rate of hatching in each treatment can be categorized as high according to the research of Hijriyanti [29] which states that the rate of hatching in the range of 30% - 50% is considered low, and  $> 60\%$  is considered high. The high

value of the rate of egg hatching is due to the maximum production of eggs. Stated that the success of hatching is influenced by the quality of the eggs and the number of fertilized eggs, if the maximum yield will also be maximum [30]. In addition, according to Marnis [31], the transgenic *Pangasianodon hypophthalmus* growth hormone (PhGH) present in transgenic fish is also influential in producing a high rate of hatchability, which is 80.68%, while non-transgenic fish is only 76.68% [32]. This is directly proportional to this study, where the G5 transgenic mutiara catfish obtained a result of above 80% with the best score of 90.87%.

### 4. CONCLUSION

The best temperature for gonad maturation was obtained in treatment B with a temperature of  $28 \pm 0.5$  °C with a gonadosomatic index value of  $52.42 \pm 5.93\%$  for females broodstock and  $5.22 \pm 0.17\%$  for male broodstock. The broodstock pair of treatment B also produced the best relative fecundity, fertilization rate (FR), and hatching rate (HR) values with sequential values of 89,179 egg/kg broodstock,  $84.98 \pm 5.01\%$ , and  $90.87 \pm 3.12\%$ .

In the next study, it was recommended to use a temperature of  $28 \pm 0.5$  °C to accelerate the maturation of the gonads in catfish broodstock.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Rukisah, Maulianawati D, Cahyadi . *In vitro* antibacterial efficacy of leaves extract of centela asiatica against *Vibrio harveyi* and *Aeromonas hydrophila*. Indonesian Aquaculture Journal. 2019;14 (2):69 – 74.
2. Sharaf SM. Effect of GnRHa, pimozone and Ovaprim on ovulation and plasma sex steroid hormones in African catfish *Clarias gariepinus*. Theriogenology. 2012;77(8): 1709-1716.
3. Buwono ID, Puteri A., dan Herman RG. The maturity of female gonads of G2 Transgenic Mutiara Catfish (*Clarias Sp.*). Asian Journal of Fisheries And Aquatic Research. 2021;12(3):9 – 19.
4. Scott JC. Accommodative institutions: A fisheries development approach that is oriented towards increasing the accessibility of community fisheries. Fisheries scientific bulletin. Faculty of Fisheries, Brawijaya University Malang; 1996.
5. Sundararaj BI. Reproductive physiology of teleost fishes, United Nations Development Program and FAO. Rome. 1981;82.
6. Buwono ID, Iskandar, Grandiosa R. Training on padjajaran mutiara catfish hatchery techniques at the Mina Sejahtera Sadaya Group, Cileunyi District, Bandung Regency. Journal of Science and Technology Applications for Society. 2023; 12(1):114–121.
7. Eze, Felix, Muhammad Yakubu Haruna, Akosubo Oghenemarho Victory. A Comparative study on the gonadosomatic index and milt volume of four populations of *clarias gariepinus* (Burchell, 1822) broodstock strains from North-East Nigeria. Asian Journal of Fisheries and Aquatic Research. 2022;17(5): 8-19. Available:https://doi.org/10.9734/ajfar/2022/v17i530415
8. Buwono ID, Puteri A, Herman R. The maturity of female gonads of G2 transgenic mutiara catfish (*Clarias Sp.*). Asian Journal Of Fisheries And Aquatic Research. 2021;12(3):9 – 19.
9. Mariskha PR, Abdulgani N. Aspects of reproduction of tiger grouper (*Epinephelus sexfasciatus*) in glondonggede waters, Tuban. Its Science and Art. 2012;1(1):27-31.
10. Flores A, Wiff R, Ganiyas K, Marshall CT. Accuracy of gonadosomatic index in maturity classification and estimation of maturity ogive. Fisheries Research. 2019 ;210:50-62.
11. Witthames R, Peter, Thorsen A, Murua H, Saborido-Rey F, Greenwood NL, Dominguez R, Korta M, Kjesbu S, Olav. Advances in methods for determining fecundity: Application of the new methods to some marine fishes. Fishery Bulletin. 2009;107(2):148 – 164.
12. Effendie MI. Fisheries biology. Nusantara Library Foundation; 2002.
13. Lin Q, Lu J, Gao Y, Shen L, Cai J, Luo J. The effect of temperature on gonads, embryonic development and survival rate of juvenile seahorses, Hippocampus of horses Bleeker. Aquaculture. 2006;254: 701–713.
14. Buwono ID, Iskandar, Agung MUK, Subhan U. Assembling transgenic catfish (*Clarias sp.*) using sperm construction electroporation technique. Journal of Biology. 2016;20(1):17 – 28.
15. Puteri A, Buwono ID, Herman RG, Iskandar. The maturity of female gonads of G2 transgenic mutiara catfish (*Clarias sp.*). Asian Journal of Fisheries and Aquatic Research, 2021;12 (3): 9–19.
16. Buwono ID, Junianto J, Iskandar I, Alimuddin A. Reproductive performance of transgenic mutiara catfish (G1) comprising the growth hormone gene. Journal of Biotech Research. 2019;10: 199–212.
17. Iswanto B, Imron H, Marnis, Suprpto R. Technical instructions for cultivating mutiara catfish. Fish Breeding Research Institute, Subang. 2014;55.
18. Ahmed YA, Samei NAA, Zayed AZ. Morphological and histomorphological structure of testes of the catfish *Clarias*



- gariiepinus* from Egypt. Pakistan Journal of Biological Sciences. 2013;16 (13): 624-629.
19. Buwono ID, Dewanti LP, Herman RG, Mariane, A. T. Use of ovaprim and Chorulon hormones for spawning induction of G4 transgenic Mutiara catfish Brood stock in rearing indoor Hatchery; 2023.
  20. Tokalov SV, Gutzeit HO. Spermatogenesis in testicular primary cell cultures of the tilapia (*Oreochromis niloticus*). Developmental Dynamics. 2005;233:1238 –1247.
  21. Indonesian National Standards (SNI). Dumbo Catfish (*Clarias sp*). National Standardization Agency. Jakarta. 2014;SNI 6484.3.
  22. Iswanto B, Suprpto R, Marnis H. Reproductive performance of mutiara catfish (*Clarias gariepinus*). Aquaculture Media. 2016;11(1):1–9.
  23. Du H, Zhang X, Leng X, Zhang S, Luo J, Liu Z, Qiao X, Kynard B, Wei Q. Gender and gonadal maturity stage identification of captive Chinese sturgeon, *Acipenser sinensis*, using ultrasound imagery and sex steroids. General and Comparative Endocrinology. 2017;245:36-43.
  24. Nainggolan C, Matling, Yusuf NS. Hatching degrees of dumbo catfish eggs incubated in different water media. Journal of Tropical Fisheries. 2023;18(1):8 - 16.
  25. Effendi, Prasetya, T. Maturation of Brood Gonads of Botia Fish (*Botia macracanthus*) in Ponds. Indonesian Aquaculture Journal. 2003;2 (2):51 – 54.
  26. Brzuska, E. Artificial propagation of African catfish (*Clarias gariepinus*): differences between reproductive effects after stimulation of ovulation with carp pituitary homogenate or GnRH-a and dopaminergic inhibitors. Czech Journal of Animal Science. 2003;48:181-190.
  27. Omitoyin BO, Adesehinwa AOK, Edibite LI. Reproductive performance and serum biochemistry of female *Clarias gariepinus* broodstock raised in pond effluent water. Tropical and Sub-tropical Agroecosystem. 2005;5:117 – 122.
  28. Gbemisola OB, Adebayo OT. Sperm quality and reproductive performance of male *Clarias gariepinus* induced with synthetic hormones (ovatide and ovaprim). International Journal of Fisheries and Aquaculture. 2014;6 (1):9-15.
  29. Hijriyati KH. Egg quality and early development of duck grouper larvae (*Cromileptes altivelis*, valenciennes (1928) in Air Saga Village, Tanjung Pandan, Belitung. Thesis. Faculty of Mathematics and Natural Sciences. University of Indonesia; 2012.
  30. Julianti. Technical instructions for goldfish cultivation. directorate general of fisheries. Jakarta; 2001.
  31. Marnis H, Iswanto B, Suprpto R, Imron I, Dewi RRSPS. Growth and cygosity of transgenic F-2 African catfish (*Clarias gariepinus*) Carrying the Siamese Patin Fish Growth Hormone Gene (*Pangasianodon hypophthalmus*). Journal of Aquaculture Research. 2015;10(2):161 – 168.
  32. Zhong CY, Song Y, Wang Y, Li Y, Liao L, Xie S, Zhu Z, Hu W. Growth hormone transgene effects on growth performance are inconsistent among offspring derived from different homozygous transgenic common carp (*Cyprinus carpio* L.). Aquaculture. 2012;356(357):404 – 411.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/119639>