



# Effect of Sowing Methods and Different Levels of Nitrogen on Yield Attributes and Yield of Maize

Bhanu Singh <sup>a++\*</sup>, Dhruvendra Singh Sachan <sup>b++</sup>,  
Sushil Kumar Yadav <sup>a++</sup> and Shabbeer Ahmad <sup>a++</sup>

<sup>a</sup> Department of Agriculture, Integral Institute of Agricultural Science and Technology, Integral University, Lucknow, U.P. - 226026, India.

<sup>b</sup> Department of Agronomy, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, U.P. – 208002, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i92859>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/122723>

**Original Research Article**

**Received: 01/07/2024**

**Accepted: 04/09/2024**

**Published: 05/09/2024**

## ABSTRACT

The field experiment was carried out at Research Farm, Integral University, Lucknow, Uttar Pradesh, India during Kharif season, 2021. The experiment envisages to study the effect of Sowing methods and different Nitrogen levels on yield and attributes of maize crop. The experiment was laid down in the Split Plot Design and the treatments was replicated three times. Among the sowing methods, the maximum length of a cob (18.52 cm) was obtained in the ridge sowing method (S1) while among nitrogen levels the maximum length of cob (19.51 cm) was obtained with the application of 25% N through granular urea + 75% N through Nano-urea (N5). Among nitrogen levels highest number of grains per row (40.76) were obtained in the application of 25% N through

<sup>++</sup> Research Scholar;

\*Corresponding author: E-mail: [bhanu6201@gmail.com](mailto:bhanu6201@gmail.com);

**Cite as:** Singh, Bhanu, Dhruvendra Singh Sachan, Sushil Kumar Yadav, and Shabbeer Ahmad. 2024. "Effect of Sowing Methods and Different Levels of Nitrogen on Yield Attributes and Yield of Maize". *Journal of Experimental Agriculture International* 46 (9):613-18. <https://doi.org/10.9734/jeai/2024/v46i92859>.

granular urea + 75% N through Nano-urea (N5). Seed Index was achieved to be highest in ridge method of sowing(S1) with the application of 25% N through granular urea + 75% N through Nano-urea (N5). Number of grains per cob were highest (385.55) in the N5 treatment among the nitrogen levels treatments. Among sowing methods, the maximum grain yield of 51.75 q ha<sup>-1</sup> was founded in the ridge method of sowing. Among nitrogen levels the highest grain yield of 57.76 q ha<sup>-1</sup> was obtained in N5 treatment. However, harvest index was found to be non-significant. Ridge method of sowing was found to be most effective than broadcasted method. Application of Nano-urea alone did not give good results. Granular urea along with nano-urea was found to be more effective in maize crop under this region.

*Keywords: Granular urea; ridge; treatments.*

## 1. INTRODUCTION

Maize, scientifically classified as *Zea mays* L., is a highly significant fodder and cereal crop renowned for its extensive availability and it belongs to the Poaceae family [1]. Globally, it ranks as the third most important cereal crop, following wheat and rice, and is celebrated for its exceptional production capabilities and adaptability to a variety of agroclimatic conditions, earning the title of 'Queen of Cereals' [2]. Worldwide, maize is cultivated across approximately 0.207 billion hectares, yielding about 1.22 billion Metric Tons with an average productivity of 5.89 tons per hectare, with United States being the largest producer (USDA, 2023-24).

In India, major maize-growing states are Karnataka, Madhya Pradesh, Bihar, Tamil Nadu, Telangana, Maharashtra and Andhra Pradesh. Production of maize is estimated to be 35.67 million tonnes in 2023-24 (3rd Advance Estimate, PIB). India has exported 14,42,671.48 metric tonnes of maize to the world for the worth of 443.53 USD Millions in 2023-24 (APEDA). For India major maize export destinations in 2023-24 were Vietnam, Nepal, Bangladesh, Malaysia and Thailand.

Sowing methods or land management system plays a major role in improving water and nutrient use efficiency of field crops. Land configuration increases water use efficiency and also increases availability of nutrients to crops Chiroma et al. [3]. The superiority of ridges and furrow system could be ascribed to proper drainage of excess water coupled with adequate aeration at the time of irrigation or heavy rainfall. Ridges and furrow method of sowing improved grain as well as stover yield of maize over the flat bed method of sowing corroborate the studies of Parihar et al. [4].

Among the plant nutrients, nitrogen (N) management is one of the most important factors required for improving crop productivity [5]. According to Guo et al. [6], Nitrogen is the most yield-restraining nutrient in crop production, globally.

## 2. MATERIALS AND METHODS

### 2.1 Experimental Site

The experiment was carried out at Research Farm, Integral University, Lucknow, Uttar Pradesh, India during Kharif, 2021. Lucknow has a humid subtropical climate with hot, sunny summers from March to May. The city receives an average of 835 millimetres of rainfall from the southwest monsoon winds between June to October. Summers are very hot with temperatures rising to 40-45 °C (104-113 °F) range.

### 2.2 Edaphic Condition

The soil in the experimental field was clayey in texture and slightly alkaline with pH of 7.8. Organic carbon in the soil was 0.31% which was estimated by rapid titration method given by Walkley and Black (1934). The available Nitrogen in soil was 138 kg ha<sup>-1</sup>, which was estimated by the Alkaline permanganate method given by Subbiah and Asija [7]. The available Phosphorus was 17.7 kg ha<sup>-1</sup> estimated by Olsen's method given by Olsen et al. [8]. The available K was 294.1 kg ha<sup>-1</sup> which was estimated by the Flame photometer method given by Jackson [9].

### 2.3 Experimental Design and Treatment Details

The experiment was designed as Split Plot Design. The treatments were replicated thrice.

**List 1. The details of the treatment are as follows**

<b>Treatments Details</b>	
<b>Main plots (Method of sowing)</b>	
<b>S1</b>	Ridge sowing
<b>S2</b>	Flat-bed sowing
<b>S3</b>	Broadcasting
<b>Sub plot (Nitrogen Management)</b>	
<b>N1</b>	100% N through granular urea
<b>N2</b>	100% N through Nano-urea
<b>N3</b>	75% N through granular urea + 25% N through Nano-urea
<b>N4</b>	50% N through granular urea + 50% N through Nano-urea
<b>N5</b>	25% N through granular urea + 75% N through Nano-urea

\*RDF= 120 N : 60 P<sub>2</sub>O<sub>5</sub> :60 K<sub>2</sub>O**2.4 Preparation of the Experimental Field and Application of Fertilizers**

The seed bed was prepared by ploughing with a cultivator followed by rotavator. The field was then manually laid out according to the plan, and *TMMH 826* Hybrid seeds were sown at a rate of 25 kg ha<sup>-1</sup> with a spacing of 45 cm × 20 cm. Granular Urea (46% N) and Nano urea (4% N) was applied as per the treatments, while DAP (46% P<sub>2</sub>O<sub>5</sub>) and MOP (60% K<sub>2</sub>O) were uniformly applied as a basal application across all treatments.

**3. RESULTS AND DISCUSSION****3.1 Effect of Sowing Methods and Nitrogen Levels on Yield Attributes of Maize****3.1.1 Cob length and cob girth**

The data on cob length (cm) have been presented in Table 1. The maximum length of a cob was obtained in the ridge sowing method (S1) with the application of 25% N through granular urea + 75% N through Nano-urea (N5). The minimum length of a cob (15.46 cm) was reported in the broadcasting method of sowing. Among the nitrogen levels the minimum cob length was reported in the N2 treatment (100% N through Nano-urea). Results showed that sole application of granular urea (46 % N) performed better than nano-urea. Similar results were also reported by Reddy et al. [10].

The data on cob girth (cm) have been presented in Table 1. The minimum girth of a cob (13.27 cm) was reported in the broadcasting method of sowing. Among the nitrogen levels the minimum cob girth of 11.65 cm was reported in the N2 treatment (100% N through Nano-urea). The maximum cob girth was obtained in the ridge

sowing method (S1) with the application of 25% N through granular urea + 75% N through Nano-urea (N5). Results showed that sole application of granular urea (46 % N) performed better than nano-urea. Similar results were also reported by Samui et al., [11], Naik et al., [12].

**3.2 Number of Grains Per Row and Number of Grains Per Cob**

The data pertaining to number of grains per row differed significantly and is given under Table 1. Among sowing methods, the maximum (35) and minimum (28) number of grains per row was obtained in the ridge method of sowing (S1) and broadcasting method, respectively. Among nitrogen levels the maximum number of grains per row (40.76) was obtained in the N5 treatment with the application of 25% N through granular urea + 75% N through Nano-urea.

Number of grains per cob was obtained in the ridge method of sowing with the application of 25% N through granular urea + 75% N through Nano-urea. Similar type of results was also obtained by Manwar and Mankar, [13], Thakur et al. [14].

**3.3 Grain weight Per Cob and Seed Index**

The data pertaining to grain weight per cob and seed index is given in Table 1. The highest grain weight per cob (249.37 g) and Seed Index (372.80 g) was reported in Ridge sowing (S1). Among the nitrogen levels, the treatment N5 was found to be superior among all the treatments. Seed index (weight of 100 seeds) was found to be maximum in ridge sowing method and the application of 25% N through granular urea + 75% N through Nano-urea. However, sole application of granular urea was comparatively higher than nano-urea. Similar types of results were also reported by Naik et al. [12]; Manwar and Mankar [13].

Table 1. Cob length, cob girth, grain weight per cob, number of grains per row, seed index and number of grains per cob

Treatments	Cob Length (cm)	Cob girth (cm)	Grain weight cob <sup>-1</sup> (g)	Number of grain row <sup>-1</sup>	Seed Index (g)	Number of grains cob <sup>-1</sup>
<b>Method of sowing</b>						
S1	18.52	15.75	249.37	35.00	302.40	372.80
S2	17.58	14.43	244.30	31.13	296.80	364.86
S3	15.46	13.27	233.33	28.00	293.26	359.73
<b>SEm±</b>	0.25	0.015	0.55	0.17	1.58	1.53
<b>CD (P=0.5)</b>	1.02	0.67	2.22	0.68	6.42	6.17
<b>Nitrogen levels</b>						
N1	15.86	12.95	183.20	26.77	289.44	350.55
N2	14.50	11.65	230.77	28.76	292.00	347.00
N3	17.61	14.11	255.08	30.11	294.00	370.22
N4	18.45	15.43	259.06	31.88	303.87	375.65
N5	19.51	16.00	283.54	40.76	308.11	385.55
<b>SEm±</b>	0.23	0.24	1.01	0.49	1.46	2.01
<b>CD (P=0.5)</b>	0.69	0.67	2.96	1.45	4.28	5.92

**Table 2. Grain yield, biological yield and Harvest Index**

Treatments	Grain yield (q/ha)	Biological Yield (q/ha)	Harvest Index (%)
<b>Method of sowing</b>			
S1	51.75	144.89	42.7
S2	48.16	134.85	41.78
S3	44.85	125.58	41.61
<b>SEm±</b>	0.48	1.35	0.51
<b>CD (P=0.5)</b>	1.94	5.47	NS
<b>Nitrogen levels</b>			
N1	44	123.2	41.72
N2	40.14	112.4	41.69
N3	47.54	133.12	41.85
N4	51.81	145.07	42.03
N5	57.76	161.75	43.27
<b>SEm±</b>	1.58	4.43	0.71
<b>CD (P=0.5)</b>	4.65	13.01	NS

### 3.4 Effect of Sowing Methods and Nitrogen Levels on Yield of Maize

#### 3.4.1 Grain yield

The data pertaining to grain yield of maize is given in Table 1. The highest grain yield was reported in Ridge sowing (S1) of 51.75 q ha<sup>-1</sup>. Among the nitrogen levels, the treatment N5 was found to be superior among all the treatments attaining the grain yield of 57.76 q ha<sup>-1</sup>. However, grain yield was comparatively higher with the sole application of granular urea than to nano-urea. Similar types of results were also reported by Naik et al. [12]; Manwar and Mankar [13].

#### 3.4.2 Biological yield and harvest index

The data for biological yield is given under Table 2. It is reported that, the sole application of granular urea was found to be better than nano urea. Biological yield was significantly higher in ridge method of sowing along with the application of 25% N through granular urea + 75% N through Nano-urea.

Harvest Index was found to be non-significant, which is expressed in terms of percentage as given under Table 2. Highest harvest index was found in the S1 and N5 treatment. Similar types of results were also obtained by Naik et al. [12]; Manwar and Mankar [13]; Thakur et al. [14].

## 4. CONCLUSION

Data from research revealed that sole application of urea was found to be better than nano-urea.

While application of urea along with nano urea was found be most effective in maize under plains zone of Uttar Pradesh region. Application of Nano-urea alone did not give good results. Hence, it is advised to grow maize under ridge sowing planting method along with the application of 25% N through granular urea + 75% N through Nano-urea.

### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have not been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Sachan DS, Khan N, Sachan R, Singh S, Gangwar P, Singh B, Kumar M. Effect of Chemical Herbicides on Diversified Weed Flora and Weed Control Efficiency in Maize (*Zea mays* L.). International Journal of Plant & Soil Science. 2023;35(17):54-61.
2. Sachan DS, Khan N, Maurya CL, Singh B. Influence of different herbicides on the growth, growth attributes and yield of maize (*Zea mays* L.) under central plains zone of Uttar Pradesh. Journal of

- Experimental Agriculture International. 2024;46(3):9-19.
3. Chiroma AM, Alhassan AB, Khan B. Yield and water use efficiency of millet as affected by land configuration treatments. J. Sustainable Agric. 2008;32(2):321 – 333.
  4. Parihar CM, Rana KS, Parihar MD. Crop productivity, quality and nutrient uptake of maize (*Zea mays* L.) cropping system as influenced by configuration and direct and residual effect of nutrient management. Indian J. agric. Sci. 2009;79(11):927-930.
  5. Amanullah Yasir M, Khan AZ, Tariq Jan M. Phenology, growth, and grain yield of maize as influenced by foliar applied urea at different growth stages. J. Plant Nutr. 2016;33:71–79.
  6. Guo C, Li P, Lu J, Ren T, Cong R, Li X. Application of controlled release urea in rice: reducing environmental risk while increasing grain yield and improving nitrogen use efficiency. Commun. Soil Sci. Plant Anal. 2016;47:1176–1183.
  7. Subbiah BV, Ashija GL. A rapid procedure for the determination of available nitrogen in soils. Current Sciences. 1956;25:259-260
  8. Olsen SR, Cole VC, Watanable FS, Dean LA. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. Cir. U.S. Dept. Agric. 1954;939.
  9. Jackson ML. Soil Chemical Analysis, Prentice Hall of India Pvt. Ltd., New Delhi;1973.
  10. Reddy BM, Elankavi S, Midde SK, Mattepally VS, Bhumireddy DV. Effects of conventional and nano fertilizers on growth and yield of maize (*Zea mays* L.). Bhartiya Krishi Anusandhan Patrika. 2022;37(4): 379-382.
  11. Samui S, Sagar L, Sankar T, Manohar A, Adhikary R, Maitra S, Praharaj S. Growth and productivity of rabi maize as influenced by foliar application of urea and nano-urea. Crop Research. 2022; 57(3):136-140.
  12. Naik MR, Hemalatha S, Reddy APK, Madhuri KN, Umamahesh V, Rakesh S. Efficient Need Based Nitrogen Management in Rabi Maize (*Zea mays* L.) using Leaf Colour Chart Under Varied Planting Density. International Journal of Bio-resource and Stress Management. 2022;13(Jun, 6):586-594.
  13. Manwar B, Pand Mankar DD. Effect of land configuration and fertilizer management in kharif maize (*Zea mays* L.): Journal of Soils and Crops. 2015; 25(1):220-225.
  14. Thakur NS, Kushwaha BB, Sinha NK. Productivity and water use in kharif sorghum (*Sorghum bicolor*) under different land configurations and mulching. Indian Journal of Agronomy. 2011;56(1):47-51.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/122723>