



# **A Decade-long Meteorological Report at the Local Scale and a Literature Review of Possible Consequences on Crops**

**A.S Arunima<sup>a++\*</sup> and D.Jacob<sup>b#†</sup>**

<sup>a</sup> Kerala Agricultural University, India.

<sup>b</sup> On Farm Research Centre, Kerala Agricultural University, India.

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/arja/2024/v17i4591>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111964>

**Original Research Article**

**Received: 10/11/2023**

**Accepted: 14/01/2024**

**Published: 14/11/2024**

## **ABSTRACT**

The purpose of the study was to look into how the physiology of crops are affected by climate change. The meteorological data for the most recent 10 years (2012–22) was given by the Department of Agricultural Meteorology, College of Agriculture Vellayani. These data were examined in order to comprehend weather patterns and investigated how the climate change, especially the temperature raise may affect plant growth and development. The results has been shown that the physiology of crops were severely affected by the temperature stress brought by climate change

<sup>++</sup> MSc Plant Physiology;

<sup>#</sup> PhD in Agronomy;

<sup>†</sup> Assistant professor and Head;

\*Corresponding author: E-mail: [arunimaas97@gmail.com](mailto:arunimaas97@gmail.com);

*Keywords: Climate change; crop physiology; temperature rise.*

## 1. INTRODUCTION

Climate change is one of the most significant challenges that plant scientists are currently facing. It is the primary cause of the biotic and abiotic pressures that have negative impact on agriculture; hence, agriculture and climate change are intrinsically linked in a number of ways. It results in variations in yearly rainfall, average temperature, heat waves, changes in weeds, pests and bacteria, worldwide changes in atmospheric CO<sub>2</sub> or ozone levels and variations in sea level etc. The agricultural production system also affected both directly and indirectly. Direct effects include the consequences on certain agricultural production systems brought on by a change in physical attributes, such as temperature levels and rainfall distribution. When changes are made to other species, such as pollinators, pests, disease vectors, and invading species, it has an indirect effect on production. Increasing greenhouse gases concentrations are the cause of extreme climatic shifts. Carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), and ozone are examples of greenhouse gases (GHG) that contribute to the greenhouse effect. The output of greenhouse gases are anticipated to rise in the near future also.

The atmospheric CO<sub>2</sub> concentrations have increased from 280ppm preindustrial reference levels to current global levels, which is above 400 ppm. According to studies, higher carbon dioxide levels promote photosynthesis especially in C3 plants. It is known as the "CO<sub>2</sub> fertilization effect". But the positive direct impact of elevated [CO<sub>2</sub>] may be countered by other climatic change effects like increased temperatures, higher tropospheric ozone concentrations, and altered precipitation patterns. The consequences of elevated atmospheric CO<sub>2</sub> and associated temperature and precipitation variations were anticipated to have an impact on the distribution, relationships, and eco physiology of plants (Intergovernmental Panel on Climate Change (IPCC), [1]). The longer eCO<sub>2</sub> treatments may result in photosynthetic adaptation because of an increase in soluble sugars, which leads to an imbalanced C: N ratio, faster leaf senescence, and/or limited growth rate, which affect crop yield and quality. It is believed that crop quality is a multifaceted and it is a complex topic that involves growth, assimilation, partitioning, storage, pre- and post-harvest, as well as nutritional, technological and environmental

components [2]. In a future high-CO<sub>2</sub> environment, the elemental (e.g., zinc, iodine) and macromolecular (e.g., protein) composition of plant tissues is anticipated to alter [3]. Crop physiologists will need to address the interests of breeders and processors more in this circumstance by analysing, characterising, and modelling changes in food quality among crop varieties and species under climate change scenarios [2]. Hence, in this study, we examined the impact of climate change, specifically temperature rises over the previous ten years, using meteorological data from the Department of Agricultural Meteorology, College of Agriculture, Vellayani on physiology of crops.

## 2. MATERIALS AND METHODS

The Department of Agricultural Meteorology, College of Agriculture Vellayani, Thiruvananthapuram maintains a class B agromet observatory where all the meteorological equipments are housed and the observer regularly takes observations twice a day at 7:22 a.m. and 2:22 p.m., respectively. This weather information is routinely sent to the IMD headquarters in New Delhi. In this study, we examined both average maximum and average minimum temperatures over the past ten years (2013-22) from the data recorded in the department (Tables 1 & 2).

## 3. RESULTS AND DISCUSSION

The data shows that over the past ten years, the maximum temperature range has been between 30°C and 35.3°C and the minimum temperature range has been between 21°C and 26.3°C. The highest maximum temperature and the greatest minimum temperature recorded in 2013 were 33.1°C and 25.4°C, respectively. All of 2013's months had a dramatic fluctuation in maximum temperatures, which ranged from 29 °C (June) to 33.1 °C. (April). Additionally, there were variations in the minimum temperature, which ranged from 21.9°C in December to 25.4°C in April. The months with the highest maximum temperature in 2014 were observed to be both march and April (32.4°C). The minimum temperature range was 21.5°C (January) to 25.2°C (June). The maximum and minimum temperatures of 2014 were somewhat lower than they were in 2013. In 2015, the maximum temperature of every month's were exceeded 30°C and most of the months it was recorded

**Table 1. Weather data- The monthly average temperature (Maximum and Minimum temperature) at college of agriculture, Vellayani from 2013-2017. The data collected from department of agricultural meteorology, college of agriculture, vellayani**

	2013		2014		2015		2016		2017	
	Max Temp (°C)	Min Temp (°C)	Max Temp (°C)	Min Temp (°C)	Max Temp (°C)	Min Temp (°C)	Max Temp (°C)	Min Temp (°C)	Max Temp (°C)	Min Temp (°C)
<b>January</b>	30.3	22.0	30.6	21.5	30.6	21.6	32.3	22.6	32.6	22.8
<b>February</b>	31.4	22.1	31.3	22.3	31.6	22.3	32.9	23.2	32.5	21.5
<b>March</b>	32.3	24.0	32.4	22.9	32.4	23.7	34.5	24.7	33.4	24.5
<b>April</b>	33.1	25.4	32.4	24.5	32.7	24.5	35.3	26.4	34.2	25.9
<b>May</b>	32.1	25.0	31.9	24.8	32.1	25.3	33.8	24.8	33.7	25.5
<b>June</b>	29.0	23.0	30.8	25.2	31.4	24.5	31.5	24.4	31.4	25.5
<b>July</b>	29.1	22.7	30.0	24.3	31.3	24.6	31.6	24.7	31.6	24.7
<b>August</b>	29.2	23.7	29.5	23.7	31.7	24.6	31.8	24.8	31.2	24.6
<b>September</b>	29.2	23.9	30.2	24.2	31.4	24.4	31.8	24.5	31.5	24.5
<b>October</b>	30.8	23.2	30.5	23.8	31.3	24.0	31.7	24.3	31.1	24.9
<b>November</b>	30.6	23.5	30.2	23.4	31.6	23.8	32.0	24.0	30.8	24.2
<b>December</b>	30.8	21.9	30.2	24.0	31.5	23.3	32.3	23.6	31.8	23.4

\*Max Temp (°C) – Maximum temperature in °C

\*Min Temp (°C) – Minimum temperature in °C

**Table 2. Weather data-The monthly average temperature (Maximum and Minimum temperature) at College of Agriculture, Vellayani from 2018-2022.The data collected from Department of Agricultural Meteorology, College of Agriculture, Vellayani**

	2018		2019		2020		2021		2022	
	Max Temp (°C)	MinTemp (°C)	Max Temp (°C)	Min Temp (°C)	Max Temp (°C)	Min Temp (°C)	Max Temp (°C)	Min Temp (°C)	Max Temp (°C)	Min Temp (°C)
<b>January</b>	31.7	21.7	32.0	21.1	32.3	22.9	31.8	23.5	32.4	22.9
<b>February</b>	32.4	23.6	33.6	23.6	33.0	23.3	33.2	22.3	32.4	21.5
<b>March</b>	33.3	24.7	34.6	24.8	33.5	24.6	34	23.3	33.4	24.4
<b>April</b>	33.9	25.9	35.0	25.8	34.2	25.6	33.8	25.9	33.1	22.5
<b>May</b>	32.8	25.3	34.1	26.3	32.6	25.6	32.1	24.9	32.2	23.5
<b>June</b>	31.1	24.7	31.9	25.4	31.5	24.8	32.3	25.6	31.6	23.4
<b>July</b>	30.7	23.8	30.8	24.8	31.7	24.7	31.3	24.9	30.4	22.7
<b>August</b>	30.3	23.6	30.8	24.2	31.7	26.3	30.9	23.9	30.5	22.5
<b>September</b>	32.4	24.3	30.9	24.4	31.1	24.6	31.1	23.5	30.9	23.2
<b>October</b>	31.4	24.3	30.7	24.1	31.3	24.8	31.1	22.3	31.2	23.6
<b>November</b>	31.6	24	31.9	24.5	32.4	25.1	30.1	22.5	30.9	22.5
<b>December</b>	32.0	23.5	31.7	23.8	32.5	24.1	32.1	23.4	31.2	21.1

\*Max Temp (°C) – Maximum temperature in °C

\*Min Temp (°C) – Minimum temperature in °C

above 31°C. The highest maximum temperature was recorded in the month of April (32.7°C) and highest minimum temperature was recorded in May (25.3°C). It has been noted that the maximum temperature has never dropped below 29°C since 2015. Most of the months in 2015 have crossed the minimum temperature of 24°C. The year 2016 was noted as having the highest maximum and minimum temperatures over the previous 10 years and it was in the month of April with 35.3°C and 26.4°C respectively. So it was recorded as one of the hottest year among 2013-2022. Because, the maximum temperature from January to December in 2016 were 32.3°C, 32.9°C, 34.5°C, 35.3°C, 33.8°C, 31.5°C, 31.6°C, 31.8°C, 31.8°C, 31.7°C, 32°C, 32.3°C and the minimum temperature were 22.6°C, 23.2°C, 24.7°C, 26.4°C, 24.8°C, 24.4°C, 24.7°C, 24.8°C, 24.5°C, 24.3°C, 24°C, 23.6°C respectively. In 2017, the highest maximum temperature and minimum temperature was recorded in the month of April (34.2°C and 25.9°C). Most of the months, the maximum temperatures were crossed 31°C and minimum temperatures were crossed 24 °C).

In 2018, the highest maximum temperature was recorded in the month of April (33.9 °C), and the lowest maximum temperature was recorded in August (30.3°C). The highest minimum temperature was observed in April (25.9 °C), and the lowest minimum temperature was observed in January (21.7°C). The year 2019 was recorded as the second hottest year among the past ten years, with a highest average maximum temperature of 35°C in April. The following maximum temperatures were experienced in 2019 from January to December: 32°C, 33.6°C, 34.6°C, 35°C, 34.1°C, 31.9°C, 30.8°C, 30.9°C, 30.7°C, 31.9°C, and 31.7°C. The highest maximum temperature was recorded in the month of April (35°C) and the lowest maximum temperature was recorded in the month of October (30.7°C). In 2019, the minimum temperatures were 21.1°C, 23.6°C, 24.8°C, 25.8°C, 26.3°C, 25.4°C, .8°C, 24.2°C, 24.4°C, 24.1°C, 24.5°C, 23.8°C respectively with a highest in May (26.3°C) and the lowest in January (21.1°C). In 2020, the maximum highest temperature was recorded in April (34.2°C) and the highest minimum temperature was recorded in August (26.3°C). The majority of the months in 2020, recorded a maximum temperature above 31.5°C, with the exception of September (31.1°C) and October (31.3°C) and minimum temperature were above 23.3 °C, except January (22.9°C). The month with the greatest maximum

(33.8°C) and minimum (25.9°C) temperatures in 2021 was April. The lowest maximum temperatures (22.3°C) were recorded in the months of February and October. A temperature range of 30.4°C to 33.4°C was reported for each month of the year 2022. The highest maximum temperature (33.4°C) and highest minimum temperature (24.4°C) was observed in March. The maximum temperature have crossed 33°C in both March (33.4°C) and April (33.1°C) of 2022. Except for February (21.5°C) and December (21.1°C), most of the months have a minimum temperature exceeding 22.5°C in 2022. These are the conclusions drawn from the data gathered from the Department of Agricultural Meteorology, College of Agriculture, Vellayani. According to the facts above, temperatures have risen during the last 10 years.

### 3.1 Impact of Elevated CO<sub>2</sub> on Elevation of Temperature and Physiology of Crops

The impact of climate change is a long-term concern, but it demands immediate action because greenhouse gases are accumulating in the atmosphere and there is the possibility of global temperatures rising by more than 2 degrees Celsius. Long-term global mean temperature estimates indicate an extra 1.1-4.8 °C rise over the previous 50 years, depending on the scale of future greenhouse gas emissions [4]. These changes in global temperature would cause further changes in the climate, resulting in an increased frequency of heat waves, fewer days of freezing temperatures, less rainfall but more intense precipitation, and a higher incidence of droughts and other weather extremes experienced around the world, all of which would have a negative impact on agricultural production (Dempewolf *et al.*, 2014).

CO<sub>2</sub> is one of the greenhouse gases, which have significant direct effects on plant development, physiology, and chemistry [5]. It helps to stimulate photosynthesis and water use efficiency at higher concentration [6]. Plants absorb atmospheric CO<sub>2</sub> and it represents not only a gain of stored chemical energy for the plant, but also supplies the carbon skeletons for the organic molecules that make up a plant's structure. Around 96% of a normal plant's dry mass is made up of carbon, hydrogen, and oxygen absorbed into organic molecules during photosynthesis [7]. Hence, photosynthesis is at the core of plants' nutritional metabolism, and

increasing the amount of CO<sub>2</sub> available for photosynthesis can have a significant impact on plant growth and many other aspects of physiology. According to Ainsworth & Rogers 2007, The development of plants under increased CO<sub>2</sub> concentrations of 475–600 ppm results in an average 40% increase in leaf photosynthetic rates. The opening of stomata, through which plants exchange gases with the environment, are regulated by carbon dioxide concentrations. The stomatal pores in plants close when CO<sub>2</sub> concentrations are higher than ambient, and they open when CO<sub>2</sub> concentrations are lower. Water may flow out of leaves through open stomata, which also allow CO<sub>2</sub> to permeate into leaves for photosynthesis. So plants can maintain a high photosynthetic rates with low stomatal conductance at higher CO<sub>2</sub> concentrations. Since photosynthesis and stomatal behavior are central to plant metabolism, growing plants under elevated CO<sub>2</sub> has a variety of secondary effects on plant physiology. The availability of additional photosynthesis allows most plants to grow faster and produce dry matter under increased CO<sub>2</sub> levels.

The long-term climatic change brought on by the steady increase in CO<sub>2</sub> concentration will balance off the increase in agricultural yield [8]. Increased CO<sub>2</sub> also leads to changes in the chemical composition of plant tissues. Due to increased photosynthetic activity, leaf non structural carbohydrates (sugars and starches) per unit leaf area increased by an average of 30–40% under FACE elevated CO<sub>2</sub> condition (Long, 2005).The nitrogen content of leaves in plant tissue often declines as CO<sub>2</sub> levels rise and it is reported an average of 13% less nitrogen per unit leaf mass [9]. This decrease in tissue nitrogen may be due to several factors such as nitrogen dilution due to increased carbohydrate concentration, reduced mineral uptake from the soil because of reduced stomatal conductivity and plants absorb less water [10] and reduced rate of assimilation of nitrate into organic compounds [11]. Plant tissue protein concentration is closely related to plant nitrogen status. In FACE experiments, protein concentrations in wheat, rice, barley grains, and potato tubers decreased by 5–14% with increasing CO<sub>2</sub> [3,10]. Plant concentrations of nutritionally important minerals such as calcium, magnesium and phosphorus may also decreased with elevated CO<sub>2</sub> levels [12]. So the rising CO<sub>2</sub> emissions, high

temperature and associated climate change over the next century are likely to impact both agricultural production and food quality.

### 3.2 Temperature Rise and Crop Physiology

Plants experience several kinds of stresses during growth and development. Temperature changes can affect agricultural productivity. In order to ensure food security across the globe, we must be aware of climate change, particularly high temperature. In recent years, scientists have observed a rapid rise in average temperatures across the country. According to the National Resources Defense Council's (NRDC) "Killer Summer Heat" report, the average temperature in the United States could rise by as much as 11 degrees Fahrenheit by the end of the 21<sup>st</sup> century [13]. Temperature is one of the key meteorological factors in crop growth and development. It impacts the enzyme functions in a leaf and causes changes in the developmental stages that are directly related to crop yield [14]. The optimum temperature for growing crops is about 15-30°C. However temperature elevations predominantly affect plant's physiological processes particularly photosynthesis, respiration, transpiration, and yield. So any changes in long-term mean annual temperature and extreme temperature events will probably have a big effect on physiological responses of crops and ultimately affect how much food and fuel crops produce around the world.

The physiological damages caused by high temperatures such as leaf abscission, leaf burning, senescence and restricted root and shoot development have a significant impact on crops and ultimately reduce production. Also it can leads to lower seed germination rates, plant emergence, weak seedling vigor, aberrant seedlings, and reduced radicle and plumule development [15]. Many food crops such as rice, wheat, soy, corn, cotton, sorghum and tomato are very sensitive to high temperatures. High temperatures may disrupt a number of metabolic processes that occur in guard cells; the stomatal response is typically regulated by transpiration rate, photosynthetic rate, plant water status, and vapour pressure deficit [16]. High temperatures increase the rate of evapotranspiration throughout the vegetative and reproductive stages, which restricts the amount of water that plants can take in. This effects dehydration, which inhibits growth of both the individual

organs and the entire plant [17]. High temperature stress can affect crop's seedling stage in mung bean, and wheat as well as the reproductive stage in wheat and other cereals [18].

The reproductive stage is the most sensitive time in a crop's life cycle, and this sensitivity causes a considerable reduction in seed set and crop output [19]. It is reported that the seedling and reproductive phases of rice can be negatively impacted by temperatures above 35°C. It has been shown to have a negative impact on various aspects of reproductive development in a variety of species including meiosis in both male and female gametes, pollen germination and pollen tube growth, pollen/pistil interactions, ovule viability, pollen grain number, formation of endosperm and embryo development, fertilization and post-fertilization processes. According to van der Merwe et al., [20], the sunflower seeds and the oil constituents are significantly impacted by high temperature during grain filling stage. It is also observed that the high temperature (33°C) decreases anther dehiscence and pollen fertility rate; it results in a reduction in the amount of pollen on the stigma, which in turn causes decreased fertilization, spikelet fertility, and sterile seed in rice [21]. Pollen viability is lost as a result of high-temperature stress due to changes in membrane integrity, protein, carbohydrate and lipid mechanisms and phospholipid profiles [22].

High temperatures also have an influence on photosynthetic membranes, grana stack growth, ion leakage and aberrant stacking. High temperature affects the activities of carbon metabolic enzymes, starch accumulation, and sucrose synthesis through down regulating certain genes involved in carbohydrate metabolism. Abscisic acid, ethylene, and salicylic

acid are the key phytohormones that are increased by high temperatures, whereas gibberellic acid, cytokinin and auxin are reduced. Reactive oxygen species are also produced as a result of high temperatures [15] results oxidative stress [23]. Increased generation of ROS, including lipid peroxidation, superoxide free radicals, and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), results in increased cell membrane damage [24]. High temperature reduces the net assimilation rate and the denaturation of proteins occur, it induces programmed cell death in some cells or tissues. This results in decreased ion flux, leakage of electrolytes, changes in relative water content, the production of toxic compounds, and disruption of homeostasis, all of which reduce cell viability. Due to lower Rubisco activation, many plant species experience a drop in stomatal conductance (gs) and net photosynthetic rate when exposed to moderate heat stress.

Extreme temperatures have a significant impact on photosynthesis in leaves. Temperature has an impact on the mechanisms involved in photosynthetic metabolism [25]. High temperatures have a significant impact on chloroplast throughout the photosynthesis process, including grana stack enlargement and aberrant stacking [26]. The influence of high temperature stress on chlorophyll production in plastids plays a significant role in light harvesting. The chlorophyll pigments in plastids are also harmed and degraded by the high-temperature stress. By eliminating the enzymes involved in the process of chlorophyll production, high temperature stress-related inhibition of chlorophyll biosynthesis was achieved. Under high-temperature stress, 5-aminolevulinic acid dehydratase (ALAD), the initial enzyme of pyrrole biosynthesis, showed reduced enzymatic activity [27].





Fig. 1. The graph depicting the month-by-month average maximum temperature from 2013-22



**Fig. 2. The graph depicting the month-by-month average minimum temperature from 2013-22**

So the high-temperature stress is one of the main abiotic variables that negatively impacts crop output. Through its impact on the physiological system, high-temperature stress poses a danger at various development phases and eventually lowers output [28-31].

#### 4. CONCLUSION

Climate change will cause temperatures to rise in the next years, which will have a significant impact on agricultural productivity. Despite the fact that many research efforts focus on biotic and abiotic stresses to shield plants from harmful circumstances. The plants need to be given the best circumstances for growth at each stage. Germination, vegetative stage, reproductive stage, and the yield are all impacted by high temperature stress and it is critically necessary to find innovative ways to adapt crops to these changes. ie, From the seedling stage up to the harvest stage, the crop has to be improved.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### CONSENT

It is not applicable.

#### ETHICAL APPROVAL

This article does not contain any studies with human participants or animals performed by any of the authors.

#### ACKNOWLEDGEMENT

Thanks to the faculties in The Department of Agricultural Meteorology, College of Agriculture, Vellayani, Kerala Agricultural University,

Thrissur, Kerala for extending their guidance and technical assistance for this study.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. IPCC 2014. Climate Change 2014: Synthesis report. Contribution of working groups I, II and III to the fifth assessment report of the intergovernmental panel on climate change.
2. Hay R, Porter J. The physiology of crop yield (2nd ed.). Oxford: Blackwell. 2006;43-49.
3. Taub D, Miller B, Allen H. Effects of elevated CO<sub>2</sub> on the protein concentration of food crops: A meta-analysis. *Global Change Biology*. 2008;14: 565–575.
4. Walsh J, Wuebbles D, Hayhoe K, Kossin J, Kunkel K, Stephens G, Somerville R. Our changing climate. U.S. Global Change Research Program; 2014.
5. Ziska LH, Kleinman DL, Cloud-Hansen KA. Rising atmospheric carbon dioxide and plant biology: the overlooked paradigm. In controversies in science and technology, from climate to chromosomes. New Rochele: Liebert. 2008;379-400.



6. Long SP, Ainsworth EA, Rogers A, Ort DR. Rising atmospheric carbon dioxide: plants FACE the future. *Annu. Rev. Plant Biol.* 2004;55:591–628.
7. Marschner H. Mineral nutrition of higher plants, 2nd ed. London, UK: Academic Press; 1995.
8. Wei PS, Chiu HH, Hsieh YC, Yen DL, Lee C, Tsai YC. Absorption coefficient of water vapor across atmospheric troposphere layer. *Heliyon.* 2019;5:e01145. DOI: 10.1016/j.heliyon.2019.e01145
9. Ainsworth EA, Long SP. What have we learned from 15 years of free-air CO<sub>2</sub> enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO<sub>2</sub>. *NewPhytologist.* 2005;165:351-372.
10. Taub DR, Wang XZ. Why are nitrogen concentrations in plant tissues lower under elevated CO<sub>2</sub>? A critical examination of the hypotheses. *Journal of Integrative Plant Biology.* 2008;50:1365-1374.
11. Bloom AJ, Burger M. Carbon dioxide inhibits nitrate assimilation in wheat and arabidopsis. *Science.* 2010;328: 899-903.
12. Loladzel. Rising atmospheric CO<sub>2</sub> and human nutrition: Toward globally imbalanced plant stoichiometry? *Trends in Ecology & Evolution.* 2002;17:457-461.
13. NRDC. The road from paris: The United States Progress toward its climate pledge; 2016
14. Zhu P, Jin Z, Zhuang Q, Ciais P, Bernacchi C, Wang X, Makowski D, Lobell D. The important but weakening maize yield benefit of grain filling prolongation in the US Midwest. *Global Change Biology.* 2018;24:4718–4730.
15. Bitá CE, Gerats T. Plant tolerance to high temperature in a changing environment: Scientific fundamentals and production of heat stress-tolerant crops. *Frontiers in Plant Science.* 2013;4:1-18. DOI: 10.3389/FPLS.2013.00273
16. Urban J, Ingwers MW, McGuire MA, Teskey RO. Increase in leaf temperature opens stomata and decouples net photosynthesis from stomatal conductance in *Pinus taeda* and *Populus deltoides* x *nigra*. *Journal of Experimental Botany.* 2017;68(7):1757-1767. DOI: 10.1093/JXB/ERX052
17. Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A. Crop production under drought and heat stress: Plant responses and management options. *Frontiers in Plant Science.* 2017;8:1-16. DOI: 10.3389/fpls.2017.01147
18. Akter N, Rafiqul Islam M. Heat stress effects and management in wheat. A review. In: *Agronomy for Sustainable Development.* France: Springer-Verlag. 2017;375. DOI: 10.1007/s13593-017-0443-9
19. Hein NT, Ciampitti IA, Jagadish SVK. Bottlenecks and opportunities in field-based high-throughput phenotyping for heat and drought stress. *Journal of Experimental Botany.* 2021;72(14):5102-5116. DOI: 10.1093/JXB/ERAB021
20. Van der Merwe R, Labuschagne MT, Herselman L, Hugo A. Effect of heat stress on seed yield components and oil composition in high- and mid-oleic sunflower hybrids. *South African Journal of Plant and Soil.* 2015;32(3):121-128. DOI: 10.1080/02571862.2015.1018354
21. Hurkman WJ, Vensel WH, Tanaka CK, Whitehand L, Altenbach SB. Effect of high temperature on albumin and globulin accumulation in the endosperm proteome of the developing wheat grain. *Journal of Cereal Science.* 2009;49(1):12-23. DOI: 10.1016/j.jcs.2008.06.014
22. Djanaguiraman M, Perumal R, Jagadish SVK, Ciampitti IA, Welti R, Prasad PVV. Sensitivity of sorghum pollen and pistil to high-temperature stress. *Plant, Cell & Environment.* 2018;41(5):1065-1082. DOI: 10.1111/PCE.13089
23. Potters G, Pasternak TP, Guisez Y, Palme KJ, Jansen MAK. Stress-induced morphogenic responses: Growing out of trouble? *Trends in Plant Science.* 2007;12(3):98-105. DOI: 10.1016/J.TPLANTS.2007.01.004
24. Narayanan S, Tamura PJ, Roth MR, Prasad PVV, Welti R. Wheat leaf lipids during heat stress: I. High day and night temperatures result in major lipid alterations. *Plant, Cell & Environment.* 2016;39(4):787-803. DOI: 10.1111/PCE.12649
25. Asthir B. Protective mechanisms of heat tolerance in crop plants. *Journal of Plant Interactions.* 2015;10(1):202-210. DOI: 10.1080/17429145.2015.1067726
26. Wahid A, Shabbir A. Induction of heat stress tolerance in barley seedlings by pre-

- sowing seed treatment with glycinebetaine. *Plant Growth Regulation*. 2005;46(2):133-141.  
DOI: 10.1007/s10725-005-8379-5
27. Ashraf M, Harris PJC. Photosynthesis under stressful environments: An overview. *Photosynthetica*. 2013;51:163-190.  
DOI: 10.1007/S11099-013-0021-6
28. Wahid A, Gelani S, Ashraf M, Foolad M. Heat tolerance in plants: An overview. *Environmental and Experimental Botany*. 2007;61(3):199-223.  
DOI: 10.1016/j.envexpbot
29. Ainsworth EA, Rogers A. The response of photosynthesis and stomatal conductance to rising (CO<sub>2</sub>): Mechanisms and environmental interactions. *Plant, Cell and Environment*. 2007;30:258-270.
30. IPCC. Summary for Policymakers. In: *Global warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*; 2018.
31. Liu W, Yin T, Zhao Y, Wang X, Wang K, Shen Y. Effects of high temperature on rice grain development and quality formation based on proteomics comparative analysis under field warming. *Frontiers in Plant Science*. 2021; 12:2320.  
DOI: 10.3389/FPLS.2021.746180/BIBTEX

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/111964>