



Microalbuminuria in Women with Risk Factors for Gestational Diabetes Mellitus in Some Selected Hospitals in Sokoto, Nigeria

M. K. Dallatu^{1*}, A. M. Kaoje², A. U. Adoke³, J. A. Kehinde¹ and J. M. Bunza¹

¹Department of Chemical Pathology, Faculty of Medical Laboratory Sciences, Usmanu Danfodiyo University, Sokoto, Nigeria.

²Department of Chemical Pathology, Usmanu Danfodiyo University Teaching Hospital, Sokoto, Nigeria.

³Department of Medicine, Usmanu Danfodiyo University Teaching Hospital, Sokoto, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Authors MKD and AMK designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors JAK and JMB managed the analyses of the study. Authors JAK and AUA managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRIMPS/2019/v6i230097

Editor(s):

(1) Dr. Aurora Martinez Romero, Professor, Department of Clinical Biochemistry, Juarez University, Durango, Mexico.

Reviewers:

(1) Javier Rodríguez Villanueva University of Alcalá, Alcalá de Henares, Madrid, Spain.

(2) Saime Paydas, Internal Medicine Nephrology Cukurova University, Turkey.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/47454>

Original Research Article

Received 19 October 2018

Accepted 12 February 2019

Published 06 March 2019

ABSTRACT

Gestational diabetes mellitus (GDM) is a common metabolic abnormality which affects approximately 2-5% of pregnancies. Risk factors such as previous infants with macrosomia, strong family history of type 2 diabetes or GDM, poor glycaemic control and high pre-pregnancy body mass index (BMI) have been implicated in the development of GDM. This study was conducted to determine the prevalence of Microalbuminuria in women with risk factors for GDM and to estimate the levels of Urinary Microalbumin in these women. Fifty (50) pregnant women with risk factors for GDM and 50 controls (pregnant women without risk factors for GDM) were evaluated for Microalbuminuria. Microalbuminuria was estimated using Turbidimetric method, Random Plasma

*Corresponding author: E-mail: muhammad.dallatu@udusok.edu.ng;

Glucose was estimated using Glucose oxidase method, serum Urea was estimated using Diacetyl Monoxime method, Creatinine was estimated using Jaffe Slot method and Albumin was estimated using Bromo Cresol Green method. The prevalence of Microalbuminuria in women with risk factors for GDM was 22%. Urinary microalbumin was significantly higher in the study subjects (56.36 ± 8.44 mg/L) than in the control (17.32 ± 4.5) mg/L. The mean \pm standard error of mean of random plasma glucose in the study subjects was (5.84 ± 0.16) mmol/L and that of the control was (4.33 ± 0.14) mmol/L. The mean \pm standard error of mean Serum Urea, Creatinine and Albumin were (4.1 ± 0.15) mmol/L, (0.70 ± 0.03) mg/dL, and (3.06 ± 0.05) g/dL respectively while that of the control was (3.47 ± 0.13) mmol/L, (0.63 ± 0.01) mg/dL and (2.78 ± 0.09) g/dL respectively. Obesity was strongly correlated to microalbuminuria.

Keywords: Gestational diabetes mellitus; pregnant women; microalbuminuria; urea; creatinine.

1. INTRODUCTION

Gestational diabetes mellitus (GDM), by definition is any degree of glucose intolerance with the onset or first recognition during pregnancy [1]. This definition applies regardless of whether the treatment involves insulin or diet modification alone; it may also apply to conditions that persist after pregnancy. Women with GDM have a 40-60% chance of developing diabetes mellitus over the period of 5-10 years after the pregnancy [2].

Approximately 4% of pregnant women in the United States have diabetes mellitus. It has been estimated that 75 to 90% of high blood glucose during pregnancy are gestational diabetes, and the remaining 10 to 25% are either type 1 or type 2 diabetes mellitus [3]. Normalizing blood glucose concentrations before and early in pregnancy can reduce these risks to levels of the general population [3]. All GDM severity levels will result in adverse neonatal outcome [4]. Global fetal and infant loss, perinatal mortality, neonatal mortality, and malformations rates are significantly greater if the mother is affected by diabetes than in the non diabetic population [5]. The predominant pathogenic factor in GDM could be inadequate insulin secretion. It has been convincingly demonstrated that GDM occurs as a result of a combination of insulin resistance and decreased insulin secretion. The cumulative incidence of Type 2 diabetes is 50% at 5 years. GDM is also a predictor, or even an early manifestation, of the metabolic (insulin resistance) syndrome. GDM is a cardiovascular risk factor and affected patients should be screened to prevent late [6].

Studies in Nigeria have shown high and low values respectively. In Ibadan, South West Nigeria the prevalence of GDM is 4.9% [7]. Consequently, in Sokoto, North-western Nigeria

the prevalence of Gestational diabetes mellitus was found to be 7.7% [8].

Risks factors for gestational diabetes mellitus includes Macrosomia (≥ 4.5 kg), Family history of type 2 diabetes mellitus, Body Mass Index (BMI) ≥ 30.0 , Pre diabetes, High blood pressure, Previous history of unexplained stillbirth and miscarriage and Hormonal disorders such as Polycystic Ovary Syndrome among others [9].

Microalbuminuria is an increased excretion of albumin above the reference range for healthy non diabetic subjects, but undetectable by the Albustix dipstick test, or as modified by expert group, as a urinary albumin excretion rate between 20 – 200 μ g/min (3-30 mg/mmol ACR) in an overnight or 24 hours sample on at least 2 of three occasions within a period 6 months [10]. Microalbuminuria is also defined as the urinary excretion of 30-300 mg of albumin per day [11,12]. The patients with longer duration of diabetes mellitus, hypertension, older age, higher body mass index (BMI) and adverse lipid profile are more prone to develop microalbuminuria [13]. Screening tests for microalbuminuria are recommended annually for patients with type 1 diabetes of greater than 5 years duration and for all patients with type 2 diabetes from the time of diagnosis.

Normal human urine contains only very small quantities of albumin, less than 30 mg of albumin being excreted by healthy adults in 24 hours. The appearance of large amounts of albumin in the urine is a cardinal sign of renal damage, especially glomerular diseases, and is not detectable by screening techniques using urinary dipsticks. Early detection of microalbuminuria allows early intervention with a goal of delaying the onset of overt diabetic nephropathy [14]. This study was therefore designed to investigate microalbuminuria in pregnant women with the

risk factors for the development of gestational diabetes.

2. MATERIALS AND METHODS

2.1 Study Area

The study was carried out in the Department of Chemical Pathology, Usmanu Danfodiyo University Teaching Hospital (UDUTH) Sokoto, Nigeria. Usmanu Danfodiyo University Teaching Hospital is a Tertiary Health Facility located in Sokoto, the Sokoto State Capital, Nigeria. It serves as a referral centre for more than 10 million people of the Nigerian States of Sokoto, Zamfara and Kebbi, and neighboring Niger and Benin Republic in the West African sub region.

Sokoto state is located at the extreme part of the North-Western Nigeria, (longitude 11°-30 to 13°-50 east and latitude 4-6° north). Sokoto state shares a border with Niger republic to the North, Kebbi state to the West, and South and Zamfara state to the south and east. Sokoto state occupies an area of short-grass savannah vegetation in the South and thorn shrub in the North generally of Arid zone. The State covers a total land area of about 32,000 square kilometers and a population of 4,953,078 million based on 2013 projection [15].

2.2 Study Subjects

The study subjects were recruited from population of Pregnant women attending the antenatal clinic (ANC) in Specialist Hospital, Sokoto and Maryam Abacha Women and Children Hospital, Sokoto. Subjects were selected by simple random sampling technique. A structured questionnaire was administered to each participant and relevant bio data and clinical informations were obtained. Informed consent for inclusion into the study was obtained from each subject using the standard protocol.

2.3 Study Design

It is a case control study. The subjects were grouped into 2:

Group 1: Fifty (50) pregnant women without risk factor for Gestational Diabetes Mellitus (control)

Group 2: Fifty (50) pregnant women with risk factors for Gestational Diabetes Mellitus

2.4 Sample Collection and Processing

A total number of one hundred blood samples and urine samples were collected. Fifty numbers

of blood and urine samples were collected from women with risk factors for Gestational Diabetes Mellitus, 50 numbers of blood and urine samples were collected from pregnant women without gestational diabetes (controls). Spot urine sample (2 mL) was collected into a universal container for the estimation of microalbuminuria while 3 ml of blood sample was collected into fluoride oxalate container for the estimation of glucose and 3 mL of blood into a plain container for the estimation of urea, creatinine and albumin respectively. The blood in the plain container was allowed to clot after which it was centrifuged, separated and the serum was stored in a freezer at -20°C till analysis. The urine sample was analyzed immediately for Microalbuminuria, glucose was also estimated immediately after separation of plasma from the whole blood.

2.5 Biochemical Analysis

Urinary microalbumin was estimated using Turbidimetric method [16]. Blood glucose was estimated by glucose oxidase method [17]. Serum Urea was estimated by Diacetyl Monoxine Method, serum albumin was estimated using Bromocresol green Method [18]. Serum Creatinine was estimated using Jaffe's Method [19].

Data obtained from this study were analyzed using statistical package for social sciences (SPSS) for Windows version 20 (SPSS Inc., Chicago, IL, USA). The data were expressed as the mean \pm standard error of mean. Paired comparison was made using student T- test, and less or equal to 0.05 ($P < .05$) was regarded as statistically significant.

3. RESULTS

The prevalence of microalbuminuria in women with risk factors for gestational diabetes mellitus was found to be 22%.

Table 1 shows the Socio- economic and Demographic characteristics of pregnant women with and without risk factors for gestational diabetes mellitus. Majority of the study population were Hausa (80%), among which most of them (44%) had no Formal Education and 62% of them are married to peasant farmers.

Table 2 shows the age distribution (in years) and gestational ages (in weeks) of pregnant women with and without risk factors for gestational diabetes mellitus. Majority of the study population

had pregnancies with gestational age between 24- 28 weeks. Majority of the study population (60%) were within the age group of 18-25 years.

Table 3 shows the mean urinary levels of microalbumin and serum levels of random plasma glucose, albumin, urea and creatinine in pregnant women with and without risk factors for gestational diabetes mellitus. The mean ± standard error of mean urinary microalbumin level in pregnant women with risk factors for gestational diabetes mellitus was significantly higher than controls p- value < 0.05. The difference between pregnant women with and without risk factors for gestational diabetes mellitus of serum concentrations of random plasma glucose, albumin, urea and creatinine were not statistically significant (p>0.05).

Table 4 shows the mean levels of BMI, blood pressure, family history of diabetes and family history of hypertension in pregnant women with and without risk factors for gestational diabetes mellitus. The mean ± standard error of mean BMI of the study population was significantly higher in pregnant women with risk factors for gestational diabetes mellitus than in controls, p- value < 0.05. The mean ± standard error of mean systolic blood pressure and diastolic blood pressure

shows no difference across the groups, p- value > 0.05.

Table 5 shows the correlation of urinary microalbumin to serum urea, creatinine, albumin and random plasma glucose.

4. DISCUSSION

In this study, the prevalence of microalbuminuria in women with risk factors for gestational diabetes mellitus was found to be 22%. This is similar to the findings of 22% prevalence of microalbuminuria in diabetic patients in Usmanu Danfodiyo University Teaching Hospital, Sokoto [20]. The prevalence of microalbuminuria in the current study is similar to the 20% prevalence found in a cohort of African- American women with GDM [21] even though, our patients just have risk factors for GDM and not GDM proper. This similarity could be due to geographical location and also the socio economic status of the population. Our finding is lower than the prevalence of 25% reported in 40 type II diabetic patients from Ilorin, Nigeria, [22]. This is also in contrast to the prevalence of 38% reported in Port Harcourt [23], 37.6% in Lagos [24]. These variations in prevalence might be attributed to factors such as differences in populations, method of urine collection or differences in ethnic susceptibility [25].

Table 1. Socio-economic characteristics of pregnant women with and without risk factors for gestational diabetes mellitus

Characteristics	Pregnant women +RFGDM (n= 50)	Pregnant women RFGDM (n= 50)
Tribes		
Hausa	43(86%)	40(80%)
Yoruba	3(6%)	4(8%)
Igbo	2(4%)	3(6%)
Others	2(4%)	3(6%)
Educational status		
Primary	20(40%)	18(36%)
Secondary	6(12%)	8(16%)
Tertiary	4(8%)	2(4%)
Informal	20(40%)	22(44%)
Occupation of husbands		
Farmer	30(60%)	31(62%)
Trader	15(30%)	14(28%)
Civil servants	3(6%)	3(6%)
Others	2(4%)	2(4%)

Values are number of subjects with percentage in parenthesis, n= number of subjects, -RFGDM = No Risk factor for gestational diabetes mellitus, +RFGDM= with risk factors for gestational diabetes mellitus. Majority of the study population were Hausa (80%), among which most of them (44%) had no formal education. Majority of them (62%) were women married to peasant farmers

Table 2. Age distribution (in years) and gestational ages (in weeks) of pregnant women with and without risk factors for gestational diabetes mellitus

Variable	Pregnant women +RFGDM (n= 50)	Pregnant women - RFGDM(n= 50)
Gestational age (weeks)		
16-20	10(10%)	14(28%)
20-24	14(28%)	13(26%)
24-28	26(52%)	23(46%)
Total	50(100%)	50(100%)
Age Class of pregnant women (years)		
18-25	18(36%)	30(60%)
26-32	25(50%)	8(16%)
33-39	5(10%)	10(20%)
40-46	2(4%)	2(4%)
Total	50(100%)	50(100%)

Values are number of subjects with percentage in parenthesis, n= number of subjects, -RFGDM = No Risk factor for gestational diabetes mellitus, +RFGDM= with risk factors for gestational diabetes mellitus. Majority of the study population (46%) had pregnancies with gestational age between 24-28 weeks. Majority of the study population (60%) were within the age of 18-25 years

Table 3. Mean urinary levels of microalbumin and serum levels of random plasma glucose, albumin, urea and creatinine in pregnant women with and without risk factors for gestational diabetes mellitus

Parameters	Pregnant women – RFGDM (n= 50)	Pregnant women + RFGDM (n= 50)
RPG (mmol/L)	4.33 ± 0.14	5.84 ± 0.16
MA(mg/L)	17.31± 0.64	56.36 ± 8.44*
Albumin(g/dL)	2.78 ± 0.09	3.06 ± 0.05
Urea (mmol/L)	3.47 ± 0.13	4.10 ± 0.15
Creatinine(mg/dL)	0.63 ± 0.01	0.70 ± 0.03

* (P < .001). n= number of subjects.

-RFGDM = No Risk factor for gestational diabetes mellitus, +RFGDM= with risk factors for gestational diabetes mellitus, RPG- Random plasma Glucose, MA- Microalbuminuria

Table 4. Mean levels of BMI, blood pressure, family history of diabetes and family history of hypertension in pregnant women with and without risk factors for gestational diabetes mellitus

Risk factors	Pregnant women - RFGDM (n= 50)	Pregnant women + RFGDM (n= 50)
BMI (kg/m ²)	22.98 ± 0.48	37.89 ± 0.60*
Systolic blood pressure (mm Hg)	124 ± 5.8	128.36 ± 11.3
Diastolic blood pressure (mm Hg)	79.2 ± 5.9	79.2 ± 10.1
Gravidity	P: 23(46%), M: 27(54%)	P: 29(58%), M: 21(42%)
Familyhistoryof diabetes	0	50
Familyhistoryof hypertension	3	35

* (P < .001)

n= number of subjects. BMI= Body Mass Index.

-RFGDM = No Risk factor for gestational diabetes mellitus, +RFGDM= with risk factors for gestational diabetes mellitus. mm Hg= millimeters of mercury. Kg= kilograms. M² =meters square.

P= primigravida. M= multigravida

Table 5. Correlation of urinary microalbumin to serum urea, creatinine, albumin and random plasma glucose

	Urea	Creatinine	Albumin	RBG
Microalbuminuria	r= 0.261 p= 0.09	r= 0.087 p= 0.387	r= 0.045 p= 0.656	r= 0.126 p= 0.212

In the this study, urine microalbumin levels was higher ($P<0.001$) in pregnant women with risk factors for gestational diabetes than in the control, and this is associated with increased in body mass index (BMI) of patients as compared to that of controls. Increased BMI is closely associated with microalbuminuria. Study on Type 2 diabetes mellitus patients reported a correlation between obesity and microalbuminuria [26]. The high mean urinary microalbumin level gotten from this study is similar to the findings of Shenaz [27] in type II diabetes mellitus in Karachi.

Random plasma glucose concentration for both women with risk factors for gestational diabetes mellitus and the control shows no statistically significant difference ($P<0.001$). This normoglycaemia seen in the study population may be due to the fact that these pregnant women only have the presence of risk factors for GDM such as family history of DM, high blood pressure, high BMI, etc; but these women do not have GDM.

In the current study, the slightly low serum albumin levels in the study population and that of controls shows no statistically significant difference ($P<0.001$). This is similar to the report of [28] which reported low serum albumin level in pregnant women. The decrease in serum albumin of the study population could be due to alteration in protein metabolism which is associated with gestational diabetes mellitus, or the socio economic status.

In the current study, serum urea and creatinine levels of the study population and the control shows no statistically significant difference ($P<0.001$). Similar findings [29,30] reported normal serum urea and creatinine levels in these category of patients but is lower than the one reported by [28]. This variation can be due to environmental factors and the socio economic status of the study population.

5. CONCLUSION

The present study concludes that the prevalence of microalbuminuria in women with risk factors for gestational diabetes mellitus in Sokoto, Nigeria was found to be 22%. The urinary microalbumin level is higher in pregnant women with risk factors for gestational diabetes mellitus than in the controls (pregnant women without risk factors for gestational diabetes mellitus). There is

a significant positive correlation between microalbuminuria and obesity ($BMI \geq 30 \text{ Kg/m}^2$).

The current study on microalbuminuria in pregnant women with risk factors for GDM is the first of its kind in our locality. Previous studies focused on microalbuminuria in pregnant women with GDM (not pregnant women with risk factors for GDM).

CONSENT

It is not applicable.

ETHICAL APPROVAL

Ethical approval was obtained from the Ethical Committee of Specialist Hospital, Sokoto and Maryam Abacha Women and Children Hospital, Sokoto before the commencement of the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Metzger BE, Lee KM. Detection and diagnostic strategies for gestational diabetes mellitus. In Hod, Moshe; Jovanovic, Lois; Renzo, Gian Carlo Di; de Leiva, Alberto; and Langer, Oded (Eds.), Textbook of Diabetes and Pregnancy, Second Edition, Informa UK Ltd. 2008;179-187.
2. National Institute of Diabetes and Digestive and Kidney Disease (NIDDK). Gestational diabetes. [Online]; 2014. Available:www.niddk.nih.gov/health-information/gestationaldiabetes [Accessed 23 June 2017]
3. International Diabetes Federation (IDF). Diabetes Atlas. Eighth Edition; 2017.
4. Langer Oded. Gestational diabetes: The consequences of not-treating. Textbook of Diabetes and Pregnancy, Second Edition, Informa UK Ltd. 2008;130-140.
5. Massi-Benedetti, Massimo F, Marco O, Renzo G. Management of gestational diabetes mellitus. Textbook of Diabetes and Pregnancy, Second Edition, Informa UK Ltd. 2008;188-195.
6. Qi Lian-Wen, Liu E-Hu, Chu Chu, Peng Yong Bo, Cai Hai-Xia, Li Ping. Anti-diabetic agents from natural products, an update

- from 2004 to 2009. *Current Topics in Medicinal Chemistry*. 2010;10(4):434-457.
7. Fawole AO, Ezeasor C, Bello FA, Roberts A, Awoyinka BS, Tongo O, Adeleye JO, Ipadeola. Effectiveness of a structured checklist of risk factors in identifying pregnant women at risk of Gestational diabetes mellitus. A cross sectional study. *Nigerian Journal of Clinical Practice*. 2014;495-501.
 8. Adoke AU, Shehu CE, Ekele BA, Nwobodo EI. Gestational diabetes mellitus and outcome of pregnancy among women attending antenatal care clinic in a tertiary institution. Paper Presentation at Nigerian Medical Association Annual Scientific Conference Held at Giginya Hotel Sokoto; 2014.
 9. Gilmartin AH, Ural SH, Repke JT. Gestational diabetes mellitus. *Reviews in Obstetrics and Gynaecology*; 2008.
 10. Ibadin OM, Onunu A, Uniegbé E, Ugheoke J. Microalbuminuria in adolescence/young adult offspring of hypertensive Nigerian adults- A preliminary report. *Nigeria Journal of Clinical Practice*. 2004;7(2):60-64.
 11. Timothy CE, Peter C. Diabetic nephropathy. *Clinical Diabetes*. 2000;18:10-11.
 12. Olatunde LO, Arogundade FA, Balogun MO, Akinsola A. Microalbuminuria and its clinical correlates in essential hypertension. *Nigeria Journal for Health Science*. 2002;2:25-29.
 13. Hitha B, Pappachan JM, Balachandran, Pillai H, Sujathan P, Ramakrishna CD, Jayaprakash K, Raihanathul KJ. Microalbuminuria in patients with essential hypertension and its relation to target organ damage: An Indian experience. *Saudi Journal of Kidney Diseases and Transplant*. 2008;19(3):411-414.
 14. Wayne DC, Tanya MO, Malcom C, Richard JM, George J. Earlier detection of microalbuminuria in diabetic patients using a new urinary albumin assay. *Kidney International*. 2004;65:1850-1855.
 15. United Nations Funds for Population Activities (UNFPA). *Population Projection and Health Services in Sokoto State, Nigeria*; 2016.
 16. Elving LD. Microalbumin and freezing. *Clinical Chemistry*. 1989;35(2):308.
 17. Trinder P. *Annals of clinical biochemistry*; 6:24: Quoted in *cheese rough*. M. *Medical Laboratory Manual for Tropical Country*; Vol. 1 (2nd Edition); ELBS, Cambridge. 1969;558-545.
 18. Doumas BT, Watson WA, Biggs HG. Albumin standards and measurement of serum albumin with bromocresol green. *Clinica Chimica Acta*. 1971;31(1):87-96.
 19. Burtis CA, Ashwood ER, Bruns DE, Sawyer BE. *Fundamentals of clinical chemistry* (6th Ed). Saunders Elsevier, Texas. 2008;376.
 20. Bunza FU, Mainasara AS, Dallatu MK, Bunza JM, Wasagu IZ. Prevalence of microalbuminuria among diabetic patients in Usmanu Danfodiyo University Teaching Hospital, Sokoto. *Bayero Journal of Pure and Applied Sciences*. 2014;7(1):1-5.
 21. Rodney CP, Renee D, Jeffrey R, David B, Chotip V, Ronald Acton. Prevalence and risk factors of microalbuminuria in a cohort of African – American Women with Gestational Diabetes. *Diabetes Care*. 2001;24:1764-1769.
 22. Adebisi SA, Okesina B, Abu E. Microalbuminuria in type 2 diabetic patients in Ilorin, Nigeria. *Diabetes International (Middle East/ Africa Edition)*. 2001;11(3):93-95.
 23. Orluwene CG, Momoh MO. Screening for microalbuminuria in newly diagnosed type II Diabetes at a staff clinic in Port Harcourt. *Port Harcourt Medical Journal*. 2008;3(1): 10-14.
 24. Iwalokun BA, Ogunfemi MK, Gbajabiamila B, Olukosi YA. Incidence and Evaluation of risk factors of microalbuminuria among diabetes and non- diabetes in Lagos, Nigeria. *Nigerian Journal of Health and Biomedical Sciences*. 2006;5(1):7-8.
 25. Newman DJ, Mattock MB, Dawnay AB, Kerry S, McGuire A, Yaqoob M. Systematic review on urine albumin testing for early detection of diabetic complications. *Health Technology Assess*. 2005;9:3-6.
 26. Mokdad AH, Ford ES, Bowman BA, Dietz, WH, Vinicor F, Bales VS. Prevalence of obesity, diabetes, and obesity related health risk factors. *Journal of American Medical Association*. 2003;289:76-79.
 27. Shenaz S, Jawed AB, Syed SH, Tehseen I, Tahseen K, Muhammad Baig. Prevalence of microalbuminuria with relation to glycemic control in type-2 diabetic patients in Karachi. *Journal of Ayub Medical College Abbottabad*. 2009;21(3):83-86.

28. Vari K, Sasirekha G. Biochemical markers of renal and hepatic function in gestational diabetes mellitus. *International Journal of Clinical Biochemistry and Research*. 2017;4(1):81-84.
29. Emre SG, Nuri D, Leyla Mollamahmutoglu. Relationship between serum uric acid, creatinine, albumin and gestational diabetes mellitus. *Clinical Chemistry and Laboratory Medicine*. 2006;44(8):974-977.
30. Radhai K, Zakkia K, Khalid J, Khurshi Ali. Effect of gestational diabetes mellitus on blood sugar, liver enzymes and renal function test. *Journal of Ayub Medical College Abbottabad*. 2012;24(2):95-98.

© 2019 Dallatu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/47454>