



## Codeine-mediated Haematotoxicity, Hepatotoxicity and Nephrotoxicity in Male Albino Rats

Abiodun Olusoji Owoade<sup>1\*</sup>, Abdulrahman Abdullateef<sup>1</sup>, Adewale Adetutu<sup>1</sup>  
and Olubukola Sinbad Olorunnisola<sup>1</sup>

<sup>1</sup>Department of Biochemistry, Ladoko Akintola University of Technology, Ogbomosho, Nigeria.

### Authors' contributions

This work was carried out in collaboration among all authors. Author AOO designed the study, performed the statistical analysis and wrote the protocol. Author AA carried out the investigation and wrote the first draft of the manuscript. Author AA managed the analyses of the study and the literature searches. Author OSO wrote the final draft. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/AJRIMPS/2019/v6i330102

#### Editor(s):

- (1) Dr. Davina Hijam, Department of Biochemistry, Regional Institute of Medical Sciences, India.  
(2) Dr. Lokendra Bahadur Sapkota, Assistant Professor, Department of Biochemistry, Chitwan Medical College, Bharatpur, Nepal.

#### Reviewers:

- (1) Dalia Mamdouh Mabrouk, National Research Centre, Egypt.  
(2) Michael Guranieri, Neurosurgery Johns Hopkins, USA.  
(3) Veeravan Lekskulchai, Srinakharinwirot University, Thailand.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48042>

Original Research Article

Received 11 January 2019  
Accepted 24 March 2019  
Published 30 March 2019

### ABSTRACT

**Objectives:** This study aimed to investigate the effects of codeine administration on some haematological and biochemical indices in rats.

**Materials and Methods:** Therapeutic dose (5 mg/kg/day), high dose (25 mg/kg/day) and extreme dose (50 mg/kg/day) of codeine were administered orally to rats for 28 days. Twenty-four hours after the last codeine administration, blood, liver and kidney were removed from the animals after an overnight fast and analysed for their haematological and biochemical parameters.

**Results:** Results obtained revealed that codeine administration significantly reduced the levels of white blood cells (WBC), red blood cell (RBC) and platelet count (PLT) and increased the levels of mean corpuscular haemoglobin (MCH) and mean corpuscular volume (MCV) while it resulted in non-significant changes in other haematological parameters examined when compared with control rats. Codeine intake significantly increased plasma levels of alkaline phosphatase (ALP), aspartate aminotransferase (AST), creatinine and urea while its reduced total protein levels. Hepatic and

\*Corresponding author: E-mail: aowoode@lajutech.edu.ng;

renal thiobarbituric acid reactive substances (TBARS) levels were significantly increased by codeine administration while levels of endogenous antioxidant enzymes superoxide dismutase (SOD), catalase (CAT), and reduced glutathione (GSH) were reduced.

**Conclusions:** This study confirmed the risk of increased oxidative stress, haematotoxicity, hepatotoxicity and nephrotoxicity due to codeine administration. Although codeine is reported to be effective in pain management, its toxicity should be kept in mind.

*Keywords: Codeine; haematological; oxidative stress; hepatotoxicity; nephrotoxicity.*

## 1. INTRODUCTION

Codeine is an opiate analgesic that is commonly used in a formulation with paracetamol (acetaminophen), although combinations with other analgesics like acetylsalicylic acid, ibuprofen, caffeine, barbiturates and sedative antihistamines may also exist [1]. The drug is often used to suppress a cough either alone or by combining it with other drugs [2]. The major pharmacological effects of codeine such as analgesia, drowsiness, mood changes, respiratory depression, nausea, and decreased gastrointestinal motility are produced on the central nervous system (CNS) and gastrointestinal tract [3].

The metabolism of codeine occurs mostly in the liver, and to a lesser extent in the intestine and CNS [1]. Although codeine metabolism resulted in several metabolites, it is morphine, a product of codeine O-demethylation by enzyme cytochrome P450 2D6 (CYP2D6), that is responsible for its analgesic effect [4,5]. Codeine dosage is highly regulated, its overdose could cause depressive effects on the central nervous system or death from respiration arrest. The adult minimum lethal oral dose for codeine is estimated to be 0.5–1.0 g, i.e 17–34 pills containing 30 mg codeine [6]. The serum codeine concentrations exceeding 0.3 mg/L have been reported to cause toxicity, while concentrations above 1.6 mg/L are considered to be lethal [2].

The abuse and misuse of prescription opioids such as codeine have reached an alarming rate in the last ten years. In the United States, it was reported that about 1.2 million visit emergency department (ED) as a result of non-medical use of prescription medications in 2011 alone [7]. The production and importation of cough syrup that include codeine as an ingredient was ban by the Nigerian Government in 2018 due to concerns regarding its use by youths to get intoxicated [8]. Many addicts of codeine in many countries are into the habit of using the drug

every day without doctor prescriptions which is of great concerns. Therefore, this study assesses the toxic effects of different doses, including overdoses of codeine, on haematological parameters, biochemical changes and oxidative damage in the liver and kidney of rats exposed to normal, high and extreme doses of codeine orally for 28 days.

## 2. MATERIALS AND METHODS

### 2.1 Materials

Centrifuge machine, human automated haematology system analyzer (ERMA PCE 210, ERMA, Japan), weighing balance, dissecting sets, cuvette, spectrophotometer, pH meter, refrigerator, homogenizer, razor blade, 1 ml syringes, 2 ml syringes, and 5 ml syringes, surgical gloves, cotton wool, measuring cylinder, test tubes, beaker, spatula, plastic cages, EDTA bottles, plain sample bottles.

### 2.2 Reagents

Thiobarbituric acid (TBA), nicotinamide adenine dinucleotide reduced (NADH) and Codeine were obtained from Sigma–Aldrich Chemical Co. Ltd. (England). Nitrobluetetrazolium (NBT), 5,5'-Dithiobis (2-nitrobenzoic acid) (DTNB) are the product of Fluka (Buchs, Switzerland). All other chemicals used were analytical grade.

### 2.3 Animals

Twenty (20) male Wistar rats with an average weight of 170-200 g were used for the experiments. They were housed in the Ladoke Akintola University of Technology, (LAUTECH) animal house. They were allowed fourteen (14) days to acclimatize before the commencement of drug administration. The animals were maintained on a standard pellet diet throughout the acclimatization and administration period. The animal experimental procedures were conducted in accordance with the National Institutes of Health guide for the care and use of

laboratory animals (NIH Publications No. 8023) revised in 2002 and approved by the institutional research committee.

## 2.4 Experimental Design

Twenty (20) male wistar strain albino rats were divided into four groups of five rats each according to their weight. Group I labelled control received saline solution for 28 days through the oral route. Group II labelled normal codeine received a normal dose of codeine at 5 mg/kg/day body weight of rat. Group III labelled high codeine received a high dose of codeine at 25 mg/kg/day. Group IV labelled extreme codeine received an extreme dose of codeine at 50 mg/kg/day. Codeine was constituted in saline solution and administered through the oral route. During the experiment, the animals were allowed free access to food and distilled water. After 28 days of codeine treatment and after an overnight fast, animals were sacrificed by cardiac puncture under light ether anaesthesia into ethylene diamine tetra-acetic acid (EDTA) sample bottles for haematological analysis and heparinised sample bottles for biochemical analysis. Liver and kidney were removed from the animals for biochemical analyses. Blood samples in heparinized bottles were centrifuged to separate plasma and red blood cells. All samples were stored at -20°C until analysed.

## 2.5 Haematological Study

Freshly collected blood samples in EDTA bottles were analysed for haematological assay using an automatic haematological assay analyser (ERMA PCE 210, ERMA, Japan). Different tested haematological parameters were as follows: White Blood Cell (WBC), Red Blood Cells (RBC), Haemoglobin (HGB), Haematocrit (HCT), Red cells (RDW%), Red cells Distribution Width (RDW), Mean Corpuscular Haemoglobin (MCH), Mean Corpuscular Haemoglobin Concentration (MCHC), Platelet (PLT), Mean Platelet Volume (MPV), Mean Corpuscular Volume (MCV), Platelet crit (PCT), Platelet distribution width (PDW).

## 2.6 Determination of Blood Biochemical Parameters

Plasma concentrations of alkaline phosphatase (ALP), aspartate aminotransferase (AST), alanine aminotransferase (ALT), urea, creatinine and total protein were determined using

enzymatic kits (CYPRESS® Diagnostics, Langdorp, Belgium) according to the manufacturer's instructions.

## 2.7 Preparation of Liver and Kidney Homogenates

Prior to biochemical analyses, the liver and kidney samples were cut into small pieces and homogenized in Phosphate buffer saline (PBS) with a homogenizer to give a 10% (w/v) liver and kidney homogenate. The homogenates were then centrifuged at 12,000 rpm for 15 min. The supernatant obtained was used for the assay of superoxide dismutase, catalase, reduced glutathione, thiobarbituric acid reactive substances (TBARS) content, and protein estimation.

## 2.8 Determination of Hepatic and Renal Antioxidant Enzyme Activities and MDA Levels

Hepatic and renal superoxide dismutase (SOD) activities were assayed in the tissue homogenates by the method of Kakkar, et al. [9] at 560 nm. One unit of enzyme activity was defined as that amount of enzyme which caused 50% inhibition of nitrobluetetrazolium reduction/mg protein. Catalase (CAT) activity was determined at room temperature by using the method of Aebi [10] and the absorbance of the sample was measured at 240 nm in a UV spectrophotometer. The concentration of reduced glutathione (GSH) in liver and kidney homogenates was measured, as described by Jollow et al. [11]. The extent of lipid peroxidation was estimated as the concentration of thiobarbituric acid-reactive product malondialdehyde (MDA), using the method of Draper and Hadley, [12]. All of the enzyme activities were expressed as per mg of protein and the tissue protein was estimated according to the method of Lowry et al., [13], using bovine serum albumin (BSA) as a standard.

## 2.9 Statistical Analysis

Results are expressed as mean  $\pm$ S.E.M. The levels of homogeneity among the groups were assessed using One-way Analysis of Variance (ANOVA) followed by Turkey's test. All analyses were done using Graph Pad Prism Software Version 5.00 and p values < 0.05 were considered statistically significant.

### 3. RESULTS

#### 3.1 Haematological Parameters

The effects of codeine administration on haematological parameter were depicted in Table 1. No significant changes in the parameters of HGB, HCT, RDW-CV, MPV, and PDW were found when compared with control animals. However, administration of codeine significantly lower ( $p < 0.05$ ) white blood cell (WBC) count, red blood cell (RBC) count, and platelet (PLT), while the value of mean corpuscular volume (MCV) and Mean Corpuscular Haemoglobin (MCH) were increased when compared with control animals.

#### 3.2 Effect of Codeine Administration on Blood Biochemical Parameters

Administration of codeine at normal, high and extreme doses significantly increased the activity of ALP by 75.39%, 149.36% and 122.65% respectively and AST activity by 31.82%, 83.41% and 145.47% respectively when compared with the normal rats. The plasma concentration of creatinine and Urea were also significantly increased by all the three doses of codeine while total proteins level was decreased by

administration of normal, high and extreme doses of codeine by 30.35%, 48.02% and 38.84% respectively when compared with the normal rats (Table 2).

#### 3.3 Effect of Codeine Administration on TBARS Levels

Hepatic TBARS levels of rats treated with normal, high and extreme doses of codeine were dose-dependently significantly increased by 58.26%, 131.42% and 234.74% respectively when compared with the normal rats. Similarly, administration of codeine at normal, high and extreme doses significantly increases renal TBARS levels by 57.59%, 114.14% and 157.23% respectively when compared with the control rats (Fig. 1).

#### 3.4 Effect of Codeine Administration on SOD Activity

Administration of codeine at normal, high and extreme doses significantly reduced hepatic SOD levels by 39.06%, 61.74% and 57.29% respectively and reduced renal SOD levels by 37.70%, 50.27% and 62.04% respectively when compared with normal rats (Fig. 2).

**Table 1. Effect of codeine administration on haematological parameters of rats**


Parameters	Control	Normal codeine	High codeine	Extreme codeine
WBC ( $\times 10^9/L$ )	8.30 $\pm$ 0.35	4.18 $\pm$ 0.22 **	4.94 $\pm$ 0.74 **	4.56 $\pm$ 0.34 **
HGB (g/dl)	10.58 $\pm$ 0.39	9.88 $\pm$ 0.08	10.66 $\pm$ 0.64	10.67 $\pm$ 0.13
RBC ( $\times 10^{12}/L$ )	6.92 $\pm$ 0.31	5.59 $\pm$ 0.05 **	5.51 $\pm$ 0.33 **	5.47 $\pm$ 0.11 **
HCT (%)	31.64 $\pm$ 0.97	29.00 $\pm$ 0.32	32.00 $\pm$ 1.58	32.00 $\pm$ 0.32
MCV (fl)	55.50 $\pm$ 0.87	56.48 $\pm$ 0.54	67.70 $\pm$ 1.15 **	67.70 $\pm$ 1.15 **
MCH (pg)	15.28 $\pm$ 0.31	17.60 $\pm$ 0.19	19.30 $\pm$ 0.28 **	19.42 $\pm$ 0.22 **
MCHC (g/L)	275.64 $\pm$ 1.93	313.00 $\pm$ 2.21	286.60 $\pm$ 2.6	290.80 $\pm$ 2.46
RDW-CV (%)	17.22 $\pm$ 0.31	16.30 $\pm$ 0.25	17.84 $\pm$ 0.38	17.32 $\pm$ 0.16
RDW-SD (fl)	31.82 $\pm$ 0.15	31.60 $\pm$ 0.43	39.18 $\pm$ 1.01	37.42 $\pm$ 0.47
PLT ( $\times 10^9/L$ )	583.00 $\pm$ 1.26	592.86 $\pm$ 18.11	544.67 $\pm$ 14.10	405.50 $\pm$ 6.30 **
MPV (fl)	7.00 $\pm$ 0.06	6.80 $\pm$ 0.11	6.88 $\pm$ 0.10	7.16 $\pm$ 0.10
PDW	16.08 $\pm$ 0.32	16.14 $\pm$ 0.19	15.96 $\pm$ 0.08	15.98 $\pm$ 0.13
PTC (%)	0.39 $\pm$ 0.00	0.40 $\pm$ 0.02	0.33 $\pm$ 0.01	0.33 $\pm$ 0.04

Each value represents the mean of five rats. \*\* = significantly different from control ( $p < 0.05$ ).


**Table 2. Effect of codeine administration on blood biochemical parameters**

PARAMETERS	Control	Normal codeine	High codeine	Extreme codeine
ALP (IU/L)	100.74 $\pm$ 7.86	176.69 $\pm$ 11.57 **	251.21 $\pm$ 12.52 **	224.30 $\pm$ 15.76 **
AST (IU/L)	44.25 $\pm$ 1.15	58.33 $\pm$ 2.42 **	81.16 $\pm$ 4.45 **	108.62 $\pm$ 3.56 **
Creatinine (mg/dL)	0.83 $\pm$ 0.08	1.61 $\pm$ 0.25 **	2.44 $\pm$ 0.21 **	2.19 $\pm$ 0.32 **
Urea (mg/dL)	23.79 $\pm$ 1.64	46.46 $\pm$ 3.94 **	41.60 $\pm$ 3.51 **	33.39 $\pm$ 2.91 **
Total protein (g/dL)	8.60 $\pm$ 0.73	5.99 $\pm$ 0.36 **	4.47 $\pm$ 0.50 **	5.26 $\pm$ 0.47 **

Each value represents the mean of five rats. \*\* = significantly different from control ( $p < 0.05$ ).



**Fig. 1. Effect of codeine administration on hepatic and renal TBARS level of rats**  
 Values are mean  $\pm$  SEM (n=5). \*\* = significantly different from control ( $p < 0.05$ )



**Fig. 2. Effect of codeine administration on hepatic and renal SOD activity of rats**  
 Values are mean  $\pm$  SEM (n=5). \*\* = significantly different from control ( $p < 0.05$ )

### 3.5 Effect of Codeine Administration on Catalase Activity


Administration of codeine at normal, high and extreme doses significantly reduced

hepatic catalase levels by 28.93%, 40.89% and 53.33% respectively and reduced renal catalase levels by 45.92%, 61.73% and 40.30% respectively when compared with normal rats (Fig. 3).


### 3.6 Effect of Codeine Administration on GSH Activity

Hepatic GSH levels of rats treated with normal, high and extreme doses of codeine were significantly reduced by 37.04%, 65.21% and

63.58% respectively when compared with the normal rats. Similarly, administration of codeine at normal, high and extreme doses significantly reduced renal GSH levels by 33.06%, 49.73% and 66.71% respectively when compared with the control rats (Fig. 4).



**Fig. 3. Effect of codeine administration on hepatic and renal catalase activity of rats**  
 Values are mean  $\pm$  SEM (n=5). \*\* = significantly different from control ( $p < 0.05$ )



**Fig. 4. Effect of codeine administration on hepatic and renal GSH activity of rats**  
 Values are mean  $\pm$  SEM (n=5). \*\* = significantly different from control ( $p < 0.05$ )

#### 4. DISCUSSION

Codeine (Opioid) is an analgesic mainly used as an antituitive drug and to manage mild to moderate pain [14,15]. It is, however, a drug of abuse because of its stimulatory effect on CNS among some adults [16]. Toxic effects of codeine use have been reported, although little is known about codeine toxicity mechanisms [2,17]. In this study, the toxic effects of codeine were examined in animal models. Codeine was studied as a drug and not as analgesics because alarming misuse of codeine recently made Nigeria Government ban production and importation of cough syrup that has codeine as an ingredient [8]. Therefore, this study evaluated the toxicity of codeine on systemic body organs because of people use of codeine without doctor prescriptions.

Free radicals and reactive oxygen species are generated by chemicals and pollutants such as factory waste, toxic gases and they are known to disrupt biochemical and haematological parameters in organisms [18]. Disruption of haematological parameters could provide valuable information and insight into the diagnosis of various diseases and pathological conditions. The deviation from normal haematological parameters levels represents the presence of toxicity or disease conditions [19]. The decrease in red blood cell count (RBC) level could be a result of an imbalance between its production and loss [20]. In this study, codeine administration caused a significant reduction in red blood cell counts (RBC). The observed decrease in the number of RBCs suggest that codeine administration resulted in blood loss due to serious gastrointestinal tract bleeding, red blood cell haemolysis and poor iron absorption in the intestine.

Codeine administration also resulted in a reduction of WBC of experimental animals in this study. White blood cells fight infections, defend the body against foreign organisms' invasion and produce antibodies in immune response [21]. Animals with low WBC are at high risk of disease infection, while high WBC results in high resistance to diseases [21]. The reduction of WBC by codeine observed in this study agrees with pervious study which revealed that abuse or long-term use of opium supresses the immune system and individuals are more susceptible to infectious disease [22].

Blood platelets are involved in blood clotting and its low level will prolong the process of clot-formation resulting in excessive blood loss during injury. Although, there was no significant variation in platelet concentration of rats administered with normal and high doses of codeine in this study, however, the extreme dosage of codeine significant reduce platelet concentration. A decreased number of platelet (thrombocytopenia) by codeine in this study is in supports of pervious work which observed that morphine administration induced thrombocytopenia [23]. Codeine administration also resulted in increased in the levels of mean corpuscular haemoglobin (MCH) and mean corpuscular volume (MCV). We strongly believe that disruption in haematological parameters observed in this study may be due to increased population of unquenched free radicals caused by codeine administration. The changes observed in other haematological parameters such as HGB, HCT, RDW%, RDW, MCHC, MPV,PCT, PDW in this study were largely found to be non-significant, an observation that may be different if codeine administration period was much longer than 28 days used in this study.

In this study, many biochemical parameters on liver and kidney functions were determined in plasma samples to assess damage to metabolizing organs. The increased in the activities of ALT, aspartate aminotransferase (AST), ALP and lactate dehydrogenase (LDH) have been reported in previous studies following exposure to opioids, including morphine and tramadol [24-26]. Administration of codeine in this study significantly increased ALP and AST activities which are in conformity with previous research that revealed that AST, ALP and ALT activities in plasma increased significantly in an addicted patient of opioid [27]. The liver is an organ that detoxified toxic elements and chemical drugs in the body, the increase in the activities AST and ALP in plasma in this study are indicative of liver damage [28]. The increased secretion of these liver enzymes may be accompanied by acute cell necrosis, therefore, the increased plasma level of these enzymes in rats treated with codeine could be due to necrosis or damage to liver cell membrane which leak the enzymes into the blood circulation [29].

The level of plasma creatinine is used to determined glomerular filtration rate and serves as renal function assessment [30]. Codeine administration significantly increased plasma

creatinine level of rats in this study and this can be taken as evidence of renal damage because the high level of creatinine in the blood implies a loss of kidney function in ensuring creatinine excretion. Similarly, administration of codeine in this study increased blood urea concentration. Urea is a nitrogenous waste and product of protein and amino acid metabolism, it is eliminated from the body through urinary excretion. It is an important clinical parameter because it can be used to determine the nephrotoxic profile of xenobiotics. The increased in blood urea concentration observed in this study following codeine administration agrees with previous research [31] and it is an indication of renal toxicity which might have instigated decrease in glomerular filtration rate leading to the build-up of creatinine and urea in the blood.

There was a decrease in plasma total protein in rats treated with codeine in this study this is in support of previous research finding which showed decrease in plasma total protein levels in opium dependent participants when compared to the control group [32]. The clinical diagnosis has shown that a decrease in plasma concentrations of protein characterized by significant increases in the urinary excretion of protein and albumin are indicators of renal dysfunction [33]. Therefore, the decrease in plasma total protein observed in this study can be taken as an indication of kidney damage.

Administration of codeine resulted in increased levels of MDA the last metabolite of lipid peroxidation chain, and inhibition of the antioxidant enzymes, superoxide dismutase (SOD), catalase (CAT) and reduced glutathione (GSH) in liver and kidney of rats. Elevated levels of MDA have been reported to be an indication of an increase in free radical generation and it is considered a useful measure of oxidative stress status [34]. SOD, CAT and GSH are important antioxidant enzymes which played a pivotal role in scavenging of oxidative free radicals [35]. The inhibition of these antioxidant enzymes observed in this study could be linked to exhaustion of these enzymes as a result of oxidative stress caused by codeine administration.

The toxic effect of codeine administration leads to a large population of unquenched free radicals leading to the state of oxidative stress. Oxidative stress form when there is an imbalance between free radical generating and scavenging systems has been implicated

in the pathogenesis of a wide range of disorders, including neurodegenerative disorders, cardiovascular diseases, cancer, and ageing [36].

## 5. CONCLUSION

Our results evidence that codeine administration may cause haematotoxicity, hepatotoxicity and nephrotoxicity and as such, its use should be limited to prescription only. Our findings underlined the need to avoid indiscriminately and prolong use of codeine, since prolonged daily use of the drug either at a therapeutic dose or the extreme dose may lead to damage accumulation.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

As per international standard or university standard ethical approval has been collected and preserved by the authors.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Frost J, Løkken TN, Helland A, Nordrum IS, Slordal L. Post-mortem levels and tissue distribution of codeine, codeine-6-glucuronide, norcodeine, morphine and morphine glucuronides in a series of codeine-related deaths. *Forensic Sci. Int.* 2016;262:128–137.
2. Frost J, Helland A, Nordrum IS, Slordal L. Investigation of morphine and morphine glucuronide levels a cytochrome P450 isoenzyme 2D6 genotype in codeine-related deaths, *Forensic Sci. Int.* 2012; 220:6–11.
3. Jaffe H, Martin WR. Opioid analgesics and antagonists. *The Pharmacological Basis of Therapeutics* (Gilman AG, Goodman LS, Gilman A, Eds.). 6<sup>th</sup> Ed., Macmillian Co., New York. 1980;494-534.
4. Poulsen L, Brosen K, Arendt-Nielsen L, Gram LF, Elbaek K, Sindrup SH. Codeine and morphine in extensive and poor metabolizers of sparteine:



- Pharmacokinetics, analgesic effect and side effects. *Eur. J. Clin. Pharmacol.* 1996; 51:289–295.
5. Sindrup SH, Arendt-Nielsen L, Brosen K, Bjerring P, Angelo HR, Eriksen B, Gram LF. The effect of quinidine on the analgesic effect of codeine. *Eur. J. Clin. Pharmacol.* 1992;42:587–591.
  6. Baselt RC. *Disposition of Toxic Drugs and Chemicals in Man*, 8<sup>th</sup> Ed., Biomedical Publications, Foster City, CA; 2008.
  7. DAWN, Drug Abuse Warning Network 2011: National estimates of drug-related emergency department visits; 2013.
  8. O'Grady S. Nigeria bans codeine cough syrup to stop an addiction epidemic. But something worse could take its place. *The Washington Post* May 4<sup>th</sup>, 2018. Available: <https://www.washingtonpost.com/news/worldviews/wp/2018/05/04/> Assessed: 28/01/2019
  9. Kakkar P, Das B, Viswanathan PN. A modified spectrophotometric assay of superoxide dismutase, *Ind. J. Biochem. Biophys.* 1984;21:130–132.
  10. Aebi H. Catalase *in vitro*. *Methods Enzymol.* 1974;105:121–126.
  11. Jollow DJ, Mitchell JR, Zampaglione N, Gillette JR. Bromobenzene induced liver necrosis: protective role of glutathione and evidence for 3,4 bromobenzeneoxide as the hepatotoxic intermediate. *Pharmacology.* 1974;11:151–169.
  12. Draper HH, Hadley M. Malondialdehyde determination as index of lipid peroxidation. *Methods Enzymol.* 1990;186: 421–431.
  13. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 1951; 193:265–275.
  14. Gasche Y, Daali Y, Fathi M, Chiappe A, Cottini S, Dayer P, Desmeules J. Codeine intoxication associated with ultra rapid CYP2D6 metabolism. *N. Engl. J. Med.* 2004;351:2827–2831.
  15. Barnhart ER. *Physicians' Desk Reference*, Medical Economics Company, Oradell, NJ. 40<sup>th</sup> Ed. 1986.
  16. Fraser HF. Patterns of abuse of narcotics: An historical overview. In *drug abuse- Proceedings of the International Conference* Lea & Febiger, Philadelphia (Zarafonitis CJD, Ed.), 1972; 141-151.
  17. Schulz M, Schmoldt A. Therapeutic and toxic blood concentrations of more than 800 drugs and other xenobiotics. *Pharmazie.* 2003;58(7):447–474.
  18. Celik I, Suzek H. The hematological effects of methyl parathion in rats. *J. Hazard. Mater.* 2008;153:1117–1121.
  19. Oyedemi SO, Bradly G, Afolayan AJ. Toxicological effects of the aqueous stem bark extract of *Strychnos henningsii* Gilg in Wistar rats. *J. Nat. Pharm.* 2010;1: 33–39.
  20. Alabi QK, Akomolafe RO, Olukiran OS, Nafiu AO, Omole JG, Adefisayo AM, et al. Assessment of haematological and biochemical effects of Kolaviron in Male Wistar Rats. *Br J Pharm Res.* 2017;16(3): 1–14.
  21. Soetan KO, Akinrinde AS, Ajibade TO. Preliminary studies on the haematological parameters of cockerels fed raw and processed guinea corn (*Sorghum bicolor*). *Proceedings of 38<sup>th</sup> Annual Conference of Nigerian Society for Animal Production.* 2013;49-52.
  22. Roy S, Wang J, Kelschenbach J, Koodie L, Martha J. Modulation of immune function by morphine implications for susceptibility to injection. *Neuroimmune pharmacol.* 2006;1:77-89.
  23. Cimo PL, Hammond JJ, Moake JL. Morphine-induced immune thrombocytopenia. *Arch Intern Med.* 1982;142(4): 832-834.
  24. Ali OK, Ahmed AJS, Mawlood AG. Effects of tramadol on histopathological and biochemical parameters in male rabbits. *Am. J. Biol. Life Sci.* 2015;3:85–90.
  25. Hafez E, Issa S, Rahman SA. Parenchymatous toxicity of tramadol: A histopathological and biochemical study. *J. Alcohol Drug Depend.* 2015;3.
  26. Salahshoor MR, Khashiadeh M, Roshankhah S, Kakabaraei S, Jalili C. Protective effect of crocin on liver toxicity induced by morphine. *Res. Pharm. Sci.* 2016;11:120–129.
  27. Sajad M, Chawla R, Zargan J, Umar S, Sadaqat M, Khan HA. Cytokinetics of adult rat SVZ after EAE. *Brain Research.* 2011; 1371:140-149.
  28. Salashoor MR, Roshankhan S, Hosseni P, Jalili C. Genistein improves liver damage in male mice exposed to morphine. *Chinese Medical Journal (Engl).* 2018; 131(13):1598-1604.
  29. Drotman RB, Lawhorn GT. Serum enzymes as indicators of chemically

- induced liver damage. *Drug and Chemical Toxicology*. 1978;1(2):163-71
30. Perrone RD, Madias NE, Levey AS. Serum creatinine as an index of renal function: New insights into old concepts. *Clin. Chem*. 1992;38(10):1933-53.
  31. Hatem SM, Saussez G, Faille MD, Prist V, Zhang X, Dispa D, Bleyenheuft Y. Rehabilitation of Motor Function after stroke: A Multiple Systematic Review Focused on Techniques to stimulate upper extremity recovery. *Frontiers in Human Neuroscience*. 2016;10:442.
  32. Afarinesh MR, Haghpanah T, Divsalar K, Dehyadegary E, Shaaikh-Aleslami A, Mahmoodi M. Changes in serum biochemical factors associated with opium addiction after addiction desertion. *Addiction and Health*. 2014;6:138-145.
  33. Garg R, Yusuf S. Overview of randomized trials of angiotensin-converting enzyme inhibitors on mortality and morbidity in patients with heart failure. Collaborative Group on ACE Inhibitor trials. *JAMA*. 1995; 273(18):1450-1456.
  34. Pan HZ, Zhang H, Chang D, Li H, Sui H. The change of oxidative stress products in diabetes mellitus and diabetic retinopathy. *Br. J. Ophthalmol*. 2008;92(4): 548–551.
  35. Kruidenier L, Kuiper I, Van Duijn W, Mieremet-Ooms MA, van Hogeza RA, Lamers CB, et al. Imbalanced secondary mucosal antioxidant response in inflammatory bowel disease. *J Pathol*. 2003;201:17–27.
  36. Halliwell B, Gutteridge JMC. Oxidative stress: Adaptation, damage, repair and death. In: Halliwell B, Gutteridge JMC, Eds. *Free Radicals in Biology and Medicine*, 3<sup>rd</sup> Edition, Oxford University Press, Oxford. 1999;246-350.

© 2019 Owoade et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/48042>