



Effects of Continuous Deep-fat Frying on the Physicochemical Properties of Assorted Brands of Edible Cooking Oils Sold in Greater Metropolitan Kampala

Timothy Omara^{1,2,3*}, Erisa Kigenyi^{3,4}, Fortunate Laker^{3,4}, Monica Adokorach³, George Otim³, Raymond Kalukusu^{3,4}, Bashir Musau^{3,4}, Sarah Kagoya^{3,5} and Brenda Victoria Nakabuye^{4,6}

¹Department of Chemical Engineering, Faculty of Engineering and Informatics, University of Bradford, West Yorkshire, BDP 1DP7, UK.

²Department of Quality Control and Quality Assurance, Product Development Directory, AgroWays Uganda Limited, 34-60, Kyabazinga Way, Jinja, Uganda.

³Department of Chemistry, Faculty of Science, Kyambogo University, 1, Kyambogo, Kampala, Uganda.

⁴Department of Quality Control and Quality Assurance, Leading Distillers Uganda Limited, 660, Mengo, Kampala, Uganda.

⁵Department of Quality Control, Sweets and Confectionaries Section, Kakira Sugar Limited, 121, Jinja-Iganga Highway, Jinja, Uganda.

⁶Department of Food Processing Technology, Faculty of Science, Kyambogo University, 1, Kyambogo, Kampala, Uganda.

Authors' contributions

This work was carried out in collaboration among all authors. This was a concerted study. Authors TO, SK, BM and EK designed the study, Authors FL, MA and EK performed the statistical analysis, Authors TO, EK, RK, FL and BVN wrote the protocol and the first draft of the manuscript. Authors GO, MA, RK, BM and FL managed the analyses of the study. Authors GO, BVN, FL, MA and SK managed the literature searches. All the authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJACR/2019/V3i230086

Editor(s):

(1) Dr. Angélica Machi Lazarin, Professor, Department of Chemistry, State University of Maringá, Brazil.

Reviewers:

(1) F. J. Owuna, Usmanu Danfodiyo University, Nigeria.

(2) Antoninho Valentini, Federal University of Ceará, Brazil.

(3) Michael Akrofi Anang, University of Cape Coast, Ghana.

(4) Vera Lúcia Viana do Nascimento, Federal Institute of Piauí, Brazil.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48900>

Original Research Article

Received 07 March 2019

Accepted 17 May 2019

Published 25 May 2019

ABSTRACT

Aims: To investigate the effects of continuous deep fat frying of white (Irish) potatoes on the physical and chemical attributes of ten brands of edible cooking oils: Fortune Butto, Roki, Tamu, Best Fry, Mukwano, Golden Fry (hard oils); Sunseed, Sunny, Sunvita and Sunlite (soft oils) sold in Kampala, Uganda.

Place and Duration of the Study: Oil samples of approximate manufacturing dates were obtained from Mega Standard supermarket in Greater Metropolitan Kampala, Uganda. Oil samples were also obtained from local Irish potato fryers in Makindye division of Kampala during ten deep frying cycles. Irish potatoes was procured from Nakasero market, Kampala. Physicochemical analyses were performed at the Quality Control Laboratory of Mukwano Industries Limited, Kampala Industrial area, Kampala. The research was conducted between May 2018 to December 2018.

Methodology: 400g of Irish potato slices (1cm × 1cm × 3cm) were submersed in 1500mL of oil maintained at 140°C for 6 minutes in an Electric Deep Fryer with a frying time of 10 minutes. The color value (CV) and the acidification of the oils as free fatty acid (FFA), peroxide value (POV), paraanisidine value (AnV), iodine adsorption value (IV) and total oxidation (TOTOX) value before and between ten successive frying cycles were determined using ISO and AOCS official methods. The maximum number of reuses of an oil was estimated from the frying round before its POV or AnV surpassed the maximum permissible statutory or Codex Alimentarius limit for edible oils.

Results: For fresh oils, the statistical physicochemical parameter ranges were: CV (0.4R 3.4Y-7.7R 70Y), FFA (0.0430±0.30-0.1508±0.30), POV (0.5951±0.03-6.6134±0.23 meqO₂/Kg), AnV (0.90±0.01-4.30±0.19) and IV (57.62±0.17-128.35±0.02gl₂/100g). By the 10th fry, the ranges were CV (3.0R 23Y-20.4R 70Y), FFA (0.2286±0.01-0.4817±0.01), POV (11.1138±0.01-15.7525±0.01meqO₂/Kg), AnV (10.31±0.03-22.16±0.01) and IV (53.66±0.01-126.03±0.02gl₂/100g). Considering oxidizability as TOTOX values, frying stability of the selected brands of cooking oils during the frying cycles followed the order: Roki > Fortune Butto > Sunvita > Sunny > Sunlite > Mukwano > Tamu > Best Fry > Golden Fry > Sunseed.

Conclusion: Reuse of the oils for continuous frying of Irish potatoes on the same day can be done only up to 7 times on average for hard oils and 6 times for soft oils with the oils still regarded as safe for human consumption. Hard oils should be preferred to soft oils for deep frying of Irish potato chips. Further research should elucidate the variation of physicochemical properties of other oil brands on the Ugandan market such as Nile, Fortune, Kimbo, Star Fry, Cow boy and Ufuta and should use other food samples such as fish, cassava, chicken, sweet plantain, dough, meat and edible grasshoppers.

Keywords: Color index; free fatty acids; iodine absorption value; paraanisidine value; peroxide index; Kampala.

1. INTRODUCTION

Kampala, the central business district and the capital city of Uganda is an area of population with over 1.53 million metropolitans surviving on a vast array of quick foods such as ready-to-drink milk, tossed salads, snacks [1] and unique Ugandan deep-fried delicacies: Fried dough (*mandazi*), sweet plantain (*gonja*), edible grasshoppers (*nseene*), banana pancakes (*kabalagala*), Nile perch fish (*emputa*), chapatti (*kikomando*), sausages, chicken nuggets (*enkoko*), meat (*enyama*), Irish potatoes and cassava chips. The fries are prepared in repeatedly used oils and sometimes fresh oil is added to the used oils. In either case, recycling of oil enhances the innocuous contamination and

interaction of moisture and air (oxygen) with the oil [2-5].

A number of physicochemical changes occur during deep frying. The color value of oils for example, usually expressed as 1 × Red + 1 × Yellow Lovibond units, registers a drastic increase in both the red and yellow units during the incipient phases of heating. According to the results of Baby Latha and Nasirullah [6] using rice bran oil, a threefold increase in red units and nearly a fourfold increase in yellow units was recorded after 2 hours of frying while darkening occurred beyond 2 hours. Physicochemical properties and oxidative degradation of the rice bran oil during the initial 2 hours of heating registered a steep increment in POV from 0.2 to

6.3 meq O₂/kg, AnV from 5.04 to 19.4 and total polar components from 1 to 4.1% [6]. Diop et al. [7] investigated the effects of deep fat frying on chemical properties of three selected brands of oils (two peanut oils A and B and sunflower oil, C) common in Senegalese preparation of fried meat, fish and potatoes. Their findings revealed that frying affects the chemical stability of cooking oils. The acid value as reported increased after 40 minutes from 0.62 to 1.08 mg/kg after frying fish, from 0.39 to 0.73 mg/kg for meat and 0.37 to 0.51 mg/kg for potatoes. POV increased only slightly for oil A and sharply for oils B and C.

Xu et al. [8] compared the oxidative stability of camellia oil composed of saturated fatty acid (SFA), monounsaturated fatty acid (MSFA) and polyunsaturated fatty acid (PUFA) in a ratio of 1:7:1 during potato deep frying with palm and peanut oils composed of SFA, MSFA and PUFA in ratios of 4:4:1 and 2:4:4 respectively. Their evaluations of acid value, Iodine Value (IV), Peroxide value (POV), peranisidine value (AnV), total oxidation (TOTOX) value, tocopherols content and fatty acids composition of the oils registered little alteration of the fatty acid contents of camellia oil with alpha-tocopherol reported to be more thermally labile compared to gamma and delta tocopherols. They concluded that the stabilities of the oils as determined by oxidizability value followed the sequence camellia oil > palm oil > peanut oil. The initially highest recorded AnV was in palm oil, which rose from 0.11 to 0.40. The recorded AnV change was initially high in peanut oil prior to frying but increased more gradually from 0.74 to 1.04 while that of camellia oil rose from 0.17 to 0.55. The IV recorded in peanut oil was the largest though it reduced from 104.74 to 80.52 gI₂/100 g. Insignificant changes of 53.83 to 45.36 gI₂/100 g and 65.40 to 55.29 gI₂/100g were registered in palm and camellia oils respectively. POV in palm oil registered an increment from 4.98 to 18.86 meqO₂/kg while that of peanut and camellia oils changed from 4.75 to 13.24 meqO₂/kg and 4.68 to 11.58 meqO₂/kg. The least AnV was in camellia oil that increased from 1.70 to 51.78 while peanut and palm oils registered 2.25 to 84.71 and 1.36 to 60.00 respectively.

Abdulkarim et al. [9] assessed the frying suitability of high-oleic *Moringa oleifera* seed oil saturated fatty acid (SFA) consisting of SFA: MUFA: PUFA in a ratio of 2:7:0 vis-a-vis soybean, palm olein and canola oils with SFA: MUFA: PUFA ratios of 1.5:2.5:6, 4:4:1 and 1:6:3 respectively. Experimental results showed that

the % FFA of the four conventional oils used comparatively in the assessment respectively increased by 66.6, 71.4, 60.0 and 65.0%. TOTOX and AnV values of the oils were registered in the order *Moringa oleifera* seed oil < palm olein < canola and soybean oils.

This research assessed the effects of continuous deep - fat frying on the physicochemical parameters of edible oils sold in Kampala, Uganda.

2. MATERIALS AND METHODS

2.1 Apparatus and Reagents

The chemicals used in this investigation were all of analytical grade. The conventional volumetric ware employed were sterilized and oven dried prior to analysis. Mettler PM200 balance (Marshall scientific, USA) was used for all weighings. Amprobe IR608A non - contact infrared thermometer with laser pointer, -18°C to 400°C (Amprobe, Everett, USA) was used for monitoring temperatures during frying.

2.2 Sampling Procedure, Sample Size and Sample Preparation

Six (6) 1L samples of hard oil brands (Fortune Butto, Roki, Tamu, Golden fry, Mukwano and Best fry) and soft oil brands (Sunny, Sunvita, Sunlite and Sunseed) of approximate manufacturing dates were procured from Mega Standard Supermarket, Yusuf Lule road, Greater Metropolitan Kampala. The brands were chosen based on their common use in deep frying according to a pre-study tour and were majorly brands from two giant oil refining companies: Bidco Uganda Limited (BUL)-Jinja and Mukwano Industries Limited-Kampala. Hence exactly sixty (60) samples of edible cooking oils were procured and maintained in their original packaging materials under ambient conditions to avoid any possible degradation. Three oil samples used up to ten times continuous frying cycles from Makindye division of greater Metropolitan Kampala were collected at intervals from three randomly selected local traders deep frying Irish potatoes. 10 kg of fresh white (Irish) potatoes (*Solanum tuberosum*) were purchased from Nakasero market, Nakasero Hill, Market square road, Kampala. The potatoes were washed and peeled manually using a clean stainless - steel knife. They were then sliced into cylindrical sizeable pieces (1 cm × 1 cm × 3 cm) corresponding to that used in the Kampala

Metropolis Irish potato culinary. The analyses were done at the Quality Control Laboratory of Mukwano Industries Limited, Plot 30, Mukwano Road, Kampala Industrial area.

2.3 Analysis of Fresh Oil Samples

The physicochemical parameters of the oils were analyzed prior to deep fat frying.

2.4 Frying Methods

Exactly 400 g of Irish potato slices were submersed in 1500mL of oil maintained at 140°C for 6 mins in Skyline VT5424 4 L Electric Deep Fryer (Skyline, New Dehli, India) with detachable oil tank and slotted spoon. A frying time of 10 mins was used with 800 g of the Irish fried in 20 mins. The physicochemical properties of the oils after deep frying were determined after cooling the oils.

2.5 Determination of Colour Index

Color value (color index) of the oil samples free from moisture and insoluble impurities were measured in a Lovibond Tintometer (The Tintometer Ltd, UK) using a 2.54 cm cell operating in the transmittance mode and recorded in Lovibond units.

2.6 Determination of Free Fatty Acids

Free fatty acid was estimated using the AOCS official method Ca 5a - 40 for free fatty acids determination in crude and refined fats and oils [10] with slight modifications. Exactly 10g of the oil sample was weighed in a 250 ml beaker. A measured 60 ml of neutralized ethanol was added and then boiled. The solution was titrated with standardized 0.025 M Sodium hydroxide using phenolphthalein indicator until the solution just turned pink. FFA was expressed as a percentage of oleic acid in the sample.

2.7 Determination of Iodine Absorption Value

Iodine absorption value was determined according to ISO 3961: 2009 iodometric procedure [11]. 0.2 g of the oil sample was weighed into a quick fit flask and dissolved in 20 ml of chloroform. 25 mL of Hanus Iodine solution was added. Three drops of Potassium Iodide were put on the mouth of the flask and kept in the dark for 30 minutes. The sample was then removed and 10 mls of 15% Potassium Iodide

solution was added followed by 100 ml of freshly distilled water. The solution was subsequently titrated with a standard 0.1 N Sodium thiosulphate solution while stirring until a golden yellow color appeared. Exactly 5 mls of starch indicator was added to the resultant solution and titration was continued until the blue-black solution turned colorless. A blank was conducted where the total halogen content of 25 mL of Hanus iodine was determined by a sodium thiosulphate solution without the addition of an oil sample. Iodine value was expressed in grams of iodine absorbed by 100 g of the oil.

2.8 Determination of Peroxide Value

Peroxide value was estimated according to ISO 3960: 2007 [12] and recorded as milliequivalents of active oxygen/kg of the edible cooking oil.

2.9 Determination of Paraanisidine Value

Paraanisidine value is a measure of the carbonyl content of the edible oils. It is based on the reactivity of the aldehyde carbonyl bond on the *p*-anisidine amine group, leading to the formation of a Schiff base that absorbs at 350 nm. Paraanisidine value was determined using the AOCS Official Method Cd 18 - 90 [13] and expressed in anisidine numbers.

2.10 Estimation of Total Oxidation Value

TOTOX value was computed from Equation 1 employed by other authors [8,14,15].

$$TOTOX = 2POV + AnV \quad (1)$$

3. RESULTS

3.1 Statistical Analysis of Results

Each determination was done in triplicate and results recorded as averages or means \pm standard deviations. Data was compared by ANOVA and Tukey test with statistical significance at $P = .05$ using Statgraphics software (Statpoint Technologies, Virginia, USA). The frying stability of the oils during frying and the average number of times the oils could be re-used for continuous deep frying were statistically computed.

3.2 Physicochemical Changes in the Edible Oils

The investigated properties of the oils are given in Table 1, Table 2 and Table 3.

4. DISCUSSION

4.1 Colour Index

There is no standard specification for color as per Uganda National Bureau of Standards (UNBS) for fresh edible oils. However, most Ugandan edible oil refining companies have internal color specifications of 7.5 R 70 Y maximum. Of the fresh oil samples, oil from the third Local chips fryer had the highest color reading of 7.7 R 70 Y (Table 3) while the lowest color reading (0.4 R 3.4 Y) was recorded in Sunseed (Table 2). Sunseed, Sunvita, Sunlite and Sunny cooking oils was still within color index specifications even after the 10th fry (Table 2). Mukwano after the 6th fry, Fortune Butto, Roki and Best fry after the 5th fry, Tamu after the 4th fry and Golden fry after the 3rd fry fell out of specification (Table 1). Oil from the first and second local fryers fell out of color index specification after a fry while oil from the third fryer did not meet the color specification (Table 3). The color values increased significantly ($P = .05$). By the 9th fry, the color recitation for oil from the Third Local fryer could hardly be determined experimentally (Table 3) and by the 10th fry, the oil from the second local fryer had become reasonably dark. Generally, by the 10th fry, all the oils had changed the color of the fried chips. It is worth noting that the yellow units of the color index of all the hard oils did not change during the ten continuous frying cycles.

The physical change in the color value of oils is a rather intuitive and swift visual index implicative of a trend of oil deterioration. Available empirical data shows that unsaturated carbonyl compounds (including ketones, conjugated dienoic acids) and degraded oxidation compounds such as hydroxides and hydroperoxides induce oil darkening [16-18].

Another close cause of the observed color regression could be due to the dispersion of Millard pigments generated *in situ* augmented by traces of carotenoids from the fries [19]. Choe and Min [8] also hinted that polymerized fats accumulated in a fryer during frying causes foam and gum formation as well as oil darkening.

Oil darkening, albeit, an experimentally valuable index while monitoring deterioration of oils heated at elevated temperatures has been underscored to not be solely attributable to oxidative degradation by Che Man and Tan [20]. Non-enzymatic browning of potato chips has been reported to be proportionate to the amount of reducing sugars in the potato, as both browning and Maillard reactions are stimulated by the level of oxidation of the food and the entrained characteristic heme pigments [21]. More so, the Maillard reaction is accompanied by loss of nutrients and the ensuing browning intensity it impacts on the fries is proportionate to the proteinaceous loss of amino acids: lysine, histidine and methionine. The results obtained in this investigation are in complete agreement with the diagnostic statement of Orthoefer and Cooper [22] that assorted frying oils and the foods fried darken oils to varying degrees during deep frying.

4.2 Free Fatty Acids

The chief composition of oils is fatty acids and the degree of unsaturation is the very first factor influencing the oxidative stability of frying oils despite other oil intrinsic and extrinsic factors. In this study, the lowest FFA of 0.0430±0.30 was recorded in Sunvita (Table 2) while the highest recorded FFA of 0.1508±0.30 was in Sunseed (Table 2). Roki, Best fry, Sunvita and oil from

Table 1. Changes in the physicochemical properties of the hard oils

Oil brand	Oil sample	Color value	FFA	POV (meqO ₂ /Kg)	AnV	IV (I ₂ /100g)	TOTOX
Fortune Butto	Fresh	4.8R 70Y	0.0977±0.01	1.6924±0.01	2.50±0.04	57.67±0.06	5.8848
	1 st fry	6.0R 70Y	0.1305±0.02	3.0157±0.02	3.50±0.01	57.56±0.03	9.5314
	2 nd fry	6.3R 70Y	0.1420±0.07	3.0481±0.01	3.90±0.02	56.46±0.01	9.9962
	3 rd fry	7.1R 70Y	0.1428±0.05	3.2682±0.04	4.20±0.02	56.42±0.05	10.736
	4 th fry	7.3R 70Y	0.1522±0.03	3.6635±0.02	4.80±0.01	56.14±0.01	12.127
	5 th fry	7.5R 70Y	0.1779±0.01	6.0672±0.02	5.24±0.11	55.57±0.02	17.374
	6 th fry	8.0R 70Y	0.2544±0.02	8.3062±0.04	7.73±0.03	55.21±0.01	24.342
	7 th fry	8.2R 70Y	0.2613±0.09	9.3708±0.02	8.94±0.06	54.80±0.01	27.682
	8 th fry	8.5R 70Y	0.3269±0.01	9.9978±0.02	10.14±0.01	54.65±0.01	30.136
	9 th fry	9.0R 70Y	0.3513±0.02	10.4800±0.02	10.35±0.01	54.12±0.20	31.310
10 th fry	9.9R 70Y	0.3776±0.02	12.2809±0.02	11.84±0.01	53.66±0.01	36.401	

Oil brand	Oil sample	Color value	FFA	POV (meqO ₂ /Kg)	AnV	IV (I ₂ /100g)	TOTOX	
Roki	Fresh	4.6R	70Y	0.1099±0.05	0.7848±0.001	1.80±0.09	60.06±0.01	3.3696
	1 st fry	6.1R	70Y	0.1305±0.02	1.9309±0.03	2.50±0.05	59.31±0.01	6.3618
	2 nd fry	6.1R	70Y	0.1356±0.01	2.8011±0.02	2.80±0.16	59.15±0.04	8.4022
	3 rd fry	6.5R	70Y	0.1449±0.02	3.0712±0.01	3.40±0.08	58.92±0.07	9.5424
	4 th fry	7.2R	70Y	0.1527±0.02	3.2109±0.01	4.00±0.02	58.92±0.11	10.4218
	5 th fry	7.5R	70Y	0.1541±0.03	5.3638±0.01	4.85±0.01	58.51±0.03	15.5776
	6 th fry	8.2R	70Y	0.1702±0.03	5.7558±0.02	5.38±0.04	57.77±0.06	16.8916
	7 th fry	8.4R	70Y	0.1865±0.01	6.9578±0.03	5.83±0.19	57.77±0.02	19.7456
	8 th fry	9.5R	70Y	0.2098±0.04	9.7719±0.02	6.44±0.01	57.60±0.01	25.9838
	9 th fry	10.0R	70Y	0.2472±0.01	9.9846±0.02	7.47±0.04	57.27±0.05	27.4392
10 th fry	10.8R	70Y	0.2844±0.01	11.1138±0.01	10.31±0.03	57.05±0.01	32.5376	
Tamu	Fresh	5.0R	70Y	0.0987±0.04	0.7137±0.01	2.30±0.20	58.88±0.09	3.7274
	1 st fry	6.3R	70Y	0.1365±0.01	2.0110±0.02	3.80±0.01	58.34±0.07	7.8220
	2 nd fry	6.6R	70Y	0.1443±0.02	4.5741±0.02	4.20±0.11	58.29±0.07	13.3482
	3 rd fry	6.9R	70Y	0.1527±0.02	5.5595±0.02	5.30±0.06	58.05±0.04	16.4190
	4 th fry	7.3R	70Y	0.1812±0.01	7.6355±0.02	6.90±0.01	57.50±0.02	22.1710
	5 th fry	8.5R	70Y	0.1993±0.03	10.0566±0.02	8.94±0.01	57.34±0.01	29.0532
	6 th fry	8.7R	70Y	0.2939±0.02	10.566±0.02	9.70±0.05	57.01±0.03	30.8332
	7 th fry	9.8R	70Y	0.3079±0.04	11.0939±0.01	11.24±0.01	56.96±0.01	33.4278
	8 th fry	10.0R	70Y	0.3053±0.02	11.2413±0.02	11.39±0.11	56.47±0.05	33.8726
	9 th fry	10.8R	70Y	0.3305±0.03	12.5219±0.03	11.59±0.03	56.26±0.21	36.6338
10 th fry	12.0R	70Y	0.3833±0.03	13.0047±0.04	14.73±0.01	55.51±0.10	40.7394	
Golden Fry	Fresh	5.4R	70Y	0.1247±0.02	1.7272±0.11	2.60±0.01	57.62±0.17	6.0544
	1 st fry	6.5R	70Y	0.1462±0.02	4.0312±0.13	3.40±0.31	57.62±0.02	11.4624
	2 nd fry	6.9R	70Y	0.1825±0.06	4.5876±0.01	4.80±0.08	57.57±0.06	13.9752
	3 rd fry	7.2R	70Y	0.2017±0.07	6.0430±0.03	6.20±0.05	57.66±0.03	18.2860
	4 th fry	7.8R	70Y	0.2668±0.12	7.2802±0.02	7.50±0.07	57.40±0.01	22.0604
	5 th fry	8.4R	70Y	0.3029±0.08	8.0133±0.01	8.40±0.03	57.06±0.05	24.4266
	6 th fry	9.1R	70Y	0.3428±0.02	9.3237±0.01	10.80±0.01	56.90±0.01	29.4474
	7 th fry	10.0R	70Y	0.3374±0.06	10.0599±0.06	11.40±0.01	56.45±0.40	31.5198
	8 th fry	11.0R	70Y	0.3874±0.01	10.6783±0.08	11.80±0.02	56.42±0.02	33.1566
	9 th fry	11.8R	70Y	0.4441±0.01	10.9245±0.04	12.40±0.01	56.47±0.06	34.2490
10 th fry	12.4R	70Y	0.4817±0.01	11.7157±0.12	12.80±0.01	56.15±0.01	36.2314	
Mukwano	Fresh	4.5R	70Y	0.0576±0.02	0.7119±0.30	0.90±0.01	61.59 ±0.03	2.3238
	1 st fry	5.8R	70Y	0.1429±0.22	3.9388±0.22	2.90±0.01	61.18±0.04	10.7776
	2 nd fry	6.2R	70Y	0.1581±0.02	4.1649±0.10	3.40±0.01	60.57±0.13	11.7298
	3 rd fry	6.5R	70Y	0.1665±0.02	4.6978±0.12	4.50±0.01	60.20±0.44	13.8956
	4 th fry	6.4R	70Y	0.1755±0.03	6.1839±0.02	6.20±0.01	60.31±0.20	18.5678
	5 th fry	7.0R	70Y	0.2275±0.03	7.4132±0.10	6.80±0.01	59.92±0.01	21.6264
	6 th fry	7.4R	70Y	0.2372±0.02	7.4361±0.02	7.90±0.26	59.22±0.39	22.7722
	7 th fry	8.3R	70Y	0.2534±0.01	9.7599±0.01	8.70±0.03	59.06±0.21	28.2198
	8 th fry	8.9R	70Y	0.2785±0.02	9.9139±0.02	10.20±0.01	58.71±0.01	30.0278
	9 th fry	9.5R	70Y	0.3216±0.01	11.009±0.21	12.69±0.03	58.34±0.04	34.7080
10 th fry	10.9R	70Y	0.3310±0.03	11.820±0.09	14.73±0.03	58.13±0.04	38.3700	
Best fry	Fresh	5.2R	70Y	0.1376±0.01	2.3542±0.02	3.40±0.03	58.34±0.09	8.1084
	1 st fry	7.4	70Y	0.1927±0.01	3.1264±0.07	4.90±0.20	58.16±0.01	11.1528
	2 nd fry	7.9R	70Y	0.2528±0.04	4.2248±0.01	5.80±0.19	57.94±0.01	14.2496
	3 rd fry	8.3R	70Y	0.2655±0.01	4.5703±0.10	7.70±0.26	57.25±0.01	16.8406
	4 th fry	8.8R	70Y	0.2829±0.02	5.8642±0.02	8.40±0.38	57.29±0.08	20.1284
	5 th fry	9.7R	70Y	0.2854±0.12	6.1624±0.13	9.11±0.10	57.13±0.04	21.4348
	6 th fry	10.5R	70Y	0.2978±0.17	6.9323±0.01	12.63±0.01	57.03±0.01	26.4946
	7 th fry	12.0R	70Y	0.2989±0.03	8.8926±0.14	14.21±0.01	57.04±0.01	31.9952
	8 th fry	15.0R	70Y	0.3595±0.12	10.422±0.04	16.12±0.03	56.87±0.01	36.9640
	9 th fry	16.0R	70Y	0.3591±0.02	11.017±0.10	17.03±0.05	56.71±0.01	39.0640
10 th fry	17.5R	70Y	0.3591±0.03	11.952±0.08	17.77±0.20	56.66±0.17	41.6740	

Table 2. Changes in the physicochemical properties of the soft oils

Oil brand	Oil sample	Color value	FFA	POV (meqO ₂ /Kg)	AnV	IV (I ₂ /100g)	TOTOX
Sunseed	Fresh	0.4R 3.4Y	0.1508±0.03	6.6134±0.23	4.30±0.19	58.30±0.02	17.5268
	1 st fry	1.0R 4.7Y	0.1567±0.07	7.8224±0.32	5.70±0.01	58.02±0.02	21.3448
	2 nd fry	1.1R 5.4Y	0.1693±0.01	8.8646±0.08	6.60±0.20	58.08±0.07	24.3292
	3 rd fry	1.2R 6.3Y	0.1829±0.02	9.2997±0.30	7.40±0.03	57.88±0.01	25.9994
	4 th fry	1.3R 7.2Y	0.1932±0.22	10.2907±0.01	9.50±0.08	57.87±0.42	30.0814
	5 th fry	2.0R 8.5Y	0.2099±0.01	10.8085±0.29	10.90±0.10	57.83±0.37	32.5170
	6 th fry	2.1R 9.9Y	0.2152±0.33	10.9622±0.10	13.80±0.19	57.71±0.22	35.7244
	7 th fry	2.1R 11Y	0.2201±0.01	10.9920±0.23	16.70±0.02	57.72±0.19	38.6840
	8 th fry	2.4R 13Y	0.2715±0.01	12.2302±0.02	17.90±0.56	57.57±0.01	42.3604
	9 th fry	2.9R 14Y	0.2769±0.02	13.4418±0.12	19.50±0.49	57.45±0.02	46.3836
10 th fry	2.6R 22Y	0.2908±0.02	13.6042±0.02	20.20±0.30	57.40±0.08	47.4084	
Sunny	Fresh	0.8R 4.9Y	0.0884±0.31	1.4980±0.06	2.50±0.06	128.35±0.02	5.4960
	1 st fry	1.6R 14Y	0.1163±0.50	2.1135±0.07	3.70±0.01	127.13±0.05	7.9270
	2 nd fry	1.9R 15Y	0.1407±0.09	3.0874±0.02	4.00±0.02	126.94±0.07	10.1748
	3 rd fry	2.1R 19Y	0.1571±0.04	4.3946±0.01	4.90±0.10	126.88±0.10	13.6892
	4 th fry	2.7R 16Y	0.1766±0.06	5.0359±0.06	6.20±0.10	126.70±0.43	16.2718
	5 th fry	2.9R 19Y	0.1875±0.06	6.7514±0.01	7.60±0.01	126.69±0.01	21.1028
	6 th fry	3.0R 23Y	0.1917±0.10	7.9430±0.02	9.40±0.11	126.33±0.56	25.2860
	7 th fry	3.0R 24Y	0.2173±0.01	9.0678±0.02	10.40±0.08	126.27±0.34	28.5356
	8 th fry	3.3R 24Y	0.2435±0.09	10.3814±0.17	12.80±0.04	126.24±0.01	33.5628
	9 th fry	3.5R 27Y	0.2477±0.04	10.8668±0.10	15.50±0.02	126.08±0.31	34.5336
10 th fry	3.6R 27Y	0.2540±0.11	11.7032±0.04	17.50±0.01	126.03±0.02	40.9064	
Sunvita	Fresh	0.9R 5.1Y	0.0430±0.30	1.4714±0.01	2.80±0.05	126.96±0.11	5.7428
	1 st fry	0.9R 6.7Y	0.0892±0.04	3.3087±0.01	3.50±0.02	126.10±0.28	10.1174
	2 nd fry	0.9R 7.5Y	0.1503±0.08	3.8037±0.01	4.70±0.09	123.03±0.21	12.3074
	3 rd fry	1R 9.1Y	0.1666±0.01	4.9809±0.08	5.70±0.12	124.88±0.52	15.6618
	4 th fry	1.1R 11Y	0.1937±0.14	5.2830±0.11	6.80±0.03	123.96±0.06	17.3660
	5 th fry	1.3R 11Y	0.1991±0.01	5.9740±0.05	7.75±0.02	123.27±0.11	19.6980
	6 th fry	1.7R 14Y	0.2136±0.03	8.2301±0.02	10.46±0.19	122.59±0.42	26.9202
	7 th fry	2.1R 15Y	0.2576±0.07	10.110±0.03	14.89±0.01	121.81±0.01	35.1100
	8 th fry	2.4R 20Y	0.3081±0.02	11.530±0.20	15.78±0.02	121.48±0.05	38.8400
	9 th fry	2.8R 22Y	0.3078±0.05	14.680±0.16	16.49±0.02	121.48±0.04	45.8500
10 th fry	3.0R 23Y	0.3396±0.02	15.370±0.31	18.66±0.02	120.28±0.01	49.4000	
Sunlite	Fresh	0.7R 5Y	0.0963±0.01	1.2723±0.01	2.60±0.01	126.80±0.22	5.1446
	1 st fry	0.7R 5.5Y	0.1098±0.06	2.0882±0.12	2.92±0.15	126.46±0.15	7.0964
	2 nd fry	0.8R 5.8Y	0.1319±0.01	3.9624±0.04	3.50±0.01	125.98±0.02	11.4248
	3 rd fry	1.0R 6.0Y	0.1458±0.03	4.4835±0.20	4.30±0.01	125.79±0.20	13.2670
	4 th fry	1.3R 6.8Y	0.1603±0.01	5.9978±0.01	5.80±0.06	125.73±0.40	17.7956
	5 th fry	1.5R 6.8Y	0.1720±0.01	7.5574±0.01	6.90±0.11	125.47±0.25	22.0148
	6 th fry	1.8R 7.2Y	0.2001±0.05	9.0822±0.03	8.30±0.20	125.29±0.09	26.4644
	7 th fry	2.0R 7.5Y	0.2159±0.06	10.5798±0.16	10.0±0.01	124.89±0.04	31.1596
	8 th fry	2.4R 13.0Y	0.2599±0.02	11.2972±0.08	12.72±0.04	124.57±0.21	35.3144
	9 th fry	2.56R 13Y	0.2828±0.02	11.9001±0.20	14.68±0.02	124.55±0.13	38.4802
10 th fry	2.9R14.5Y	0.3202±0.03	12.7721±0.10	16.02±0.02	124.21±0.25	41.5642	

the Second Local fryer were still within the maximum permissible FFA of 0.30 for edible oils even after the 10th fry (Table 1, Table 2, Table 3). Sunlite after the 9th fry, Mukwano after the 8th fry, Fortune Butto, Sunny and oil from the First Local fryer after the 7th fry had FFA greater than the permissible limit (Table 1, Table 2, Table 3). Tamu, Sunseed and oil from the Third Local fryer

after the 6th fry and Golden fry after the 4th fry had FFA greater than the specification of 0.30 (Table 1, Table 2, Table 3).

The differences observed in the FFA values of the fresh edible oils could be attributed to blending of several edible oils by the refining companies. Blending of edible oils is known to

Table 3. Changes in the physicochemical properties of oils from the local fryers

Oilbrand	Oil sample	Colorvalue	FFA	POV (meqO ₂ /Kg)	AnV	IV (I ₂ /100g)	TOTOX
First Local Fryer	Fresh	6.8R 70Y	0.1399±0.02	3.1037±0.10	4.30±0.01	58.42±0.22	10.5074
	1 st fry	7.4R 70Y	0.1499±0.04	3.5849±0.06	5.10±0.11	58.42±0.92	12.2698
	2 nd fry	7.9R 70Y	0.1556±0.01	4.4937±0.01	6.80±0.02	58.38±0.34	15.7874
	3 rd fry	8.1R 70Y	0.1698±0.02	5.8645±0.01	7.50±0.02	58.30±0.01	19.2290
	4 th fry	8.9R 70Y	0.2013±0.07	8.6028±0.04	7.90±0.02	58.34±0.20	25.1056
	5 th fry	9.7R 70Y	0.2432±0.01	9.0689±0.01	9.00±0.02	58.15±0.05	27.1378
	6 th fry	10.7R 70Y	0.2449±0.01	10.738±0.11	10.40±0.06	57.67±0.02	31.876
	7 th fry	12.0R 70Y	0.2466±0.03	11.0264±0.01	14.20±0.04	57.66±0.31	36.2528
	8 th fry	12.9R 70Y	0.2635±0.06	12.2278±0.01	16.50±0.01	57.55±0.02	40.9556
	9 th fry	14.8R 70Y	0.3094±0.01	14.6568±0.03	17.50±0.03	57.55±0.01	46.8136
10 th fry	15.2R 70Y	0.3334±0.03	15.7525±0.01	19.90±0.01	57.35±0.02	51.405	
Second Local Fryer	Fresh	4.5R 70Y	0.1007±0.01	0.5951±0.03	2.80±0.01	59.43±0.17	3.9902
	1 st fry	6.0R 70Y	0.1282±0.01	0.9021±0.02	3.40±0.02	59.23±0.07	5.2042
	2 nd fry	6.0R 70Y	0.1300±0.05	1.5840±0.01	3.80±0.01	58.90±0.01	6.9680
	3 rd fry	6.2R 70Y	0.1354±0.01	2.2182±0.02	4.50±0.03	58.97±0.32	8.9364
	4 th fry	7.1R 70Y	0.1398±0.02	2.9961±0.01	5.10±0.02	58.55±0.21	11.0922
	5 th fry	7.3R 70Y	0.1403±0.03	3.3954±0.01	6.30±0.06	58.37±0.40	13.0908
	6 th fry	7.8R 70Y	0.1625±0.01	4.4962±0.01	6.50±0.01	58.27±0.07	15.4924
	7 th fry	8.2R 70Y	0.1687±0.03	5.1984±0.05	7.40±0.01	58.15±0.03	17.7968
	8 th fry	8.7R 70Y	0.1812±0.04	6.5811±0.01	8.50±0.04	58.10±0.01	21.6622
	9 th fry	9.0R 70Y	0.2014±0.01	6.9981±0.02	9.10±0.01	58.10±0.63	23.0962
10 th fry	9.5R 70Y	0.2286±0.01	7.995±0.12	10.70±0.22	58.12±0.18	26.6900	
Third Local Fryer	Fresh	7.7R 70Y	0.0496±0.01	1.1957±0.01	2.70±0.04	128.16±0.22	5.0914
	1 st fry	9.0R 70Y	0.1362±0.02	1.6099±0.01	3.60±0.01	127.18±0.01	6.8198
	2 nd fry	10.1R 70Y	0.1535±0.01	1.7636±0.01	3.90±0.02	126.33±0.21	7.4272
	3 rd fry	10.9R 70Y	0.1634±0.01	2.0909±0.01	4.80±0.02	124.79±0.17	8.9818
	4 th fry	12.5R 70Y	0.1697±0.03	2.5227±0.01	5.90±0.07	124.22±0.01	10.9454
	5 th fry	14.3R 70Y	0.2030±0.01	2.4450±0.02	7.46±0.01	123.30±0.03	12.3500
	6 th fry	16.0R 70Y	0.2980±0.02	2.9739±0.06	8.65±0.02	121.40±0.04	14.5978
	7 th fry	17.0R 70Y	0.3235±0.04	6.0875±0.01	14.20±0.02	120.13±0.11	26.3750
	8 th fry	18.8R 70Y	0.3874±0.01	6.3546±0.03	17.98±0.16	118.77±0.05	30.6892
	9 th fry	19.2R 70Y	0.4165±0.01	7.1760±0.01	20.70±0.03	119.00±0.01	35.0520
10 th fry	20.4R 70Y	0.4748±0.06	8.5800±0.04	22.16±0.01	119.14±0.02	39.3200	

alter their fatty acid profiles [23,24] and can steeply retard oxidation of oils during deep-fat frying. The increase in FFA values of oils during the frying cycles can be attributed to the breakdown of long carbon chains into shorter chains due to thermal and oxidative decomposition. During elevated temperature heating of oils, FFA formation is attributed to the cleavage and oxidation of double bonds to form carbonyl compounds, which are subsequently oxidized to fatty acids of low molecular masses [25-27]. FFA is preferred frequently by food processors for indication of oil acidity and oil authenticity verification [28,29]. Filtration of frying oils have been reported to reduce FFA content of oils and improve their frying stability. The results of this collaborative study is concordant with that of Stevenson et al. [30] who reported that edible oils with FFA values less than 0.05 and POV of 1.0 meq O₂/kg or less are best suited for deep frying. It also agrees well with other studies that the FFA content of oils increases with the number of frying cycles [31] as well as the frying time [32].

4.3 Peroxide Value

For fresh samples, oil from the Second Local fryer had the lowest POV of 0.5951±0.03 meq O₂/Kg (Table 3) while the highest POV of 6.6134±0.23 meq O₂/Kg was observed in Sunseed (Table 2). Aside from the nature of the oil, oils to be utilized in deep frying should have a Codex regulatory maximum POV of 15 meq O₂/Kg [33]. In this study, POVs increased significantly ($P = .05$) during the ten successive fries (Table 1, Table 2, Table 3). However, up to the 10th fry, the POVs of all the oils were still concordant with the maximum Codex standard POV for vegetable oils except for Sunvita and Oil from the First Local Fryer that were above the maximum Codex standard after the 9th fry (Table 2 and Table 3).

The increase in the POVs of edible oils following frying is because of the oxidation of carbon atoms adjacent to the double bonds in the triacyl glyceride structure leading to the formation of hydroperoxides. These hydroperoxides are the direct cause of AnV shoot up as they decompose further to secondary oxidative compounds which constitute the paraanisidine components. POV is implicative of incipient oxidation which directly translates into the buildup and breakdown of oxidation products. Peroxides are reasonably unstable, and fissions at typical frying temperatures. In addition, it is a useful biomarker

of the preliminary stages of rancidity occurring under mild conditions and the freshness of the lipid matrix. Thus, the greater the POV, the faster will the oxidation of the oil occur [34].

The observed increase in POV during heating of oils have been reported by other authors [8,28, 35,36]. Furthermore, polyunsaturated oils exhibit readily depressed stability at elevated temperatures because the unsaturated fatty acids are readily oxidized to peroxides [36,37]. Empirical data also point that POVs of fresh oils may be higher than the Codex standard primarily owing to improper storage and/or packaging that triggers degradation via photo-oxidation. Auto-oxidation of oils may supposedly occur in storage due to chemical interaction with air, peculiarly oxygen [38]. However, such oils with high POVs are typically unstable and becomes rancid easily [39].

4.4 Paraanisidine Value

Oxidative degradation of frying oils is innocuously deleterious as it impacts sensorial and organoleptic attributes of fries [40]. Primary oxidation quantifies the amount of hydroperoxides as POV. Further degradation of hydroperoxides yields aldehydes, ketones, carboxylic acids, short chain alkanes and alkenes better quantitatively described by paraanisidine value (AnV).

For fresh oils, the lowest AnV of 0.90±0.01 was observed in Mukwano (Table 1) while the highest recorded AnV were 4.30±0.01 and 4.30±0.19 in Sunseed and oil from the First Local fryer (Table 2 and Table 3) respectively. Sunseed after the 4th fry, Golden fry, Best fry, Oil from the First Local Fryer and Sunvita after the 5th fry were above the maximum permissible limit of 10 [41]. Tamu, Sunny and Oil from the Third Local fryer after the 6th fry were out of specifications. Sunlite, Fortune Butto and Mukwano were out of specifications after the 7th fry whereas Roki and oil from the Second Local fryer after the 9th fry recorded AnV above the maximum acceptable limit.

Secondary oxidation products are principally non-volatile aldehydes, principally 2,4-dienals and 2-alkenals [42,43] which AnV is a quantitative measure. The AnV was observed to increase insignificantly ($P = .05$) between the successive fries. Initial stages of heating resulted in faster increase in AnV followed by a more gradual increment (Table 1, Table 2 and Table

3). This could be due to further decomposition of the carbonyls and polymerization reactions. The observed trend is due to the fact that the less stable primary oxidative products (hydroperoxides) decompose further to form aldehydic compounds. These compounds are not easily decomposed, thus the test to determine the AnV is more meaningful than that to determine the POV. Similar results have been reported by Xu et al. [8] in their comparison of oxidative stability of edible oils under continuous deep frying conditions.

4.5 Iodine Absorption Value

The iodine adsorption value, iodine number or sometimes iodine index, is chemically the mass of iodine in grams that is consumed by 100 grams of a chemical substance by mass as oleic acid. Iodine numbers are often used to determine the amount of unsaturation in fatty acids. The higher the iodine index (the greater the unsaturation), the faster is the tendency of oil oxidation during heating at elevated temperatures as in deep frying [44]. Iodine index, is a frequently considered vital analytical measure of the unsaturation of oils [45] and any significant differences in IV can be translated into increased oxidation rate.

At fresh conditions, the maximum IV observed was in Sunny ($128.35 \pm 0.02 \text{ gl}_2 / 100 \text{ g}$) (Table 2) and the least IV of $57.62 \pm 0.17 \text{ gl}_2 / 100 \text{ g}$ was recorded in Golden fry (Table 1). It was observed in this investigation that there were no significant differences ($P = .05$) in IV during the frying cycles. The observed decrement in the iodine indices is concordant with the decrement in double bonds attributed to oxidation and thermal decomposition and has been reported elsewhere [8]. However, this decrement implies there was little oxidation because significant changes in iodine values can only be appreciated during deep frying when there is excessive deterioration of the oil [17].

4.6 Frying Stability of the Edible Oils

The average number of times the hard oils and soft oils could be re-used for continuous deep-fat frying were statistically calculated. Possible reuse of the oils were estimated from the frying round when a quality criterion (POV or AnV) surpassed UNBS or Codex Alimentarius specification. The analysis gave an average of 6.5 times (≈ 7 times for hard oils and 5.5 times (≈ 6 times) for soft oils.

The stability (oxidizability) of the oils were reflected by the changes in their TOTOX values during the frying cycles. The TOTOX values of frying oils take into consideration the actual state of the oils (using POV) as well as the history of the oil (employing the AnV). Thus, the significance of TOTOX value is much stronger than the individual Peroxide and Anisidine values. A 2006 position paper for the German Society for fat sciences (cited in [45]) recommended a TOTOX value of less than 20 for refined and virgin vegetable oils. In this study, a maximum TOTOX value of 20 was considered acceptable for deep frying.

Among the hard oil brands investigated, Roki was the most stable followed by Fortune Butto, Mukwano, Tamu, Best fry and lastly Golden fry (Table 1). For soft oils, Sunseed was the most unstable as it could only be used for frying once (Table 2). It was followed by Sunny, Sunlite and lastly Sunvita (Table 2). Oils from the Local fryers followed the order: Oil from the Second Fryer > Oil from the Third Fryer > Oil from the First Local Fryer. Overall, the stability of the oils investigated followed the order: Oil from the Second Local Fryer > Roki > Oil from the Third Local fryer > Fortune Butto > Sunvita > Sunny > Sunlite > Mukwano > Tamu > Best Fry > Golden Fry > Oil from the First Local Fryer > Sunseed.

5. CONCLUSIONS AND RECOMMENDATIONS

Prior to deep frying, all the selected brands of edible oils sold in Kampala Metropolis met Codex Alimentarius specifications in terms of the assessed properties. The statistical parameter ranges were: CV (0.4 R 3.4 Y - 7.7 R 70 Y), FFA ($0.0430 \pm 0.30 - 0.1508 \pm 0.30$), POV ($0.5951 \pm 0.03 - 6.6134 \pm 0.23 \text{ meqO}_2/\text{Kg}$), AnV ($0.90 \pm 0.01 - 4.30 \pm 0.19$) and IV ($57.62 \pm 0.17 - 128.35 \pm 0.02 \text{ gl}_2 / 100 \text{ g}$). All the physicochemical parameters of the oils increased with the number of frying cycles except IV that decreased slightly. By the 10th fry, the statistical parameter ranges were CV (3.0 R 23 Y - 20.4 R 70 Y), FFA ($0.2286 \pm 0.01 - 0.4817 \pm 0.01$), POV ($11.1138 \pm 0.01 - 15.7525 \pm 0.01 \text{ meqO}_2/\text{Kg}$), AnV ($10.31 \pm 0.03 - 22.16 \pm 0.01$) and IV ($53.66 \pm 0.01 - 126.03 \pm 0.02 \text{ gl}_2 / 100 \text{ g}$). Repeated re-use of the oils for consecutive deep frying of Irish potato chips on the same day can be done only up to a maximum of 7 times on average for the investigated hard oils and 6 times for the soft oils with the oils still regarded as safe for frying potato chips for human consumption. Thus, hard oils should be

preferred to soft oils for deep frying of Irish potato chips.

Further comprehensive research should elucidate the variation of physicochemical properties of other edible cooking oil brands on the Ugandan market such as Nile, Fortune, Kimbo, Star Fry, Cow boy and Ufuta. Further research should be done with other food samples such as fish, cassava, chicken, sweet plantain, dough, meat and edible grasshoppers as the nature of the food sample influences the quality of an oil utilized for deep frying. The organoleptic test of smell should not be used physiognomically in making conclusions about the suitability of cooking oils after deep fat frying as it is hard to clearly deduce since individual sensorium vary widely and thus judgement is made so indifferently. Other physicochemical properties of the investigated edible cooking oils such as smoke point, viscosity, moisture content, volatile matter content, total polar components and saponification value should be determined. Waste oils not recommended for further use for deep frying of Irish potatoes could be harnessed for biodiesel production.

ACKNOWLEDGEMENTS

We appreciate the analytical services of Mukwano Industries Limited-Kampala. We owe a lot to Ngarombo John Peter (RIP), the standards inspector of UNBS for his technical support during the research.

COMPETING INTERESTS

Authors have declared that no competing interests exist. The products used in this research are common and predominantly used products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because the authors do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing companies rather it was funded by personal efforts of the authors.

REFERENCES

1. Omara T, Nnankabirwa R, Nassazi W, Nakabuye BV, Musau B. Spectroscopic screening of assorted pigmented vegetables and fruits common in metropolitan kampala culinary recipes for anthocyanins. *Int J Food Sci Nutr.* 2018; 3(6):285-290. Available:<http://www.foodsciencejournal.com/download/506/3-6-62-643.pdf>
2. Peers KE, Swoboda PAT. Deterioration of sunflower seed oil under simulated frying conditions and during small-scale frying of potato chips. *J Sci Food Agric.* 1982;33: 389–95. Available:<https://doi.org/10.1002/jsfa.2740330414>
3. Cuesta C, Sanchez-Muniz FJ, Garrido-Polonio C, Lopez-Varela S, Arroyo R. Thermooxidative and hydrolytic changes in sunflower oil used in frying with a fast turnover of fresh oil. *J Amer Oil Chem Soc.* 1993;70:1069–73.
4. Sanchez-Muniz FJ, Cuesta C, Lopez-Varela MC, Garrido-Polonio MC, Arroyo R. Evaluation of the thermal oxidation rate of sunflower oil using various frying Methods. In: Applewhite TH, editor. *Proceedings of World conference on oilseed and technology and utilization.* Champaign, Ill: American Oil Chemists Society; 1993.
5. Houhoula DP, Oreopoulou V, Tzia C. The effect of process time and temperature on the accumulation of polar compounds in cottonseed oil during deep-fat frying. *J Sci Food Agric.* 2003;83(4),314–19. Available:<https://doi.org/10.1002/jsfa.1314>
6. Baby Latha R, Nasirullah. Physico-chemical changes in rice bran oil during heating at frying temperature. *J Food Sci Tech.* 2014;51:335-40. Available:<https://doi.org/10.1007/s13197-011-0495-9>
7. Diop A, Ndao S, Cissé M, Baldé M, Ndiaye B, Diop YM. Effect of deep-fat frying on chemical properties of edible vegetable oils used by senegalese households. *Afr J Food Agric Nutr Dev.* 2014;14(6):9418-38. Available:<https://www.ajol.info/index.php/ajfand/article/view/109767/99516>
8. Xu T, Li J, Fan Y, Zheng T, Deng Z. Comparison of oxidative stability among edible oils under continuous frying conditions. *Int J Food Prop.* 2015;18: 1478–90. Available:<https://www.tandfonline.com/doi/ci-edby/10.1080/10942912.2014.913181?scroll=top&needAccess=true>
9. Abdulkarim SM, Long K, Lai OM, Muhammad SKS, Ghazali HM. Frying quality and stability of high-oleic *Moringa oleifera* seed oil in comparison with other

- vegetable oils. Food Chem. 2007;105: 1382–89.
Available:<https://doi.org/10.1016/j.foodchem.2007.05.013>
10. AOCS official method Ca 5a-40. Free fatty acids. American Oil Chemists' Society; 2009.
 11. ISO. Animal and Vegetable Fats and Oils-Determination of Iodine Value. ISO 3961: 2009. 4th Edn. 2009, Geneva, Switzerland.
 12. ISO. Animal and vegetable fats and oils-determination of peroxide value. ISO 3960: 2007. 2007, Geneva, Switzerland.
 13. AOCS. AOCS. *p*-Anisidine value. In: AOCS official method Cd 18-90: Official methods and recommended practices of the American oil chemists' society. Champaign, IL, USA; 1998.
 14. Shahidi F, Wanasundara UN. Methods for measuring oxidative rancidity in fats and oils. In: Food lipids: Chemistry, Nutrition, and Biotechnology; Akoh CC, Min DB, Eds. Marcel Dekker Inc: New York, 2002; 465–482.
 15. Pereira de Abreu DA, Paseiro Losada P, Maroto J, Cruz JM. Evaluation of the effectiveness of a new active packaging film containing natural antioxidants (From Barley Husks) that retard lipid damage in frozen Atlantic Salmon (*Salmo salar* L.). Food Res Int. 2010;43(5):1277–82. Available:<http://dx.doi.org/10.1016/j.foodres.2010.03.019>
 16. Gutierrez R, Gonzalez-Quijano J and Dobermans MC. Analytical procedures for the evaluation of used frying fats. In: Varela G, Bender AE, Morton ID (Eds.), Frying of food: Principles, changes, new approaches. England: VCH-Ellis Horwood Ltd. 141-154.
 17. Augustin MA, Berry SK. Efficacy of the antioxidants BHA and BHT in palm olein during heating and frying. J Amer Oil Chem Soc. 1983;60:1520-23.
 18. Farhoosh R, Kenari RE, Pooazrang R. Frying stability of canola oil blended with palm olein, olive and corn oils. J Amer Oil Chem Soc. 2009;86(1):71-76. Available:<https://doi.org/10.1007/s11746-008-1315-x>.
 19. Lalas S, Gortzi O, Tsankanis J. Frying stability of *Moringa stenopetala* seed oil. Plant Foods Hum Nutr. 2006;61(2):99-108. Available:<https://doi.org/10.1007/s11130-006-0022-8>
 20. Che Man YB, Tan CP. Effects of natural antioxidants on changes in refined, Bleached and deodorized palm olein during deep-fat frying of potato chips. J Amer Oil Chem Soc. 1999;76:331-39. Available:<https://doi.org/10.1007/s11746-999-0240-y>.
 21. Reda SY. Comparative study of vegetable oils subjected to thermal stress. Masters dissertation. Universidade Estadual de Ponta Grossa; 2000.
 22. Orthoefer ET, Cooper DS. Evaluation of used frying oil. In: Perkins EG, Erickson MD. Deep frying: Chemistry, Nutrition and practical applications. AOAC Press: Champaign.1996;285-296.
 23. Shiota M, Konishi H, Tatsumi K. Oxidative stability of fish oil blended with butter. J Dairy Sci. 1999;82:1877–81. Available:[https://doi.org/10.3168/jds.S0022-0302\(99\)75421-4](https://doi.org/10.3168/jds.S0022-0302(99)75421-4)
 24. Mamat H, Aini IN, Said M, Jamaludin R. Physicochemical characteristics of palm oil and sunflower oil blends fractionated at different temperatures. Food Chem. 2005; 91:731–36. Available:<https://doi.org/10.1016/j.foodchem.2004.06.045>
 25. Baby Latha R, Nasirullah. Effect of heat on physico-chemical and thermo-oxidative stability of repeatedly heated rice bran oil (RBO). Int J Food Nutr Sci. 2016;5: 49-60. Available:https://www.ijfans.com/ijfansadmin/upload/ijfans_572329985c27f.pdf
 26. Faridah DN, Lioe HN, Palupi NS, Kahfi J. Detection of FFA and peroxide values using FTIR for quality measurement in palm oil frying activities. J Oil Palm Res. 2015;27(2):156-67. Available:https://inis.iaea.org/search/search.aspx?orig_q=RN:48005572
 27. Lalas S. Quality of frying oil. Advances in deep-fat frying of foods. New York, USA: CRC Press; 2009.
 28. Park J, Kim J. Monitoring of used frying oils and frying times for frying chicken nuggets using POV and acid value. Korean J Food Sci Ani. 2016;5(36):612–16. Available:<https://dx.doi.org/10.5851%2Fkofs.2016.36.5.612>.
 29. Ahmad-Tarmizi AH, Siew WL. Quality assessment of palm products upon prolonged heat treatment. J Oleo Sci. 2008;57(12):639-48. Available:<https://doi.org/10.5650/jos.57.639>
 - Stevenson SG, Vaisey-Genser M, Eskin NAM. Quality control in the use of deep

- frying oils. J Amer Oil Chem Soc. 1984; 61:1102–08.
Available:<https://doi.org/10.1007/BF02636232>
30. Codex alimentarius. Codex standard for named vegetable oils, Codex stan 210-1999. Amended in 2005 and 2011. Rome, Italy; 1999.
31. Chung J, Lee J, Choe E. Oxidative stability of soybean and sesame oil mixture during frying of flour dough. J Food Sci. 2004; 69:574-78.
Available:<https://doi.org/10.1111/j.1365-2621.2004.tb13652.x>
32. Mazza G, Qi H. Effect of after-cooking darkening inhibitors on stability of frying Oil and quality of french fries. J Amer Oil Chem Soc. 1992;69:847–53.
Available:<https://doi.org/10.1007/BF02636331>
33. Atinafu DG, Bedemo B. Estimation of total free fatty acid and cholesterol content in some commercial edible oils in Ethiopia, Bahir DAR. J Cereals Oilseeds. 2001;2:71-76.
Available:<http://www.academicjournals.org/JCO>
34. Sulthana SN, Sen DP. Studies on deep Fat frying: Changes during heating of oil. J Food Sci Tech. 1979;16:208.
35. Serjouie A, Tan CP, Mirhosseini H, Che Man YB. Effect of vegetable-based oil blends on physicochemical properties of oils during deep-fat frying. Amer J Food Tech. 2010;5(5):310-23.
Available:<http://dx.doi.org/10.3923/ajft.2010.310.323>
36. Che-Man YB, Wan-Hussin WR. Comparison of the frying performance of refined, bleached and deodorized palm olein and coconut oil. J Food Lipids. 1998; 5:197–210.
Available:<https://doi.org/10.1111/j.1745-4522.1998.tb00120.x>
37. Ojeh O. Effects of refining on the physical and chemical properties of cashew kernel oil. J Fats Oils Tech. 1981;16(5):513–17.
Available:<https://doi.org/10.1111/j.1365-2621.1981.tb01844.x>
38. Susheelamma NS, Asha MR, Ravi R, Kumar AKV. Comparative studies on physical properties of vegetable oils and their blends after frying. J Food Lipids. 2002;9:259–76.
Available:<https://doi.org/10.1111/j.1745-4522.2002.tb00225.x>
39. Dana D, Saguy IS. Frying of nutritious foods: Obstacles and feasibility. Food Sci Tech Res. 2001;7(4):265-79.
Available:<https://doi.org/10.3136/fstr.7.265>
40. Rossell JB. Measurement in rancidity. In: Allen JC, Hamilton RJ (eds) rancidity in food. Elsevier, London. 1989;22–51.
41. Sulieman MR, Makhzangy AE, Ramadan MF. Antiradical performance and physicochemical characteristics of vegetable oils upon frying of French fries: A preliminary comparative study. J Food Lipids. 2006;13(3):259-76.
Available:<https://doi.org/10.1111/j.1745-4522.2006.00050.x>
42. Bou R, Navas JA, Tres A, Codony R, Guardiola F. Quality assessment of frying fats and fried snacks during continuous deep-fat frying at different large-scale producers. Food Control. 2012;27(1):254–67.
Available:<https://doi.org/10.1111/j.1745-4522.2006.00050.x>
43. Tomkins PC, Perkins EG. The evaluation of frying oils with the P-anisidine value. J Amer Oil Chem Soc. 1999, 76 (8), 945-947.
Available:<https://doi.org/10.1007/s11746-999-0111-6>
44. Otunola AG, Adebayo GB, Olufemi OG. Evaluation of some physicochemical parameters of selected brands of vegetable oils sold in ilorin metropolis. Int J Phys Sci. 2009;4(5):327-29.
45. Marthias B. In: Oxidation in foods and beverages and antioxidant applications: Management in different industry sectors; 2010.

© 2019 Omara et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48900>